

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification<sup>5</sup> : C07H 1/00, A61K 31/70 C12Q 1/68</p>	<p>A1</p>	<p>(11) International Publication Number: WO 91/10671 (43) International Publication Date: 25 July 1991 (25.07.91)</p>														
<table border="0"> <tr> <td data-bbox="183 447 784 468"> <p>(21) International Application Number: PCT/US91/00243</p> </td> <td data-bbox="784 447 1395 468"></td> </tr> <tr> <td data-bbox="183 489 784 510"> <p>(22) International Filing Date: 11 January 1991 (11.01.91)</p> </td> <td data-bbox="784 489 1395 510"></td> </tr> <tr> <td data-bbox="183 531 784 594"> <p>(30) Priority data: 463,358 11 January 1990 (11.01.90) US 566,977 13 August 1990 (13.08.90) US</p> </td> <td data-bbox="784 531 1395 594"> <p>(74) Agent: CALDWELL, John, W.; Woodcock Washburn Kurtz Mackiewicz &amp; Norris, One Liberty Place - 46th Floor, Philadelphia, PA 19103 (US).</p> </td> </tr> <tr> <td data-bbox="183 615 784 741"> <p>(60) Parent Applications or Grants (63) Related by Continuation US 566,977 (CIP) Filed on 11 January 1990 (11.01.90) US 463,358 (CIP) Filed on 11 January 1990 (11.01.90)</p> </td> <td data-bbox="784 615 1395 741"> <p>(81) Designated States: AT (European patent), AU, BE (European patent), BR, CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GB (European patent), GR (European patent), HU, IT (European patent), JP, KR, LU (European patent), NL (European patent), NO, SE (European patent), US.</p> </td> </tr> <tr> <td data-bbox="183 762 784 825"> <p>(71) Applicant (for all designated States except US): ISIS PHARMACEUTICALS, INC. [US/US]; 2280 Faraday Avenue, Carlsbad, CA 92008 (US).</p> </td> <td data-bbox="784 762 1395 825"> <p>Published With international search report.</p> </td> </tr> <tr> <td data-bbox="183 846 784 867"> <p>(72) Inventors; and</p> </td> <td data-bbox="784 846 1395 867"></td> </tr> <tr> <td data-bbox="183 867 784 1119"> <p>(75) Inventors/Applicants (for US only) : COOK, Philip, Dan [US/US]; 6430 La Paloma Street, Carlsbad, CA 92009 (US). ECKER, David, J. [US/US]; 2609 Colibri Lane, Carlsbad, CA 92009 (US). GUINOSSO, Charles, John [US/US]; 313 Encino Drive, Vista, CA 92083 (US). ACEVEDO, Oscar, Leobardo [US/US]; 12117 Brickellia Street, San Diego, CA 92129 (US). KAWASAKI, Andrew, Mamoto [US/US]; 2395 Rancho Del Oro Road, #13, Oceanside, CA 92056 (US). RAMASAMY, Kandasamy [US/US]; 5 Rocky Creek Lane, Laguna Hills, CA 92653 (US).</p> </td> <td data-bbox="784 867 1395 1119"></td> </tr> </table>			<p>(21) International Application Number: PCT/US91/00243</p>		<p>(22) International Filing Date: 11 January 1991 (11.01.91)</p>		<p>(30) Priority data: 463,358 11 January 1990 (11.01.90) US 566,977 13 August 1990 (13.08.90) US</p>	<p>(74) Agent: CALDWELL, John, W.; Woodcock Washburn Kurtz Mackiewicz &amp; Norris, One Liberty Place - 46th Floor, Philadelphia, PA 19103 (US).</p>	<p>(60) Parent Applications or Grants (63) Related by Continuation US 566,977 (CIP) Filed on 11 January 1990 (11.01.90) US 463,358 (CIP) Filed on 11 January 1990 (11.01.90)</p>	<p>(81) Designated States: AT (European patent), AU, BE (European patent), BR, CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GB (European patent), GR (European patent), HU, IT (European patent), JP, KR, LU (European patent), NL (European patent), NO, SE (European patent), US.</p>	<p>(71) Applicant (for all designated States except US): ISIS PHARMACEUTICALS, INC. [US/US]; 2280 Faraday Avenue, Carlsbad, CA 92008 (US).</p>	<p>Published With international search report.</p>	<p>(72) Inventors; and</p>		<p>(75) Inventors/Applicants (for US only) : COOK, Philip, Dan [US/US]; 6430 La Paloma Street, Carlsbad, CA 92009 (US). ECKER, David, J. [US/US]; 2609 Colibri Lane, Carlsbad, CA 92009 (US). GUINOSSO, Charles, John [US/US]; 313 Encino Drive, Vista, CA 92083 (US). ACEVEDO, Oscar, Leobardo [US/US]; 12117 Brickellia Street, San Diego, CA 92129 (US). KAWASAKI, Andrew, Mamoto [US/US]; 2395 Rancho Del Oro Road, #13, Oceanside, CA 92056 (US). RAMASAMY, Kandasamy [US/US]; 5 Rocky Creek Lane, Laguna Hills, CA 92653 (US).</p>	
<p>(21) International Application Number: PCT/US91/00243</p>																
<p>(22) International Filing Date: 11 January 1991 (11.01.91)</p>																
<p>(30) Priority data: 463,358 11 January 1990 (11.01.90) US 566,977 13 August 1990 (13.08.90) US</p>	<p>(74) Agent: CALDWELL, John, W.; Woodcock Washburn Kurtz Mackiewicz &amp; Norris, One Liberty Place - 46th Floor, Philadelphia, PA 19103 (US).</p>															
<p>(60) Parent Applications or Grants (63) Related by Continuation US 566,977 (CIP) Filed on 11 January 1990 (11.01.90) US 463,358 (CIP) Filed on 11 January 1990 (11.01.90)</p>	<p>(81) Designated States: AT (European patent), AU, BE (European patent), BR, CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GB (European patent), GR (European patent), HU, IT (European patent), JP, KR, LU (European patent), NL (European patent), NO, SE (European patent), US.</p>															
<p>(71) Applicant (for all designated States except US): ISIS PHARMACEUTICALS, INC. [US/US]; 2280 Faraday Avenue, Carlsbad, CA 92008 (US).</p>	<p>Published With international search report.</p>															
<p>(72) Inventors; and</p>																
<p>(75) Inventors/Applicants (for US only) : COOK, Philip, Dan [US/US]; 6430 La Paloma Street, Carlsbad, CA 92009 (US). ECKER, David, J. [US/US]; 2609 Colibri Lane, Carlsbad, CA 92009 (US). GUINOSSO, Charles, John [US/US]; 313 Encino Drive, Vista, CA 92083 (US). ACEVEDO, Oscar, Leobardo [US/US]; 12117 Brickellia Street, San Diego, CA 92129 (US). KAWASAKI, Andrew, Mamoto [US/US]; 2395 Rancho Del Oro Road, #13, Oceanside, CA 92056 (US). RAMASAMY, Kandasamy [US/US]; 5 Rocky Creek Lane, Laguna Hills, CA 92653 (US).</p>																
<p>(54) Title: COMPOSITIONS AND METHODS FOR DETECTING AND MODULATING RNA ACTIVITY AND GENE EXPRESSION</p>																
<p>(57) Abstract</p> <p>Compositions and methods for modulating the activity of RNA and DNA are disclosed. In accordance with preferred embodiments, antisense compositions are prepared comprising targeting and reactive portions. Reactive portions which act, alternatively, through phosphodiester bond cleavage, through backbone sugar bond cleavage or through base modification are preferably employed. Groups which improve the pharmacodynamic and pharmacokinetic properties of the oligonucleotides are also useful in accordance with certain embodiments of this invention. Delivery of the reactive or non-reactive functionalities into the minor groove formed by the hybridization of the composition with the target RNA is also preferably accomplished. Therapeutics, diagnostics and research methods are also disclosed. Synthetic nucleosides and nucleoside fragments are also provided useful for elaboration of oligonucleotides and oligonucleotide analogs for such purposes.</p>																

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MN	Mongolia
BE	Belgium	GA	Gabon	MR	Mauritania
BF	Burkina Faso	GB	United Kingdom	MW	Malawi
BG	Bulgaria	GN	Guinea	NL	Netherlands
BJ	Benin	GR	Greece	NO	Norway
BR	Brazil	HU	Hungary	PL	Poland
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TC	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark				

**COMPOSITIONS AND METHODS FOR  
DETECTING AND MODULATING RNA ACTIVITY  
AND GENE EXPRESSION**

This application is a continuation-in-part of U.S.  
5 Patent Application Serial No. 463,358, filed January 11,  
1990 (ISIS-3), and U.S. Patent Application Serial No.  
566,977, filed August 13, 1990 (ISIS-25).

**BACKGROUND OF THE INVENTION**

10 This invention relates to materials and methods for  
detecting and modulating the activity of RNA. The  
invention generally relates to the field of "antisense"  
compounds, compounds which are capable of specific  
hybridization with a nucleotide sequence of an RNA. In  
accordance with preferred embodiments, this invention is  
15 directed to the design, synthesis and application of  
oligonucleotides and to methods for achieving therapeutic  
treatment of disease, regulating gene expression in  
experimental systems, assaying for RNA and for RNA  
products through the employment of antisense interactions  
20 with such RNA, diagnosing diseases, modulating the  
production of proteins and cleaving RNA in site specific  
fashions.

It is well known that most of the bodily states in  
mammals including most disease states, are effected by  
25 proteins. Such proteins, either acting directly or  
through their enzymatic functions, contribute in major  
proportion to many diseases in animals and man.  
Classical therapeutics has generally focused upon  
interactions with such proteins in efforts to moderate

- 2 -

their disease causing or disease potentiating functions. Recently, however, attempts have been made to moderate the actual production of such proteins by interactions with molecules that direct their synthesis, intracellular  
5 RNA. By interfering with the production of proteins, it has been hoped to effect therapeutic results with maximum effect and minimal side effects. It is the general object of such therapeutic approaches to interfere with or otherwise modulate gene expression leading to  
10 undesired protein formation.

One method for inhibiting specific gene expression is the use of oligonucleotides and oligonucleotide analogs as "antisense" agents. The oligonucleotides or oligonucleotide analogs complimentary to a specific, target,  
15 messenger RNA, mRNA sequence are used. A number of workers have reported such attempts. Pertinent reviews include C.A. Stein & J.S. Cohen, Cancer Research, vol. 48, pp. 2659-2668 (1988); J. Walder, Genes & Development, vol. 2, pp. 502-504 (1988); C.J. Marcus-Sekura, Anal.  
20 Biochemistry, vol. 172, 289-295 (1988); G. Zon, Journal of Protein Chemistry, vol. 6, pp-131-145 (1987); G. Zon, Pharmaceutical Research, vol. 5, pp. 539-549 (1988); A. R. Van der Krol, J.N. Mol, & A.R. Stuitje, BioTechniques, vol. 6, pp. 958-973 (1988) and D.S. Loose-Mitchell, TIPS,  
25 vol. 9, pp. 45-47 (1988). Each of the foregoing provide background concerning general antisense theory and prior techniques.

Thus, antisense methodology has been directed to the complementary hybridization of relatively short oligo-  
30 nucleotides to single-stranded mRNA or single-stranded DNA such that the normal, essential functions of these intracellular nucleic acids are disrupted. Hybridization is the sequence specific hydrogen bonding of oligonucleotides to Watson-Crick base pairs of RNA or single-  
35 stranded DNA. Such base pairs are said to be complementary to one another.

- 3 -

Prior attempts at antisense therapy have provided oligonucleotides or oligonucleotide analogs which are designed to bind in a specific fashion to--which are specifically hybridizable with-- a specific mRNA by hybridization. Such analogs are intended to inhibit the activity of the selected mRNA-- to interfere with translation reactions by which proteins coded by the mRNA are produced-- by any of a number of mechanisms. The inhibition of the formation of the specific proteins which are coded for by the mRNA sequences interfered with have been hoped to lead to therapeutic benefits.

A number of chemical modifications have been introduced into antisense oligonucleotides to increase their therapeutic activity. Such modifications are designed to increase cell penetration of the antisense oligonucleotides, to stabilize them from nucleases and other enzymes that degrade or interfere with the structure or activity of the oligonucleotide analogs in the body, to enhance their binding to targeted RNA, to provide a mode of disruption (terminating event) once sequence-specifically bound to targeted RNA, and to improve their pharmacokinetic properties. At present, however, no generalized antisense oligonucleotide therapeutic or diagnostic scheme has been found. The most serious deficiency of prior efforts has been the complete lack of a termination event once appropriate hybridization takes place or the presence of only a termination event that is so inefficient that a useful potency cannot be achieved due to the inability of oligonucleotides to be taken into cells at effective concentrations. The activity of the antisense oligonucleotides presently available has not been sufficient for effective therapeutic, research reagent, or diagnostic use in any practical sense. Accordingly, there has been and continues to be a long-felt need for oligonucleotides and oligonucleotide analogs which are

- 4 -

capable of effective therapeutic and diagnostic antisense use.

This long-felt need has not been satisfied by prior work in the field of antisense oligonucleotide therapy and diagnostics. Others have failed to provide materials which are, at once, therapeutically or diagnostically effective at reasonable rates of application.

Initially, only two mechanisms or terminating events have been thought to be operating in the antisense approach to therapeutics. These are the hybridization arrest mechanism and the cleavage of hybridized RNA by the cellular enzyme, ribonuclease H (RNase H). It is likely that additional "natural" events may be involved in the disruption of targeted RNA, however.

These naturally occurring events are discussed by Cohen in Oligonucleotides: Antisense Inhibitors of Gene Expression, CRC Press, Inc., Boca Raton, FL (1989). The first, hybridization arrest, denotes the terminating event in which the oligonucleotide inhibitor binds to the target nucleic acid and thus prevents, by simple steric hindrance, the binding of essential proteins, most often ribosomes, to the nucleic acid. Methyl phosphonate oligonucleotides; P.S. Miller & P.O.P. Ts'0, Anti-Cancer Drug Design, 2:117-128 (1987), and  $\alpha$ -anomer oligonucleotides are the two most extensively studied antisense agents which are thought to disrupt nucleic acid function by hybridization arrest.

The second "natural" type of terminating event is the activation of RNase H by the heteroduplex formed between the DNA type oligonucleotide and the targeted RNA with subsequent cleavage of target RNA by the enzyme. The oligonucleotide or oligonucleotide analog, which must be of the deoxyribo type, hybridizes with the targeted RNA and this duplex activates the RNase H enzyme to cleave the RNA strand, thus destroying the normal function of the RNA. Phosphorothioate oligonucleotides are the most prominent example of antisense agents which are thought

- 5 -

to operate by this type of antisense terminating event. R.Y. Walder and J.A. Walder, in Proceedings of the National Academy of Sciences of the U.S.A., Vol. 85, pp.5011-5015 (1988) and C.A. Stein, C. Subasinghe, K. Shenozuka, and J. Cohen, in Nucleic Acids Research, Vol 16, pp.3209-3221 (1988) describe the role the RNase H plays in the antisense approach.

To increase the potency via the "natural" termination events the most often used oligonucleotide modification is modification at the phosphorus atoms. One oligonucleotide analog that has been developed in an effort to secure hybridization arrests is a methyl phosphonate oligonucleotide. Such analogs of oligonucleotides, analogs in the sense that the ordinary structure of the oligonucleotide has been modified into one or more methylphosphonate-substituted structures, have been extensively reported on. A number of authors including K.L. Agarwal & F. Riftina, Nucleic Acids Research, vol. 6, pp. 3009-3024 (1979); P.S. Miller, J. Yano, E. Yano, C. Carroll, C. Jayaraman, K. & P.O.P. Ts'o, Biochemistry, vol. 18, pp. 5134-5143 (1979); K. Jayaraman, K. McParland & P.O.P. Ts'o, Proceedings of the National Academy of Sciences of the U.S.A., vol. 78, pp. 1537-1541 (1981); P. S. Miller, K. B. McParland, K. Jayaraman and P.O.P. Ts'o, Biochemistry, vol. 20, pp. 1874-1880 (1981); P. S. Miller, K. N. Fang, N. S. Kondo and P.O.P. Ts'o, Journal of the American Chemical Society, vol. 93, pp. 6657-6665 (1971); C. H. Agris, K. R. Blake, P. S. Miller, M. P. Reddy, and P.O.P. Ts'o, Biochemistry, vol. 25, 6268-6275 (1986); C. C. Smith, L. Aurelian, M. P. Reddy, P. S. Miller & P.O.P. Ts'o, Proceedings of the National Academy of Sciences of the U.S.A., vol. 83, pp. 2787-2791 (1986); and S. W. Ruby and J. Abelson, Science, vol. 242, pp. 1028-1035 (1988) have reported on such oligonucleotide modifications. In such modifications, a non-bonding phosphoryl oxygen of the phosphorodiester linking moiety is replaced or the nucleotide elements together are



- 6 -

replaced, either in total or in part, by methyl groups. This modification gives the molecule a greater resistance to nucleases. Inhibition of gene expression has been demonstrated with methylphosphonate oligonucleotides targeted to several mRNA's as reported, for example, by D. M. Tidd, T. Hawley, H. M. Warenius & I. Gibson, *Anti-Cancer Drug Design*, vol. 3, pp. 117-127, (1988). Methyl phosphonate modified oligonucleotides do not activate RNase H on hybridization to RNA and only operate in a strictly hybridization arrest mode. Oligonucleotides from this modification class typically possess inferior binding to targeted RNA, likely due to an R/S stereoisomer at each phosphorus atom, as well as increased steric bulk about the phosphate linkage.

Other workers have made other types of modifications to the phosphorus atom of the phosphate backbone of oligonucleotides in attempts to increase the efficiency of oligonucleotide therapy. The most prominent example is the use of phosphorothioate oligonucleotides. These have included HCPF Roelen, E. DeVroom, G. A. Van der Marel & J. H. VanBoom, *Nucleic Acid Research*, vol. 16, pp. 7633-7645 (1988) who employed methyl phosphorothionates. S. Agarwal, J. Goodchild, M. P. Civeira, A. H. Thornton, P. S. Sarin & P. C. Zamecnik, in *Proceedings of the National Academy of Sciences of the U.S.*, vol. 85, pp. 7079-7083 (1988); M. Matsukura, K. Shinozuka, G. Zon, H. Mitsuya, M. Reitz, J. S. Cohen & S. Broder, in *Proceedings of the National Academy of Sciences of the U.S.*, vol. 84, pp. 7706-7710 (1987); and C. J. Marcus-Sekura, A. M. Woerner, K. Shinozuka, G. Zon & G. V. Quinnan, in *Nucleic Acid Research*, vol. 15, pp. 5749-5763 (1987) employed phosphorothioates. See also *Biosis/CA Selects* abstract 110:88603e, reflecting U.S. Serial Number 30,075 filed September 1, 1987 which relates to phosphorothioate modified oligonucleotides. The phosphorothioate modified oligonucleotides are thought to terminate RNA by activation of RNase H upon

- 7 -

hybridization to RNA although hybridization arrest of RNA function may play some part in their activity.

Phosphorodithioates for such use have been disclosed by W.K.D. Brill, J. Y. Yang, Y.X. Ma, & M. H. Caruthers, 5 Journal of the American Chemical Society, vol. 111, pp. 2321-2322 (1989). The phosphorothioate and phosphorodithioate type modifications possess a non-antisense mode of action in that they bind to and inhibit protein function. Protein interactions of oligonucleotides of 10 this type unfortunately undermines the concept of the antisense approach which originally attracted researchers to this novel area. In addition, phosphoroamidates have been disclosed for such uses by A. Jager, M. J. Levy and S. M. Hecht in Biochemistry, vol. 27, pp. 7237-7246 15 (1988); and R. L. Letsinger, S. A. Bach & J. S. Eadie, in Nucleic Acids Research, vol. 14 pp. 3487-3499 (1986).

In contemplating the application of oligonucleotides and oligonucleotide analogs as antisense agents for therapeutic purposes, all applications of oligonucleo- 20 tides as diagnostic, research reagents, and potential therapeutic agents require that the oligonucleotides or oligonucleotide analogs be synthesized in large quantities, be transported across cell membranes or taken up by cells, appropriately hybridize to targeted RNA or 25 DNA, and subsequently terminate or disrupt nucleic acid function. These critical functions depend on the initial stability of oligonucleotides toward nuclease degradation.

A serious deficiency of oligonucleotides for these 30 purposes, particularly antisense therapeutics, is the enzymatic degradation of the administered oligonucleotide by a variety of ubiquitous nucleolytic enzymes, intracellularly and extracellularly located, hereinafter referred to as "nucleases". It is unlikely that 35 unmodified, "wild type", oligonucleotides will be useful therapeutic agents because they are rapidly degraded by nucleases. Modification of oligonucleotides to render

- 8 -

them resistant to nucleases is therefore currently a primary focus of antisense research.

Modifications of oligonucleotides to enhance nuclease resistance have heretofore exclusively taken place on the sugar-phosphate backbone, particularly on the phosphorus atom. Phosphorothioates, methyl phosphonates, phosphorimidates, and phosphorotriesters (phosphate methylated DNA) have been reported to have various levels of resistance to nucleases. However, while the ability of an antisense oligonucleotide to bind to specific DNA or RNA with fidelity is fundamental to antisense methodology, modified phosphorous oligonucleotides, while providing various degrees of nuclease resistance, suffer from inferior hybridization properties.

Phosphorus atom modifications such as methyl phosphonates, phosphorothioates, and phosphoramidates, as described above, confer chirality at the phosphorus atom. Due to this prochiral nature of the phosphorous atom, modifications on the internal phosphorus atoms of modified phosphorous oligonucleotides result in Rp and Sp stereoisomers. Since a practical synthesis of stereoregular oligonucleotides (all Rp or Sp phosphate linkages) is unknown, oligonucleotides with modified phosphorus atoms have  $n^2$  isomers with  $n$  equal to the number of the phosphorus atoms in the oligonucleotide so modified. Furthermore, modifications on the phosphorus atom have unnatural bulk about the phosphorodiester linkage which interferes with the conformation of the sugar-phosphate backbone and consequently, effects the stability of the duplex. The effects of phosphorus atom modifications cause inferior hybridization to the targeted nucleic acids relative to the unmodified oligonucleotide hybridizing to the same target.

The relative ability of an oligonucleotide to bind to complementary nucleic acids is compared by determining the melting temperature of a particular hybridization complex. The melting temperature ( $T_m$ ), a characteristic

- 9 -

physical property of double helixes, denotes the temperature in degrees centigrade at which 50% helical versus coil (unhybridized) forms are present.  $T_m$  is measured by using the UV spectrum to determine the formation and breakdown (melting) of hybridization. Base stacking, which occurs during hybridization, is accompanied by a reduction in UV absorption (hypochromicity). Consequently a reduction in UV absorption indicates a higher  $T_m$ . The higher the  $T_m$ , the greater the strength of the binding of the strands. Non-Watson-Crick base pairing has a strong destabilizing effect on the  $T_m$ . Consequently, absolute fidelity of base pairing is necessary to have optimal binding of an antisense oligonucleotide to its targeted RNA.

Considerable reduction in the hybridization properties of methyl phosphonates and phosphorothioates has been reported by Cohen. Methyl phosphonates have a further disadvantage in that the duplex formed with RNA does not activate degradation by RNase H as a terminating event, but instead acts by hybridization arrest which can be reversed due to a helical melting activity located on the ribosome. Phosphorothioates are highly resistant to most nucleases. However, phosphorothioates typically exhibit non-antisense modes of action, particularly the inhibition of various enzyme functions due to nonspecific binding. Enzyme inhibition by sequence-specific oligonucleotides undermines the very basis of antisense chemotherapy.

While known modifications to oligonucleotides have been shown to have some effect on improving their inhibition of translation, and while such materials have shown some inhibitory activity towards the production of proteins coded by the mRNA, activities which are sufficient for diagnostic or therapeutic use have not been demonstrated.

Oligonucleotides modified to exhibit resistance to nucleases, to activate the RNase H terminating event, and

- 10 -

to hybridize with appropriate strength and fidelity to its targeted RNA (or DNA) are still greatly desired for antisense oligonucleotide purposes.

M. Ikehara et al., European Journal of Biochemistry  
5 139:447-450(1984) report the synthesis of a mixed octamer containing one 2'-deoxy-2'-fluoroguanosine residue or one 2'-deoxy-2'-fluoroadenine residue. W. Guschlbauer and K. Jankowski, Nucleic Acids Res. Vol. 8, p. 1421 (1980) have shown that the contribution of the N form (3'-endo,  
10 2'-exo) increases with the electronegativeness of the 2'-substituent. Thus, 2'-deoxy-2'-fluorouridine contains 85% of the C3'-endo conformer. M. Ikehara et al., Tetrahedron Letters Vol. 42, p. 4073 (1979) have shown that a linear relationship between the  
15 electronegativeness of 2'-substituents and the % N conformation (3'-endo-2'-exo) of a series of 2'-deoxy-adenosines. M. Ikehara et al., Nucleic Acids Research Vol. 5, p. 1877 (1978) have chemically transformed 2'-deoxy-2'-fluoro-adenosine to its 5'-diphosphate. This  
20 was subsequently enzymatically polymerized to provide poly(2'-deoxy- 2'- fluoroadenylic acid).

Furthermore, evidence was presented which indicates that 2'-substituted 2'-deoxyadenosines polynucleotides resemble double stranded RNA rather than DNA. M. Ikehara  
25 et al., Nucleic Acids Res. Vol. 5, p. 3315 (1978) show that a 2'-fluorine substituent in poly A, poly I, and poly C duplexed to their U, C, or I complement are significantly more stable than the ribo or deoxy poly duplexes as determined by standard melting assays. M.  
30 Ikehara et al., Nucleic Acids Res. 4:4249 (1978) show that a 2'-chloro or bromo substituents in poly(2'-deoxy-adenylyc acid) provides nuclease resistance. F. Eckstein et al., Biochemistry Vol. 11, p. 4336 (1972) show that poly(2'-chloro-2'- deoxyuridylic acid) and poly-  
35 (2'-chloro-2'-deoxycytidylic acid) are resistant to various nucleases. H. Inoue et al., Nucleic Acids Research Vol. 15, p.6131 (1987) describe the synthesis of

- 11 -

mixed oligonucleotide sequences containing 2'-OMe at every nucleotide unit. The mixed 2'-OMe substituted sequences hybridized to their ribooligonucleotide complement (RNA) as strongly as the ribo-ribo duplex (RNA-RNA) which is significantly stronger than the same sequence ribo- deoxyribo heteroduplex ( $T_m$ s, 49.0 and 50.1 versus 33.0 degrees for nonamers). S. Shibahara et al., Nucleic Acids Research Vol. 17, p. 239 (1987) describe the synthesis of mixed oligonucleotides sequences containing 2'-OMe at every nucleotide unit. The mixed 2'-OMe substituted sequences were designed to inhibit HIV replication.

It is thought that the composite of the hydroxyl group's steric effect, its hydrogen bonding capabilities, and its electronegateness versus the properties of the hydrogen atom is responsible for the gross structural difference between RNA and DNA. Thermal melting studies indicate that the order of duplex stability (hybridization) of 2'-methoxy oligonucleotides is in the order of RNA-RNA, RNA-DNA, DNA-DNA. The 2'-deoxy-2'-halo, azido, amino, methoxy homopolymers of several natural occurring nucleosides have been prepared by polymerase processes. The required 2'-modified nucleosides monomers have not been incorporated into oligonucleotides via nucleic acids synthesizer machines. Thus, mixed sequence (sequence-specific) oligonucleotides containing 2'-modifications at each sugar are not known except for 2'-deoxy- 2'-methoxy analogs.

Other synthetic terminating events, as compared to hybridization arrest and RNase H cleavage, have been studied in an attempt to increase the potency of oligonucleotides for use in antisense diagnostics and therapeutics. Thus, an area of research has developed in which a second domain to the oligonucleotide, generally referred to as a pendant group, has been introduced.

The pendant group is not involved with the specific Watson-Crick hybridization of the oligonucleotide analog

- 12 -

with the mRNA but is carried along by the oligonucleotide analog to serve as a reactive functionality. The pendant group is intended to interact with the mRNA in some manner more effectively to inhibit translation of the mRNA into protein. Such pendant groups have also been attached to molecules targeted to either single or double stranded DNA.

The type of pendant group known as an intercalating agent has been disclosed by Cazenave, N. Loreau, N. T. Thuong, J. J. Toulme and C. Helene in *Nucleic Acid Research*, vol. 15, pp. 4717-4736 (1987) and J. P. Constant, P. Laugaa, B. P. Roques & J. Lhomme, in *Biochemistry*, vol. 27, pp. 3997-4003 (1988). The disclosed purpose of such intercalating agents is to add binding stability to the hybrid formed between the oligonucleotide analog and the target nucleic acid by binding to the duplex formed between them.

It has also been disclosed to provide a pendant group to oligonucleotide analogs which is capable of cross-linking. Thus, a pendant agent such as psoralin has been disclosed by A. T. Yeung, B. K. Jones and C. T. Chu in *Biochemistry*, vol. 27, pp. 2304-3210 (1988). It is believed that after hybridization of the oligonucleotide analog to the target mRNA, the psoralin is photoactivated to cross-link with the mRNA forming a covalent bond between the oligonucleotide analog and the mRNA thereby permanently inactivating the mRNA molecule and precluding the further formation of protein coded by that particular portion of RNA.

It has also been proposed to employ an alkylating agent as a pendant group for oligonucleotide analogs for use in antisense approaches to diagnostics and therapeutics as disclosed by R.B. Meyer in the *Journal of American Chemical Society*, Vol. 111, pp 8517-8519 (1989) and D.G. Knorre and V.V. Vlassov, *Progress in Nucleic Acid Research and Molecular Biology*, Vol.32, pp.291-320 (1985).

- 13 -

The object of employing alkylating agents and pendant groups in oligonucleotide analogs in antisense approaches is to cause the alkylating agent to react irreversibly with the target mRNA. Such irreversible binding between  
5 the antisense oligonucleotide and the mRNA is generally covalent and leads to permanent inactivation of the mRNA with a concomitant halt in protein production from the portion of mRNA thus inactivated.

A further strategy which has been proposed is to use  
10 chemical reagents which, under selected conditions, can generate a radical species for reaction with the target nucleic acid to cause cleavage or otherwise to inactivate it. Proposed pendant groups of this category include coordination complexes containing a metal ion with  
15 associated ligands. A metal ion can change oxidation state to generate reactive oxygen-containing radical ions or other radical species. P.L. Doan, L. Perrouault, M. Chassignol, N. T. Thuong, & C. Helene, in Nucleic Acids Research, Vol. 15, pp. 8643-8659 (1987) have disclosed  
20 iron/EDTA and iron/porphrin species for this purpose. Copper/phenanthroline complexes have been disclosed by D. S. Sigman, in Accounts of Chemical Research, Vol. 19, pp. 180-186 (1986). G.B. Dreyer and P.B. Dervan, in Proceedings of the National Academy of Sciences, U.S.A.,  
25 Vol. 82, pp.968-972 (1985) have investigated the EDTA/Fe moiety to cleave nucleic acids.

Prior approaches using cross-linking agents, alkylating agents, and radical generating species as pendant groups on oligonucleotides for antisense  
30 diagnostics and therapeutics have several significant shortcomings. The sites of attachment of the pendant groups to oligonucleotides play an important, yet imperfectly known, part in the effectiveness of oligonucleotides for therapeutics and diagnostics.  
35 Prior workers have described most pendant groups as being attached to a phosphorus atom which, as noted above, affords oligonucleotides with inferior hybridization



- 14 -

properties. Prior attempts have been relatively insensitive, that is the reactive pendant groups have not been effectively delivered to sites on the messenger RNA molecules for alkylation or cleavage in an effective proportion. Moreover, even if the reactivity of such materials were perfect, i.e. if each reactive functionality were to actually react with a messenger RNA molecule, the effect would be no better than stoichiometric. That is, only one mRNA molecule would be inactivated for each molecule of oligonucleotide. It is also likely that the non-specific interactions of the modified oligonucleotides with molecules other than the target RNA, for example with other molecules that may be alkylated or which may react with radical species, as well as self-destruction, not only diminishes the diagnostic or therapeutic effect of the antisense treatment but also leads to undesired toxic reactions in the cell or in vitro. This is especially acute with the radical species which are believed to be able to diffuse beyond the locus of the specific hybridization to cause undesired damage to non-target materials, other cellular molecules, and cellular metabolites. This perceived lack of specificity and stoichiometric limit to the efficacy of such prior alkylating agent and radical generating-types of antisense oligonucleotides is a significant drawback to their employment.

Accordingly, there remains a great need for antisense oligonucleotide formulations which are capable of improved specificity and effectiveness both in binding and in mRNA modulation or inactivation without the imposition of undesirable side effects.

#### OBJECTS OF THE INVENTION

It is a principal object of this invention to provide oligonucleotides and oligonucleotide analogs for use in antisense oligonucleotide diagnostics and therapeutics.

- 15 -

It is a further object of the invention to provide nuclease resistant, sugar modified oligonucleotides or oligonucleotide analogs for use in antisense oligonucleotide diagnostics, research reagents, and  
5 therapeutics.

It is a further object of this invention to provide such oligonucleotides and oligonucleotide analogs which are effective in modulating the activity of a DNA or an RNA.

10 A further object of this invention is to provide such oligonucleotides and oligonucleotide analogs which are less likely to invoke undesired or toxic side reactions.

A further object is to provide research and diagnostic methods and materials for assaying bodily states in  
15 animals, especially diseased states.

A further object is to provide means for modifying nucleic acids for effecting substitutions on selective portions thereof.

Yet another object is to provide therapeutic and  
20 research methods and materials for the treatment of diseases through modulation of the activity of DNA and RNA.

Still another object is to provide means for the selective cleavage of RNA.

25 These and other objects will become apparent to persons of ordinary skill in the art from a review of the present specification and attendant claims.

#### SUMMARY OF THE INVENTION

In accordance with the present invention, compositions  
30 for modulating the activity of DNA and RNA are provided. These compositions comprise a targeting portion specifically hybridizable with a preselected nucleotide sequence of RNA. The composition further provides a reactive portion capable of catalyzing, alkylating, or  
35 therwis effecting the cleavage of RNA, especially f its phosphodiester bonds. The compositions may also

- 16 -

include a tether, a means for connecting the targeting and reactive portions together to form the composition. In accordance with preferred embodiments, the compositions are adapted for placement of the reactive  
5 portion into the minor groove site of the hybrid structure formed from the hybridization of the RNA and the composition.

The targeting portion of the compositions of this invention preferably comprises an oligonucleotide analog  
10 comprising from about 3 to about 50 base units with 8 to 40 subunits being preferred and 12 to 25 being still more preferred. Oligonucleotides or oligonucleotide analogs having about 15 base units are preferable for the practice of certain embodiments of the present invention.  
15 In accordance with other preferred embodiments, the targeting portion is an analog of an oligonucleotide wherein at least some of the phosphodiester bonds of the oligonucleotide have been substituted with a structure which functions to enhance the ability of the  
20 compositions to penetrate into the intracellular region of cells where the RNA whose activity is to be modulated is located. It is preferred that such substitutions comprise phosphorothioate bonds or short chain alkyl or cycloalkyl structures. In accordance with other  
25 preferred embodiments, the phosphodiester bonds are substituted with structures which are, at once, substantially non-ionic and non-chiral.

In accordance with other preferred embodiments the reactive portion of the composition comprises a  
30 functionality capable of catalyzing the hydrolysis, cleavage, of phosphodiester bonds in RNA. Such functionalities may either be basic, acidic, or amphoteric. Heteroatomic species can be formulated to be either basic or acidic or, indeed, to be amphoteric for  
35 such purposes.

This invention also comprehends the employment of alkylating and free radical forming functionalities as

- 17 -

the reactive portions of the subject compositions where said alkylating or free radical forming materials are delivered into the minor groove of the hybrid formed between the compositions of the invention and the RNA to be modulated.

In accordance with other embodiments of the invention, compositions are provided for modulating the activity of an RNA comprising certain heterocyclic structures having at least one RNA cleaving moiety or other moiety capable of interacting with an RNA appended thereto. These compositions are adapted for placement of the reactive, RNA cleaving moiety or other reactive moiety into the minor groove site of the hybrid structure formed from the RNA and the composition through careful selection of the sites of attachment of the RNA cleaving moieties. Such compositions may include sugar or sugar analog portions and novel nucleoside base portions. Accordingly, novel nucleosides and nucleoside analogs are provided. Such nucleosides and nucleoside analogs may be incorporated into oligonucleotides and oligonucleotide analogs which are useful in the practice of the invention.

The invention also is directed to methods for modulating the activity of an RNA comprising contacting an organism having the RNA with a composition formulated in accordance with the foregoing considerations. It is preferred that the RNA which is to be modulated be preselected to comprise preferably messenger RNA which codes for a protein whose formation is to be modulated. The invention may also be applied to pre-messenger RNA and, indeed, to RNA generically and to double stranded DNA. The targeting portion of the composition to be employed is, thus, selected to be complementary to the preselected portion of RNA, that is, to be an antisense oligonucleotide for that portion.

This invention is also directed to methods for treating an organism having a disease characterized by the undesired production of a protein comprising

- 18 -

contacting the organism with a composition in accordance with the foregoing considerations, preferably one which is designed to specifically bind with messenger RNA which codes for the protein whose production is to be inhibited.

The invention is also directed to the utilization of groups appended to oligonucleotides which do not include a reactive function. Such pendant groups may lead to enhanced oligonucleotide uptake, enhanced resistance of oligonucleotide to degradation by nucleases, and stronger binding of the oligonucleotides to targeted RNA. Such oligonucleotide can disrupt RNA by hybridization arrest and/or by activation of RNase H as well as by cleavage of the RNA phosphodiester bonds or other destruction thereof by other reactive functionalities.

The invention is also directed to functionalities which serve to attach reporter groups such as biotin and various fluorophores to sequence-specific oligonucleotides for diagnostic purposes. More than one non-reactive functionality may be attached to each oligonucleotide, two or more non-reactive functionalities may be attached to a single nucleoside unit, and a combination of non-reactive functionalities and reactive functionalities may be attached to a single nucleoside unit or a single oligonucleotide. Non-reactive functionalities and reactive functionalities attached to sequence-specific oligonucleotides are preferably designed to reside in the minor groove or on the minor groove side of the heteroduplex.

In accordance with other preferred embodiments, the present invention, compositions which are resistant to nuclease degradation and which modulate the activity of DNA and RNA are provided. These compositions are comprised of sugar modified oligonucleotides or oligonucleotid analogs, the targeting portions of which are specifically hybridizable with preselected nucleotide sequences of single-stranded or double-stranded DNA or

- 19 -

RNA. The sugar modified oligonucleotides recognize and form double strands with single-stranded DNA and RNA or triple strands with double-stranded DNA and RNA.

The nuclease resistant oligonucleotides of this invention consist of a single strand of nucleic acid bases linked together through linking groups. The target portion of the nuclease resistant oligonucleotide may range in length from about 5 to about 50 nucleic acid bases. However, in accordance with the preferred embodiment of this invention, a target sequence of about 15 bases in length is optimal.

The nucleic acid bases may be pyrimidines such as thymine, uracil or cytosine, or purines such as guanine or adenine, or both, arranged in a specific sequence. Additionally, they may be any of the novel bases of the invention. The sugar moiety of such bases may be of the deoxyribose or ribose type. The groups linking the bases together may be the usual sugar phosphate nucleic acid backbone, but may also be modified as a phosphorothioate, methylphosphonate, or phosphate alkylated moiety to further enhance the sugar modified oligonucleotide properties, along with removal of a 5'-methylene group and/or carbocyclic sugar.

In accordance with one embodiment of this invention, the targeting portion is an analog of an oligonucleotide wherein at least one of the 2'-deoxy ribofuranosyl moieties of the nucleoside unit is modified. A hydrogen or a hydroxyl, halo, azido, amino, methoxy or alkyl group may be added. For example, H, OH, lower alkyl, substituted lower alkyl, F, Cl, Br, CN, CF<sub>3</sub>, OCF<sub>3</sub>, OCN, O-alkyl, S-alkyl, SOME, SO<sub>2</sub>Me, ONO<sub>2</sub>, NO<sub>2</sub>, N<sub>3</sub>, NH<sub>2</sub>, NH-alkyl, OCH<sub>2</sub>CH=CH<sub>2</sub>, OCH=CH<sub>2</sub>, OCH<sub>2</sub>CCH, OCCH, aralkyl, heteroaralkyl, heterocycloalkyl, aminoalkylamino, heterocycloalkylamino, polyalkylamino, substituted silyl, a RNA cleaving moiety or a group for improving the pharmacodynamic properties of an oligonucleotide or a group for improving the pharmacokinetic properties of an

- 20 -

oligonucleotide where alkyl is a straight or branched chain of C1 to C12 may be used, with unsaturation within the carbon chain, such as allyloxy being particularly preferred.

- 5     The resulting novel oligonucleotides or oligonucleotide analogs are resistant to nuclease degradation and exhibit hybridization properties of higher quality relative to wild type (DNA-DNA and RNA-DNA) duplexes and the phosphorus modified oligo-  
10 nucleotide antisense duplexes containing phosphorothioates, methylphosphonates, phosphoramidates and phosphorotriesters.

- The invention further is directed to diagnostic methods for detecting the presence or absence of abnormal  
15 RNA molecules or abnormal or inappropriate expression of normal RNA molecules in organisms or cells. It is also directed to methods for the selective cleaving of RNA such as for research and diagnostics and to the RNA thus formed. Such selective cleaving is accomplished by  
20 interacting RNA with compositions of the invention which have reactive portions capable of effecting such cleavage and targeting portions designed to enforce selectivity.

- The compositions useful for modulating the activity of an RNA or detecting its presence in accordance with this  
25 invention generally comprise three portions. The first portion, the targeting portion, is a portion which is specifically hybridizable with a preselected nucleotide sequence of the RNA to be modulated. It is generally desirable to select a sequence of RNA which codes for or  
30 which is involved in the production of proteins whose synthesis is ultimately to be modulated or inhibited in entirety. The targeting portion of the composition is an oligonucleotide analog. It is designed and prepared conveniently, through solid state synthesis of known  
35 methodology, to be complementary to or at least to be specifically hybridizable with the preselected nucleotide sequence of the RNA. Nucleic acid synthesizers are

- 21 -

commercially available and their use is generally understood by persons of ordinary skill in the art to be capable of generating nearly any oligonucleotide of reasonable length which may be desired.

5 The invention is also directed to methods for modulating the production of a protein by an organism comprising contacting the organism with a composition formulated in accordance with the foregoing considerations. It is preferred that the RNA or DNA  
10 portion which is to be modulated be preselected to comprise that portion of DNA or RNA which codes for the protein whose formation is to be modulated. The targeting portion of the composition to be employed is, thus, selected to be complementary to the preselected  
15 portion of DNA or RNA, that is to be an antisense oligonucleotide for that portion.

This invention is also directed to methods of treating an organism having a disease characterized by the undesired production of a protein. This method comprises  
20 contacting the organism with a composition in accordance with the foregoing considerations. The composition is preferably one which is designed to specifically bind with messenger RNA which codes for the protein whose production is to be inhibited.

25 The invention further is directed to diagnostic methods for detecting the presence or absence of abnormal RNA molecules or abnormal or inappropriate expression of normal RNA molecules in organisms or cells.

The invention is also directed to methods for the  
30 selective binding of RNA for research and diagnostic purposes. Such selective, strong binding is accomplished by interacting such RNA or DNA with compositions of the invention which are resistant to degradative nucleases and hybridize stronger and with greater fidelity than any  
35 other known oligonucleotide or oligonucleotide analog.

In accordance with a further embodiment of the invention, novel processes are provided for the synthesis



- 22 -

of novel nucleoside analogs that are substituted in the 2' position and which are useful for incorporation into oligonucleotides and analogs of the invention. Such process provides for introduction of a 2'substituent in the absence of blocking of either the 3' or 5'hydroxyl groups of a ribofuranosyl nucleoside. Such processes utilize treatment with sodium hydride followed by use of an alkyl halide or with uracil, stannous chloride. No protecting groups on the nucleoside bases are necessary except for the exo-cyclic amino group of guanosine. The reactions are conducted at or near room temperature. These conditions are contrasted to prior known process that require strong alkylating agents, as for instance diazomethane. Such strong alkylating agents necessitate the complete protection of all reactive sites on the nucleoside bases and the 3' and 5' sugar hydroxyls.

In accordance with an even further embodiment of the invention, a novel process is provided for the synthesis of novel 3-deaza nucleosides and oligonucleotides that incorporate these nucleosides. In this process compounds such as 5-cyanomethyl-1- (2'-deoxy-3',5'-di-O-p-toluoyl-D-erythro-pentofuranosyl) imidazole-4-carboxylate or 5-cyano-[reactive substituent] methyl-1- (2'-deoxy- D-erythro-pento furanosyl) imidazole-4-carboxylate are prepared. The 5-cyano-[reactive substituent]-methyl compounds are prepared from the corresponding 5-cyanomethyl compound utilizing sodium hydride and an appropriate alkyl halide. The carboxylates are deblocked and converted to the corresponding carboxamides. The 5' position of the compounds are then protected and the compounds derivatized at the 3' position with an appropriate phosphoramidite or other activated phosphate group suitable for use on a DNA synthesizer. An oligonucleotide is constructed on the synthesizer. Such oligonucleotide incorporates one or more of the 5-cyano- [reactive substituent]-methyl imidazole compounds in its sequence. The imidazole

- 23 -

compound cyclizes to the corresponding 3-deaza purine or 3- substituted -3-deaza purine concurrently with cleavage of the oligonucleotide from the synthesizer support upon treatment with excess concentrated ammonium hydroxide.

5 The modified nucleotides (modified monomers) possessing the required functionality in the sugar portion and/or the heterocyclic portion may be used to prepare desired novel, oligonucleotides and analogs of this invention. The structure and synthesis of the  
10 modified nucleotides have not been known previously and are herewithin described for the first time. The sites of functionality in the sugars and heterocycles are novel, and have been preferably designed such that the functionalities will reside in or on the minor groove  
15 formed by the heteroduplex between the modified oligonucleotide and the targeted RNA. The minor side or minor groove of the duplexes formed between such modified oligonucleotides and the targeted RNA has been found to be the greatly preferred site for functional group  
20 activity.

One group of compositions useful for modulating the activity of an RNA or DNA molecule in accordance with this invention generally comprise a sugar modified oligonucleotide containing a target sequence which is  
25 specifically hybridizable with a preselected nucleotide sequence of single stranded or double stranded DNA or RNA molecule and which is nuclease resistant.

It is generally desirable to select a sequence of DNA or RNA for or which is involved in the production of  
30 proteins whose synthesis is ultimately to be modulated or inhibited in entirety. The targeting portion of the composition is generally an oligonucleotide analog. It is synthesized, conveniently through solid state synthesis of known methodology, to be complementary to or  
35 at least to be specifically hybridizable with the preselected nucleotide sequence of the RNA or DNA. Nucleic acid synthesizers are commercially available and

- 24 -

their use is generally understood by persons of ordinary skill in the art as being effective in generating nearly any oligonucleotide of reasonable length which may be desired.

5 In the context of this invention, the term "oligonucleotide" refers to a plurality of joined nucleotide units formed in a specific sequence from naturally occurring bases and pentofuranosyl groups joined through sugar groups by native phosphodiester  
10 bonds. These nucleotide units may be nucleic acid bases such as guanine, adenine, cytosine, thymine or uracil. The sugar group may be deoxyribose or ribose. This term refers to both naturally occurring or synthetic species formed from naturally occurring subunits.

15 "Oligonucleotide analog" or "modified oligonucleotides" as these terms are used in connection with this invention, refer to moieties which function similarly to natural oligonucleotides but which have non-naturally occurring portions. Oligonucleotide analogs  
20 or modified oligonucleotides may have altered sugar moieties, altered bases, both altered sugars and bases or altered inter-sugar linkages, for example phosphorothioates and other sulfur containing species which are known for use in the art. The oligonucleotide analogs  
25 of the invention may increase nuclease resistance of the oligonucleotide composition and thus facilitate antisense therapeutic, diagnostic use or research reagent use of the compositions of this invention.

It is greatly preferred in the present invention to  
30 employ oligonucleotide analogs or modified oligonucleotides rather than the oligonucleotides themselves. In this context, oligonucleotide analog refers to a structure which is generally similar to native oligonucleotides, but which has been modified in one or more  
35 significant ways.

It is generally preferred for use in some embodiments of this invention that some positions of the nucleotide

- 25 -

base be substituted in order to increase the nuclease resistance of the composition while maintaining the integrity of the oligonucleotide binding capabilities.

It is preferred in some embodiments of the present invention to employ further modified oligonucleotides. In this context, modified oligonucleotide analogs refers to a structure which is generally similar to native oligonucleotides, but which have been modified in one or more significant ways.

10 In order to enhance penetration into the intracellular spaces of cells where the messenger RNA or DNA, which are the targets of the overall composition reside, preferred embodiments of the invention provide modifications of the oligonucleotides. Such modifications may provide  
15 oligonucleotides that are substantially less ionic than native forms. Such modifications facilitate penetration of oligonucleotides into the intracellular spaces. Any of the existing or yet to be discovered methods for accomplishing this goal may be employed in accordance  
20 with the practice of the present invention. At present, it has been found preferable to employ substitutions for the phosphorodiester bond, which substitutions are not only relatively less ionic than the naturally occurring bonds but are also substantially nonchiral.

25 As will be appreciated, the phosphorus atom in the phosphorodiester linkage is "prochiral". Modifications at the phosphorus, such as is done in methyl phosphonates and phosphorothioates type oligonucleotides, result in essentially chiral structures. Chirality results in the  
30 existence of two isomers at each chiral center which may interact differently with cellular molecules. Such an unresolved mixture of isomers may inhibit the transport of the resulting compositions into the intracellular spaces or decrease the affinity and specificity of  
35 hybridization to the specific target RNA or DNA. Thus, it is preferred in some embodiments of this invention to employ substantially non-ionic, substantially non-chiral

- 26 -


entities in lieu of some or all of the phosphorodiester bonds. For this purpose, short chain alkyl or cycloalkyl structures especially C<sub>2</sub>-C<sub>4</sub> structures are preferred. As is set forth in an application having U.S. Serial Number 5 558,663 and assigned to a common assignee hereof, said application being entitled "Polyamine Oligonucleotides to Enhance Cellular Uptake," filed July 27, 1990 the modification of the sugar structure including the elimination of one of the oxygen functionalities may 10 permit the introduction of such substantially non-chiral, non-ionic substituents in this position. The entirety of the disclosure of that application is incorporated herein by reference in order to disclose more fully such modifications.

15 In keeping with the goals of the invention are the standard backbone modifications such as substituting P for S, Me-P, MeO-P, H<sub>2</sub>N-P, etc. These substitutions are thought in some cases to enhance the sugar modified oligonucleotide properties.

20 The targeting portions of the compositions of the present invention, are preferably oligonucleotide analogs having from about 3 to about 50 base units. It is more preferred that such analogs have from about 8 to about 40 base units and even more preferred that from about 12 to 25 about 25 be employed. At present, it is believed that oligonucleotides or analogs having about 15 base units will likely be found to be best for the practice of certain embodiments of the present invention. It is desired that the targeting portion be adapted so as to be 30 specifically hybridizable with the preselected nucleotide sequence of the RNA selected for modulation.

At present, the oligonucleotide analogs which are believed to be particularly suitable for the practice of one or more embodiments of the present invention will 35 comprise one or more subunits having gross schematic structures as shown in Formula 1.

- 27 -



1


- 5 whereir. B is any of the purine or pyrimidine bases, modified bases or base analogs including those which are known in naturally-occurring oligonucleotides or for use in modified oligonucleotides, or which exhibit similar functions; E is a RNA cleaving moiety, a group for
- 10 improving the pharmacokinetic properties of said oligonucleotide, a group for improving the pharmacodynamics of said oligonucleotide, H, or OH and other small base substituent groups; S is a sugar or sugar analog; and L
- 15 is a sugar linking group. The sugar linking group L may be any of those structures either naturally occurring, described herein, or otherwise known which are capable of linking sugar moieties of oligonucleotides or sugar analogs, together to form the targeting portion of the compositions of this invention. It is preferred that
- 20 these sugar linking functions either comprise a phosphodiester structure or a substantially non-ionic, substantially non-chiral structure as described hereinbefore, such as lower alkyl and cycloalkyl, especially C<sub>2</sub> - C<sub>4</sub> alkyl. Note that when lower alkyl structures are
- 25 employed, the 5' methylenes of one or more sugars may be eliminated. Preferable, at least some of the phosphodiester bonds of said oligonucleotide are substituted phosphorothionate, methyl phosphonate, or alkyl phosphate.
- 30 As will be appreciated by persons of ordinary skill in the art, variations in the structures of the sugar moieties useful in the preparation of the subject compositions may be made with ut deviating from the

- 28 -


spirit of the invention. Once again, it is not necessary that every sugar linking function be in a modified form. A substantial number and even a predominance of such linking groups may exist in the native, phosphodiester form as long as the overall targeting portion of the compositions of the molecules exhibits an effective ability to penetrate into the intracellular spaces of cells of the organism in question or otherwise to contact the target RNA and to specifically bind therewith to form a hybrid capable of detecting and modulating the RNA activity. Of course, fully unmodified, native phosphodiester structures may be used as well.

It is not necessary to tether more than one, or perhaps two RNA cleaving functionalities, groups for improving the pharmacodynamics of oligonucleotides or groups for improving the pharmacokinetics of oligonucleotides, to oligonucleotides in accordance with this invention in order to provide the benefits of the invention. Thus, an RNA cleaving moiety or pharmacodynamic improving group or pharmacokinetic improving group will preferably be tethered to a relatively small proportion of the subunits, generally only one or two, comprising the oligonucleotide analog which is the targeting portion of the compositions of the invention. In other embodiments of the invention, however, all of the nucleotides in an oligonucleotide can be modified to include one or more RNA cleaving moiety groups, pharmacodynamic improving groups or pharmacokinetic improving groups.


It is believed desirable in accordance with certain preferred embodiments, to attach the RNA cleaving portion or pharmacodynamic improving group portion or pharmacokinetic improving group portion of the compositions of this invention to one of the nucleosides forming the subunits of the targeting portion. Such an attachment may be depicted in accordance with Formulas 2 through 7.




2




4




6



3



5



7

Where G and K are, independently, C or N;

5 J is N or CR<sub>3</sub>;

R<sub>1</sub> is OH or NH<sub>2</sub>;

R<sub>2</sub> and R<sub>3</sub> are H, NH<sub>2</sub>, lower alkyl, substituted lower alkyl, lower alkenyl, aralkyl, alkylamino, aralkylamin ,



- 30 -

substituted alkylamino, heterocycloalkyl, heterocyclo-  
alkylamino, aminoalkylamino, hetrocycloalkylamino,  
polyalkylamino, a RNA cleaving moiety, a group for  
improving the pharmacokinetic properties of an oligo-  
5 nucleotide or a group for improving the pharmacodynamic  
properties of an oligonucleotide;

$R_4$  and  $R_5$  are H, OH,  $NH_2$ , lower alkyl, substituted lower  
alkyl, substituted amino, a RNA cleaving moiety, a group  
for improving the pharmacokinetic properties of an  
10 oligonucleotide or a group for improving the pharmaco-  
dynamic properties of an oligonucleotide;

$R_6$  and  $R_7$  are H, OH,  $NH_2$ , SH, halogen,  $CONH_2$ ,  $C(NH)NH_2$ ,  
 $C(O)O$ -alkyl,  $CSNH_2$ , CN,  $C(NH)NHOH$ , lower alkyl,  
substituted lower alkyl, substituted amino, a RNA  
15 cleaving moiety, a group for improving the pharmaco-  
kinetic properties of an oligonucleotide or a group for  
improving the pharmacodynamic properties of an oligo-  
nucleotide;

X is a sugar or a sugar analog moiety and where said  
20 sugar analog moiety is a sugar substituted with at least  
one substituent comprising a RNA cleaving moiety, a  
group for improving the pharmacodynamic properties of an  
oligonucleotide, or a group for improving the  
pharmacokinetic properties of an oligonucleotide;

25 provided that when said composition is represented by  
Formula 2 and when G is N and J is N then X is said sugar  
analog moiety substituted with at least one group other  
than F and  $OCH_3$ ;

further provided that if said oligonucleotide includes  
30 only one of said modified nucleotide and  $G=N$  and  $J=N$  and  
all of the sugar linking groups are unsubstituted phos-  
phordiester linkages then X is not a sugar or a sugar  
analog moiety containing a  $2'-OCH_3$  or  $2'-F$ ;

further provided when said composition is represented  
35 by Formula 2 and G is N and J is  $CR_3$  and  $R_3$  is H then X is  
said sugar analog moiety;

- 31 -


further provided that when said composition is represented by Formula 4 and when G is N and X is said sugar, R<sub>2</sub> is not H;

5 further provided that when said composition is represented by Formula 6 and R<sub>6</sub> is H and R<sub>2</sub> is NH<sub>2</sub> and R<sub>7</sub> is CONH<sub>2</sub>, CSNH<sub>2</sub>, C(O)O-alkyl, C(NH)NH<sub>2</sub> or C(NH)NHOH then X is said sugar analog moiety;


10 further provided that when said composition is represented by Formula 6 and R<sub>6</sub> is H, OH or SH and R<sub>7</sub> is C(O)O-alkyl or C(NH)NH<sub>2</sub> and R<sub>2</sub> is -CH<sub>2</sub>CN then X is said sugar analog moiety; and

further provided that when said composition is represented by Formula 7 and R<sub>3</sub> is H and G is C then X is said sugar analog moiety.

15 It is believed that attachment of the RNA cleaving functionality to the 2' position of the furanosyl portion of a nucleoside forming part of the oligonucleotides of this invention is preferred in certain embodiments in order to permit the tether portion to deliver the  
20 reactive functionality of the compositions of the invention into the minor groove formed by the hybridization of the composition with the messenger RNA. Other sites of attachment on the "sugar" portion of the nucleosides forming the oligonucleotides of this  
25 invention, which are believed to be preferred for placing a reactive functionality of the invention into the minor groove upon hybridization of the composition with a messenger RNA are depicted in Formulas 8 and 9. These moieties may include sugar analogs which are not strictly  
30 within the conventional definition of sugars but which serve substantially similar functions.



8



9

Where Q is O or  $\text{CHR}_{11}$ ;

Z is a point of attachment which can be a naturally occurring or synthetic variable base moiety;

- 5  $R_8$  and  $R_9$  are H, lower alkyl, substituted lower alkyl, a RNA cleaving moiety, a group for improving the pharmacodynamic properties of an oligonucleotide or a group for improving the pharmacokinetic properties of an oligonucleotide;
- 10  $R_{10}$  is H, OH, lower alkyl, substituted lower alkyl, F, Cl, Br, CN,  $\text{CF}_3$ ,  $\text{OCF}_3$ , OCN, O-alkyl, S-alkyl, SOME,  $\text{SO}_2\text{Me}$ ,  $\text{ONO}_2$ ,  $\text{NO}_2$ ,  $\text{N}_3$ ,  $\text{NH}_2$ , NH-alkyl,  $\text{OCH}_2\text{CH}=\text{CH}_2$ ,  $\text{OCH}=\text{CH}_2$ ,  $\text{OCH}_2\text{CCH}$ ,  $\text{OCCH}$ , aralkyl, heteroaralkyl, heterocycloalkyl, amino-alkylamino, heterocycloalkyl, polyalkylamino, substituted silyl, a RNA cleaving moiety, a group for improving the pharmacodynamic properties of an oligonucleotide or a group for improving the pharmacokinetic properties of an oligonucleotide; and
- 15  $R_{11}$  is H,  $\text{OH}$ , lower alkyl, substituted lower alkyl, a RNA cleaving moiety, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide.
- 20 It is preferred that  $R_{10}$  be lower alkyl, substituted lower alkyl, CN,  $\text{CF}_3$ ,  $\text{OCF}_3$ , OCN, O- $\text{C}_3$ - $\text{C}_{12}$ -alkyl, S-alkyl, SOME,  $\text{SO}_2\text{Me}$ ,  $\text{ONO}_2$ ,  $\text{NO}_2$ , NH-alkyl,  $\text{OCH}=\text{CH}_2$ ,  $\text{OCH}_2\text{CCH}$ ,  $\text{OCCH}$ , aralkyl, heteroaralkyl, heterocyclo-alkyl, polyalkyl-amino, substituted silyl, a RNA cleaving moiety or a
- 25

- 33 -

group for improving the pharmacodynamic properties of an oligonucleotide or a group for improving the pharmacokinetic properties of an oligonucleotide.

Exemplary base moieties are those set forth in Formulas 2 through 7. The sugar, X, preferably is ribofuranosyl or 2'-deoxyribofuranosyl. A preferred sugar analog moiety is 2'-deoxy-2'-substituted ribofuranosyl. Other sugar analog moieties are as above in Formulas 8 and 9. The base B of Formulas 8 and 9 are any of the natural pyrimidinyl-1- or purinyl-9 bases including uracil, thymine, cytosine, adenine and guanine, 5-alkylcytosine such as 5-methylcytosine as well novel bases of the invention as depicted in Formulas 2 through 7.

Alkyl groups of the invention include but are not limited to C<sub>1</sub>-C<sub>12</sub> straight and branched chained alkyls such as methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, undecyl, dodecyl, isopropyl, 2-butyl, isobutyl, 2-methylbutyl, isopentyl, 2-methylpentyl, 3-methylpentyl, 2-ethylhexyl, 2-propylpentyl. Alkenyl groups include but are not limited to unsaturated moieties derived from the above alkyl groups including but not limited to vinyl, allyl, crotyl, propargyl. Aryl groups include but are not limited to phenyl, tolyl, benzyl, naphthyl, anthracyl, phenanthryl, and xylyl. Halogens include fluorine, chlorine and bromine. Suitable heterocyclic groups include but are not limited to imidazole, tetrazole, triazole, pyrrolidine, piperidine, piperazine and morpholine. Amines include amines of all of the above alkyl, alkenyl and aryl groups including primary and secondary amines and "masked amines" such as phthalimide. Amines are also meant to include polyalkylamino compounds and aminoalkylamines such as aminopropylamine and further heterocyclic alkylamines such as imidazol-1, 2 or 4-yl-propylamine. Substituent groups for the above include but are not limited to other alkyl, haloalkyl, alkenyl, alkoxy,

- 34 -

thioalkoxy, haloalkoxy and aryl groups as well as halogen, hydroxyl, amino, azido, carboxy, cyano, nitro, mercapto, sulfides, sulfones and sulfoxides. Other suitable substituent groups also include rhodamines, coumarins, acridones, pyrenes, stilbenes, oxazolo-pyridocarbazoles, anthraquinones, phenanthridines, phenazines, azidobenzenes, psoralens, porphyrins and cholesterol.

The sites that functionality may be attached to in the nucleosidic units which in turn may be converted to modified oligonucleotide is critical in the design of compositions for sequence-specific destruction or modulation of targeted RNA. The functionality must not interfere with Watson-Crick base pair hydrogen bonding rules as this is the sequence-specific recognition/binding factor essential for selection of the desired RNA to be disrupted.

Furthermore, approaches to perfect complementation between the modified oligonucleotides and targeted RNA will result in the most stable heteroduplexes. This is desired because the heteroduplex must have a sufficient half life to allow the reactive or non-reactive functionalities of this invention to initiate the cleavage or otherwise disruption of RNA function.

The half life of the perfectly formed duplex will be greatly effected by the positioning of the tethered functional group. Inappropriate positioning of functional groups, such as placement on the Watson/Crick base pair sites, would preclude duplex formation. Other attachment sites may allow sequence-specific binding but may be of such low stability that the reactive functionality will not have sufficient time to initiate RNA disruption.

For RNA inactivation, a further important factor concerning the placement of the tethered functionality is that it must be in appropriate proximity to reactive substrate located in the targeted RNA, especially the

- 35 -

most sensitive "trigger" point, the 2'-hydroxyl group. A variety of structural studies such as X-ray diffraction, chemical reaction, and molecular modeling studies suggests that the 2'-hydroxyl group of RNA in a duplex or heteroduplex resides in the minor groove. Therefore, functionality placed on sequence-specific oligonucleotides (via modified nucleosides) should preferably reside in the minor groove formed between the oligonucleotide and the targeted RNA, not interfere with duplex formation and stability, and initiate cleavage or disruption of the RNA.

It has now been found that certain positions on the nucleosides of double stranded nucleic acids are exposed in the minor groove and may be substituted without affecting Watson-Crick base-pairing or duplex stability. Reactive or non-reactive functionalities placed in these positions in accordance with this invention can best initiate cleavage and destruction of targeted RNA or interfere with its activity.

Reactive functionalities or pendant groups of oligonucleotides previously described in the literature have been almost exclusively placed on a phosphorus atom, the 5-position of thymine, and the 7-position of purines. A phosphorus atom attachment site can allow a reactive group to access both the major and minor grooves. However, internal phosphorus modification results in greatly reduced heteroduplex stability. Attachments at the 3' and/or 5' ends are limiting in that only one or two functional groups can be accommodated in the oligonucleotide. Functionality placed in the 5-position or 7-position of heterocycles (bases) pyrimidine and purine respectively will reside in the major groove of the duplex and will not be in proximity to the RNA 2'-hydroxyl substrate. Further such placement can interfere with Watson-Crick binding.

Pendant groups that do not possess reactive functionality but serve to enhance the pharmacodynamic

- 36 -

and pharmacokinetic properties of the oligonucleotides are also preferred for use in accordance with certain embodiments of this invention. Pharmacodynamic property improvement means, in this context, improved  
5 oligonucleotide uptake, enhanced oligonucleotide resistance to degradation, and/or strengthened sequence-specific hybridization with RNA. Pharmacokinetic property improvement means, in this context, improved  
10 oligonucleotide uptake, distribution, metabolism or excretion. Such pendant groups do not initiate chemical reactions. They preferably include alkyl chains, polyamines, ethylene glycols, polyamides, aminoalkyl chains, amphipathic moieties, points for reporter group attachment, and intercalators attached to any of the  
15 preferred sites for attachment.

It is believed that attaching RNA cleaving moieties in accordance with the foregoing considerations will permit those moieties to lie in the minor groove of the hybrid formed from the composition of the present invention and  
20 the messenger RNA for which modulation is desired. It is possible that other positions for attachment of the RNA cleaving moieties having a similar effect may be found, especially when further modifications of the purine or pyrimidine structure is undertaken as may be done by  
25 persons of ordinary skill in the art without deviating from the spirit of the present invention. Once again, it is to be understood that preferably one, or at most a few RNA cleaving moieties are generally to be employed. Thus, artisans in the field will have great latitude in  
30 selecting means of attachment of the RNA cleaving moieties, the pharmacodynamic improving group or the pharmacokinetic improving group in accordance with this invention.

The RNA cleaving moieties of the compositions of the  
35 present invention are designed in such a fashion that they can be effective in performing their proximate task, leading to the desired modulation of RNA activity. It

- 37 -

is believed to be useful to employ heteroatomic substitutions in the RNA cleaving moieties of these molecules, especially amides and polyamides, and indeed some may be preferred in order to ensure even tighter binding between  
5 the target mRNA and the compositions of the invention.

Oligonucleotide analogs particularly suited for the practice of one or more embodiments of the present invention comprise 2'-sugar modified oligonucleotides wherein one or more of the 2'-deoxy ribofuranosyl  
10 moieties of the nucleoside unit is modified with a hydrogen or hydroxyl, halo, azido, amino, methoxy or alkyl group. For example, the substitutions which may occur include H, OH, lower alkyl, substituted lower alkyl, F, Cl, Br, CN, CF<sub>3</sub>, OCF<sub>3</sub>, OCN, O-alkyl, S-alkyl,  
15 SOME, SO<sub>2</sub>Me, ONO<sub>2</sub>, NO<sub>2</sub>, N<sub>3</sub>, NH<sub>2</sub>, NH-alkyl, OCH<sub>2</sub>CH=CH<sub>2</sub>, OCH=CH<sub>2</sub>, OCH<sub>2</sub>CCH, OCCH, aralkyl, heteroaralkyl, heterocycloalkyl, aminoalkylamino, heterocycloalkylamino, polyalkylamino, substituted silyl, a RNA cleaving moiety or a group for improving the pharmacodynamic properties of  
20 an oligonucleotide or a group for improving the pharmacokinetic properties of an oligonucleotide where alkyl is a straight or branched chain of C<sub>1</sub> to C<sub>12</sub> with unsaturation within the carbon chain such as allyloxy.

These modified bases are linked together and to the  
25 rest of the oligonucleotide or oligonucleotide analog through a sugar linking group. The linking group may be any of those structures described herein which are capable of linking sugar moieties of oligonucleotides together to form the targeting portion of the  
30 compositions of this invention. It is preferred that these sugar linking groups comprise the phosphodiester structure or a derivative of such. Derivatives of the phosphodiester structure may include substitution of a sulphur, methyl, methyl oxide, or amine group for an  
35 oxygen. The sugar phosphate nucleic acid backbone may be modified as a phosphorothioate, methylphosphonate, or phosphate alkylated moiety. The phosphodiester linkage



- 38 -

may also be replaced by a carbon or ether linkage as discussed above.

In order to attach tethered functionality to the exocyclic 2-position of 2'-deoxyadenosines, 2'-deoxy-  
5 guanosines, and other purines and purine analogs, the following exemplary procedure can be utilized. 2,6-Dichloropurine (Aldrich Chemical Co.) is deoxyribosylated with 3,5-di-O-(4-methylbenzoyl)- $\alpha$ -D-erythro-pentofuranosyl chloride in accordance with the procedure  
10 set forth in Synthetic Procedures in Nucleic Acid Chemistry, Vol.1, p. 521, (1968) and subsequently aminated with methanolic ammonia according to the procedure of the Journal of American Chemical Society, Vol. 106, pg. 6379 (1984). The resulting 2-chloro-2-  
15 deoxyadenosine is converted into a series of 2-substituted 2-deoxyadenosines by nucleophilic displacement of the 2-chloro group with selected nucleophilic reactive functionalities. In this manner, a variety of substituted species such as amines, oxygens, sulfurs, and  
20 seleniums can be attached directly to the 2-position carbon of the bicyclic ring. Reactive functionalities attached to the 2-position of 2'-deoxyadenosine via a carbon atom can be obtained from 5-amino -1-(2-deoxy- $\beta$ -D-erythro-pentofuranosyl) imidazole 4-carboxamide and  
25 substituted aldehydes according to the published procedure of the Journal of American Chemical Society, Vol. 90 p. 4962, (1974) or from palladium-catalyzed cross coupling reactions as set forth in Nucleosides and Nucleotides, Vol. 8 p. 699, (1989) with 2-iodo-2'-  
30 deoxyadenosine. See U.S. Patent 4,719,295. The 2-reactive functionality 2'-deoxyadenosines can be deaminated with adenosine deaminase or nitrous acid to afford deoxyguanosine counterparts. As per selected examples below 2-substituted-2- deoxyguanosines and -  
35 2-adenosines can be converted to the 5'-di-methoxy-trityl-3'-cyanoethylphosphoramidites (N6-benzoyl for adenine types and N2-isobutryl for guanin types). These

- 39 -

are routinely inserted into oligonucleotides via automated, solid phase DNA synthesizers according to well-known procedures.

Reactive functionalities emanating from the 3-position of 2'-deoxyguanosine can be obtained by a multi-step synthesis starting with the alkylation of the methylene moiety of methyl 5-(cyanomethyl)-1-(2-deoxy-3,5-di-O-p-toluoyl- $\beta$ -D-erythropento-furanosyl) imidazole-4-carboxylate with halogenated reactive functionalities. The starting imidazole material can be synthesized according to a known procedure. Journal of Medicinal Chemistry, Vol. 27 p. 1389 (1984). The alkylated imidazoles are treated with methanolic ammonia to remove the toluoyl groups and to provide imidazole carboxamides. These are inserted into oligonucleotides as 5'-DMT-3'-cyanoethyl phosphoramidites through routine solid state synthetic techniques.

Following synthesis, the prepared oligonucleotides, which are in blocked form, are removed from the solid support upon which they are elaborated such as by ammonium hydroxide treatment. Further basic treatment deblocks and cyclizes the imidazole carboxamide into a 3-deaza-3-(reactive functionality)-guanine residue within the desired oligonucleotide sequence. Alternately, cleavage from the support with concentrate ammonium hydroxide directly cyclizes the imidazole carboxamide to the 3-deaza-guanine.

Alkylation of the imidazole active methylene with  $\alpha$ -halo- $\alpha$ - [reactive functionality] acetaldehyde dimethylketal or  $\alpha$ -halo- methyl-[reactive functionality] ketone with subsequent amination provides imidazole carboxamides. These can be converted to 5'-DMT-3'-cyanoethylphosphoramidites and inserted into sequence specific oligonucleotides. Basic treatment as above removes the oligonucleotides from the solid support and cyclizes the imidazole moiety to a 3-deaza-3- [reactive functionality]-guanine. Further cyclization between the

- 40 -

resulting N-2-exocyclic amine group and the aldehydic or ketonic carbonyl provides tricyclic heterocycle with reactive functionality, (pyrrolo[2,3- $\beta$ ]-imidazo[2,3- $\delta$ ]pyridin-2-one (5H)-7-(or 8)-[reactive functionality]).

Direct deoxyribosylation of 6-isobutryl pyrrolo[2,3- $\beta$ ]imidazo [2,3- $\delta$ ] pyridine 7-(or 8)-[reactive functionality] provides the 1-(2'- deoxy- $\beta$ -D-erythro-pentofuransyl) derivative after basic deblocking of the toluoyl groups. The tricyclic heterocycle can be obtained from the alkylation of the tetrahydropyranyl derivative of methyl 5-cyanomethylimidazole 4-carboxylate in accordance with the procedure of the Journal of Medicinal Chemistry, Vol. 21, p. 1212 (1978), with  $\alpha$ -halo- $\alpha$ -[reactive functionality]-acetaldehyde dimethylketals or  $\alpha$ -halomethyl-[reactive functionality] ketones with subsequent amination. Acid treatment removes the tetrahydropyranyl blocking group and reducing conditions provides the 7,8-di-hydro 7-(or 8)-[reactive functionality] tricyclic heterocycle. The dihydropyrrole ring nitrogen can be protected with an isobutryl group. The 5'-DMT-3'-phosphoramidite-6-isobutryl nucleoside can be inserted into sequence-specific oligonucleotides via standard automated synthesis. Oligonucleotides prepared in this manner contain a pyrrolo[2.3-b]imidazo[2,3-d]-pyridin-4-one(5H)-7-(or 8)- [reactive functionality], the 7,8-dihydro ring replacing a normal guanine residue.

Tether functionalities attached to the 3-position of 3-deaza-adenine modified oligonucleotides can be obtained such as by a multistep synthesis involving the attachment of reactive functionalities to the methylene moiety of the tetrahydropyranyl derivative of dimethyl 5-carboxymethylimidazole-4-carboxylates. See The Chemistry of Heterocyclic Compounds, A. Weissberger, Ed., Imidazole and Derivatives, Part 1, Interscience, N.Y. (1953). Reactive functionalities having suitable leaving groups such as halogens, sulfonates, trichloroacetimidates r

- 41 -

conjugated double bond systems are employed. The modified imidazoles can be aminated with ammonium/heat forming the 3-deaza-3-(reactive functionality)-1 (or3)-tetrahydropyranylxanthanines. These can be  
5 chlorinated with phosphoryl chloride to afford the 2,6-dichloro-3-deazapurines substituted in the 3-position. These compounds can be selectively aminated with ammonia, subjected to catalytic hydrogenolysis to remove the chlorine atom, and then protected with a  
10 conventional diphenylcarbamoyl group. Deoxyribosylation of these compounds under basic conditions with 3'5-di-O-toluoyl- $\alpha$ -D-erythro-pento-furanosyl chloride followed by selective deblocking of the sugar blocking groups provides 4-(di-phenylcarbamoyl)-1- (2-deoxy- $\beta$ -D-  
15 erythro-pentofuranosyl)-7-[reactive functionality] imidazo [4,5-d] pyridines. These can be converted in conventional fashion to their 5'-DMT-3'-cyanoethyl-phosphoramidites and incorporated into oligonucleotides.

In order to attach tethered functionality in the  
20 2'-position of ribo-oligonucleotides several exemplary synthetic procedures were devised.

**Synthesis 1.** Nucleophilic Displacement of 2'-Leaving Group in Arabino Purine Nucleosides.

Nucleophilic displacement of a leaving group in the  
25 2'-up position (2'-deoxy-2'-(leaving group)arabino sugar) of adenine or guanine or their analog nucleosides can be so employed. General synthetic procedures of this type have been described by M. Ikehara et al., Tetrahedron Vol. 34 pp. 1133-1138 (1978); *ibid.*, Vol. 31, pp. 1369-1372 (1975); Chemistry and Pharmaceutical Bulletin, Vol. 26, pp. 2449-2453 (1978); *ibid.*, Vol. 26, pp. 240-244 (1978); M. Ikehara Accounts of Chemical Research, Vol. 2, pp. 47-53 (1969); and R. Ranganathan Tetrahedron Letters, Vol. 15, pp. 1291-1294 (1977). Thus  $\beta$ -D-ara-  
30 binofuranosyl derivatives of guanine, adenine, cytosine, and thymine (Aldrich Chemical Co.) can be protected as the N2-isobutryl, N6-benzoyl, and N4-benzoyl, respec-

- 42 -

tively, by known procedures. Simultaneous protection of the 3',5'-hydroxyls of the arabinofuranosyl nucleosides can be accomplished with 1,3-dichloro-1,1,3,3-tetra-isopropyldisiloxane according to the known procedure set forth in Nucleic Acid Chemistry, Improved and New Synthetic Procedures, Methods, and Techniques, Part 3, page 229 (1986). The 2'-hydroxy group is activated towards nucleophilic displacement by treatment with trichloroacetonitrile and sodium hydride, Tetrahedron Letters., Vol. 23, pp. 409-412 (1982), or trifluoromethylsulfonic acid anhydride and sodium hydride. These leaving groups can be displaced with inversion by substituted hetero-atoms such as nitrogen, sulfur, oxygen, or selenium to provide tethered reactive functionality in the 2'-position of the 3',5'-silyl blocked nucleosides. These nucleosides can then be deblocked such as by fluoride ion and converted to their 5'-DMT-3'-cyanoethylphosphoramidites for insertion into oligo-nucleotides.

**Synthesis 2. Nucleophilic Displacement of 2,2'-Anhydro Pyrimidines.**

Nucleosides thymine, uracil, cytosine or their analogs are converted to 2'-substituted nucleosides by the intermediacy of 2,2'-cycloanhydro nucleoside as described by J.J. Fox, et al., Journal of Organic Chemistry, Vol. 29, pp. 558-564 (1964).

**Synthesis 3. 2'-Coupling Reactions.**

Appropriately 3',5'-sugar and base protected purine and pyrimidine nucleosides having a unprotected 2'-hydroxyl group are coupled with electrophilic reagents such as methyl iodide and diazomethane to provide the mixed sequences containing a 2'-OMe group as per the procedure of Inoue, et al., Nucleic Acids Research Vol. 15, pp. 6131-6148 (1987).

**Synthesis 4. 2-Deoxy-2-substituted Ribosylations.**

2-Substituted-2-deoxyribosylation of the appropriately protected nucleic acid bases and nucleic acids base

- 43 -

analogs has been reported by E. T. Jarvi, et al., Nucleosides & Nucleotides Vol. 8, pp. 1111-1114 (1989) and L. W. Hertel, et al., Journal of Organic Chemistry Vol. 53, pp. 2406-2409 (1988).

5     **Synthesis 5. Enzymatic Synthesis of 2'-Deoxy-2'-Substituted Nucleosides.**

The 2-Deoxy-2-substituted glycosyl transfer from one nucleoside to another with the aid of pyrimidine and purine ribo or deoxyribo phosphorolyses has been  
10 described by J. R. Rideout and T. A. Krenitsky, United States Patent # 4,381,344 (1983).

**Synthesis 6. Conversion of 2'-Substituents Into New Substituents.**

2'-Substituted-2'-deoxynucleosides are converted into  
15 new substituents via standard chemical manipulations. For example, S. Chladek et al., Journal of Carbohydrates, Nucleosides & Nucleotides Vol. 7, pp. 63-75 (1980) describes the conversion of 2'-deoxy-2'- azidoadenosine, prepared from arabinofuranosyladenine, into 2'-deoxy-2'-  
20 aminoadenosine.

**Synthesis 7. Free Radical Reactions.**

Conversions of halogen substituted nucleosides into 2'-deoxy- 2'-substituted nucleosides via free radical reactions has been described by K. E. B. Parkes and K.  
25 Taylor, Tetrahedron Letters Vol. 29, pp. 2995-2996 (1988).

**Synthesis 8. Conversion of Ribonucleosides to 2'-Deoxy-2'- Substituted Nucleoside.**

Appropriately 3',5'-sugar and base protected purine  
30 and pyrimidine nucleosides having a unprotected 2'-hydroxyl group are converted to 2'-deoxy-2'- substituted nucleosides by the process of oxidation to the 2'-keto group, reaction with nucleophilic reagents, and finally 2'-deoxygenation. Procedures of this type have  
35 been described by F. De las Heras, et al., Tetrahedron Letters Vol. 29, pp. 941-944 (1988).

- 44 -

Functionality in the 1'-position of a 2'-deoxyribo-  
furanosyl moiety can be obtained from the nucleoside  
antibiotic, psicofuranine, which differs from adenosine  
by having a hydroxyl methyl group in the 1'-position.  
5 The 3',5'- hydroxyls can be protected by 1,3-dichloro-  
1,1,3,3-tetra-isopropyldisiloxane, followed by protection  
of the 1'-hydroxymethyl with a triphenylmethyl group.  
The 2'-hydroxy group can be deoxygenated by formation of  
the thiocarbonate and subsequent treatment with  
10 tri-n-butyl hydride according to known procedures. See,  
J. Amer. Chem. Soc., Vol. 105, pp. 4059-4065 (1983). The  
N6-amino group can be benzoylated, the trityl group  
removed with acid, and the resulting primary hydroxy  
group converted to the trichloroacetamidate with sodium  
15 hydride and trichloroacetonitrile. Displacement  
reactions with various tether/ reactive functionality can  
be readily accomplished in the next step. The disilyl  
blocking group can be removed with fluoride ion and the  
resulting 2'-deoxy nucleoside converted to the 5'-DMT-3'-  
20 phosphoramidite for insertion into oligonucleotides via  
known procedures.

Reactive functionality in the 4'-position can be  
obtained from 2'-deoxynucleotides (Sigma Chemical  
Company) by selectivity protecting the 5'-hydroxyl with  
25 the t-butyldimethyl silyl group and the 3'-hydroxyl with  
the tetrahydropyranyl group. The silyl group is removed  
with fluoride ion. The Pfitzner-Moffatt oxidation  
procedure, Journal of American Chemical Society, Vol. 85  
p. 3027 (1963), can be used to provide the 5'-aldehydes  
30 which are then treated with formaldehyde affording the  
4'-hydroxymethyl derivatives. Tetrahedron Letters, 435  
(1977). These deoxynucleosides, as their 5'-trityls, can  
be converted to the 4'- trichloroacetamides with  
trichloroacetonitrile and sodium hydride. Reactive  
35 functionality are then placed in the 4'-position via  
nucleophilic displacements of the acetamidates. Acid  
treatment removes the blocking groups in the 5' and

- 45 -

3'-positions. These can be subsequently prepared as the 5'-DMT-3'-cyanoethylphosphoramidite and inserted into modified oligonucleotides via a automated, solid phase DNA synthesizers.

5 The synthesis of abbreviated guanosine nucleosides (monocyclics) substituted with reactive functionality such as 2'-deoxy-2-imidazolone-4-carboxamide 5-reactive  
10 reactivity can be prepared from the reaction of ethyl 2-bromomethyl-2-imidazolone (1H,3H)-4-carboxylate with reactive functionality nucleophiles such as histamine, hydroxyethylimidazolyl, and mercaptoethylimidazolyl. Other heteroatomic nucleophiles, neutral or anionic, with attached reactive functionality may be employed as well. These compounds are protected as the 2-o-diphenyl-  
15 carbamoyl derivatives with diphenylcarbamoyl chloride and base and 2'-deoxyribosylated with 3,5-di-o-(4-methyl-benzoyl)- $\alpha$ -D-erythro-pentofuranosyl chloride. The 3',5'-dihydroxys of the sugar are selectively deblocked with base and reprotected as the 5'-DMT-3'-cyanoethyl-  
20 phosphoramidite. These imidazole nucleosides can be inserted into oligonucleotides according to known procedures. The strong basic conditions required to remove the acyl blocking groups on the cytosine, adenine, guanine, and imidazole bases convert the 4-imidazole  
25 carboxylate to the 4-carboxamide moiety.

Reactive functionality in the 5-position of 4-amino-imidazol-2-one modified oligonucleotides can be prepared by the deoxyribosylation of 2,5-dibromo-4-nitroimidazole, Journal of Chemical Society Vol. 121, p. 947 (1922), with  
30 3,5-O-(p-methyl-benzoyl)- $\alpha$ -D-erythro-pentofuranosyl chloride under basic conditions. Selective displacement of the 5-bromo group with nucleophilic reactive functionality afforded a variety of bromonitro-deoxy-nucleotides. Displacement of the 2-bromo group and  
35 subsequent reduction of the 4-nitro group provided 4-amino-1-(2-d oxy- $\beta$ -D -erythro-pentofuranosyl)imidazol-



- 46 -

2-one(3H) which can be inserted into oligonucleotides via their 5'-DMT-3'-cyanoethylphosphoramidites.

Reactive functionality placed in the 3-position of 2-aza-3-deaza-2'-deoxyadenosine can be prepared from the deoxyribosylation of 4,7-dichloroimidazo[4,5-d]-pyridazine, J. Org. Chem., Vol. 23 p. 1534 (1958) with 3,5-di-O-(methylbenzoyl)- $\alpha$ -D-erythropentofuranosyl chloride under basic conditions. This is aminated with liquid ammonia to provide 4-amino-7-chloro-1-(2-deoxy- $\beta$ -D-erythro-pento furanosyl)imidazo[4,5-d]pyridazine. Reactive functionality are then inserted in the 7-position by nucleophilic displacement of the chlorine atom. Examples are heteroatoms such as nitrogen, oxygen, sulfur, selenium, etc., attached to the reactive functionality placed on the 7-carbon position. Protection of 4-amino-1-(2-deoxy- $\beta$ -D-erythropentofuranosyl)-7-chloroimidazo-[4,5-d] pyridazine with t-butyldimethylsilyl groups followed by displacement of the chlorine atom with triphenylphosphorane reagents affords carbon-linked reactive functionality. The 7-position tethered reactive functionality 4-amino-1-(2-deoxy- $\beta$ -D-erythropentofuranosyl-7-chloroimidazo[4,5-d]-pyridazines are then inserted, via 5'-DMT-3'-phosphoramidites, into oligonucleotides.

In order to attach reactive functionality in the 2'-position of  $\sigma$ -arabinofuranosyloligonucleotides, silylated heterocycles can be glycosylated with 1-O-acetyl-2,3,5-tri-O-benzoyl-D-arabinofuranose (Pfanstiehl Laboratories, Inc.) with Lewis acid catalysis. The benzoyl groups of the sugar of the resulting  $\alpha$ -nucleoside are selectively removed with methanolic ammonia; the 3',5'-hydroxyls are then reprotected with 1,1,3,3-tetraisopropylidisiloxane. Reactive functionality is attached to the 2'-hydroxyl group via the Mitsunobu reaction. Synthesis 1 (1981). Deprotection with fluoride ion then affords the 2'-[tethered reactive functionality]-1-( $\alpha$ -D-arabino-

- 47 -

furanosyl) nucleosides, which can be inserted into oligonucleotides via their 5'-DMT-3'-cyanoethylphosphoramidites.

Functionality may be attached to the 6'-position of the natural occurring nucleosides or base analogs thereof by direct nucleophilic attack of protected heterocycles on carbocyclic 3,5-di-O-benzyl-2-deoxy- $\alpha$ -1,6-epoxy-D-erythropentofuranose as set forth by K. Biggadike, et al., J. Chem. Soc., Chem. Commun., 1083 (1987). Reactions of this type will afford carbocyclic 2'-deoxynucleosides with a hydroxyl group located at the 6'-position on the  $\alpha$  face of the cyclopentane ring. Functionality can now be attached to the unprotected hydroxyl group by direct coupling (i.e., Mitsunobu conditions, triphenylphosphine/DEAD) or double displacement reactions. Debenzylation of compound of this type, followed by formation of the 5'-DMT and the 3'- $\beta$ -cyanoethoxy diisopropyl-phosphoramidite by standard procedures provides monomers which can be inserted into oligonucleotides by solid phase, automated DNA synthesizers.

Pyrimidine C-nucleoside type compounds which have the deoxyribofuranosyl moiety located at the 4-position have the ring nitrogen atom in the 3-position available to attach functionality. These type nucleosides may be synthesized by a displacement, via a 4-carbanion of the protected pyrimidine, of the chlorine atom of 3,5-di-O-toluoyl-2-deoxy- $\alpha$ -erythropentofuranosyl chloride. G. D. Daves and C.C. Cheng, Progress in Medicinal Chemistry, Vol. 14, pp. 304-349 (1978). Appropriate functionality can be placed in the 3-position of the protected pyrimidine by commonly known procedures. Selective removal of the toluoyl groups and subsequent DMT and phosphitylation will provide desired monomers for insertion into oligonucleotides.

Quinazoline deoxyribosides with functionality located in the 8-position may be obtained by palladium mediated coupling of furanid glycals with 7-tributylstannylated

- 48 -

heterocycles as set forth by R. O. Outten and G. D. Daves, Jr. in the Journal of Organic Chemistry, Vol. 52, p. 5064 (1987).

Without desiring to be bound by any particular theory  
5 of operation, it is believed that the reactive RNA  
cleaving functionalities, described in this invention  
work by mechanisms involving either:

1. phosphodiester bond cleavage via acid/base catalysis;
- 10 2. backbone sugar cleavage;
3. base alkylation cleavage; or
4. sugar alkylation, ie. 2'-hydroxyl, cross-linking.


One important aspect of this invention is the position  
and orientation of an appropriate reactive functionality  
15 minor groove or side formed between the targeting portion  
of this invention and the target RNA. A summary of this  
is provided in Table 1, and each reactive functionality  
is described according to the suggested mechanism of  
action.

Table 1

EXAMPLE NUMBER	REACTION	MECHANISM	PREFERRED FUNCTIONALITY	PREFERRED LOCATION
5	1 Phosphodiester bond cleavage	2' proton abstraction on ribose sugar	Organic base	Minor groove
2	Phosphodiester bond cleavage	2' proton abstraction on ribose sugar	Coordination complex	Minor groove
3	Phosphodiester bond cleavage	5' proton donation on ribose sugar	Organic acid	Either
10	4 Phosphodiester bond cleavage	Electrophilic activation of phosphate	Coordination complex with net positive charge	Either
5	Oxidative cleavage of ribose sugar	Oxygen radical generation & hydrogen abstraction from ribose	Organic radical generating moiety	Minor groove
15	6 Oxidative cleavage of ribose sugar	Oxygen radical generation & hydrogen abstraction from ribose	Coordination complex	Minor groove
20	7 Electrophilic attack on bases	Base removal	Alkylating agent	Minor groove
	8 Electrophilic attack on 2'-hydroxyl group of RNA	Covalent cross-link	Alkylating agent	Minor groove

- 50 -

Phosphodiester bond cleavage can be accomplished by strategically positioning either proton accepting, proton donating, or electron accepting functional groups, represented by X,Y, and Z respectively in Scheme 1 adjacent to such phosphodiester bonds.



SCHEME 1

Where B<sub>1</sub> and B<sub>2</sub> are base units. In some applications, one of the chemical groups is sufficient to catalyze RNA cleavage. However, in other applications of the invention, the combination of two or even three groups may be preferred. Artisans in the field will have great latitude in selecting the specific reactive functionalities X,Y, and/or Z. There is also great latitude in the election to use one or more reactive functionalities in the same molecule.

Examples of proton accepting functionalities which can form the phosphodiester bond cleaving reactive moiety in accordance with this invention include heteroatomic bases which are capable of base-catalysis of the phosphodiester bond. Ammoniacal species generally, such as alkyl and arylamines and alkyl and arylhydrazines, are preferred. Heterocyclic nitrogenous compositions, especially imidazoles, pyridines, azines, oxazoles, thiazoles, and the like are more preferred. Of these species, the 1-, 2-, and 5-imidazoles are most preferred. Each of these compositions may be widely substituted in order to vary the electronegativity of the various nitrogen atoms

- 51 -

available for base catalysis. Alkyl and aryl amidium species are also most preferred. Carboxylic acid and their derivatives of the above ammoniacal species are most preferred.

- 5 Another type of species which may be useful as the proton accepting function in the compositions and methods of the present invention are coordination complexes. These are represented as Example 2 of Table 1. Coordination complexes are useful in this regard by  
10 having a metal ion with ligands designed so that not all of the coordination sites on the metal are occupied by the ligands, thus leaving one or more empty sites. In solution, the empty sites are occupied by water molecules leading to -OH moieties being coordinated to the metal.  
15 The -OH group can function as a base to abstract the 2' proton and effect phosphodiester bond cleavage. The selection of ligands for coordination with the metal ions should follow three important considerations. First, they should be capable of holding the metal near the  
20 target RNA and "steer" the hydroxyl group on the coordination complex (which can be represented as HO-ML) near to the 2' proton which is to be abstracted. The ligands are designed to interact favorably with the duplex formed between the compositions of the invention  
25 and the target RNA.

A second important function of the ligands is to regulate the pKa of the hydrolyzed metal. The coordination complex, HO-ML, can function to catalyze the cleavage of the target RNA not only by providing a  
30 hydroxyl group in the proper orientation for proton abstraction, but at the proper pKa. The proper pKa depends upon the pH which exists during the practice of the methods of the invention. The proper pKa to be maintained is preferably such that the hydroxyl group is  
35 sufficiently ionized to allow for proton abstraction, but not so low that the hydroxyl group has no tendency to abstract the proton. The ligands are preferably selected in accordance with this invention as to adjust the pKa of the hydrolyzed metal complex to achieve these ends.

- 52 -

The third important function of the ligands is to regulate redox reactions. In the case of metal ions that have two or more accessible oxidation states, the ligands stabilize one oxidation state in preference to others and  
5 prevent or retard the generation of diffusible, toxic, usually oxygen-containing, radicals.

Other coordination complexes, including some which are intended to generate diffusible oxygen-containing radicals, can be used in accordance with other  
10 embodiments of this invention, but these are believed to cleave the target RNA by a different mechanism described hereinafter.

The metal ions preferred for use in this invention include  $\text{Ca}^{+2}$ ,  $\text{Sc}^{+3}$ ,  $\text{Cr}^{+3}$ ,  $\text{Mn}^{+2}$ ,  $\text{Fe}^{+3/+2}$ ,  $\text{Co}^{+2/+1}$ ,  $\text{Zn}^{+2}$ ,  $\text{Al}^{+3}$ ,  
15  $\text{Ga}^{+3}$ ,  $\text{Rh}^{+3}$ ,  $\text{Mg}^{+2}$ . Ligands include multidentate carboxylates, aminos, hydroxamic acids, catecholates, heterocyclic amines and peptides. Ligands of these types include, but are not limited to, EDTA, NTA, bipyridyl, phenanthroline, desferrioxamine, enterobactin (and its  
20 analogs), gly gly his and gly gly gly.

These reactive functionalities are attached to the balance of the molecule comprising the compositions of the invention in any convenient way, usually through an amide function. Persons of ordinary skill in the art  
25 will have no difficulty in selecting appropriate means for the attachment of the reactive functionalities to the targeting portions of the compositions of these invention through convenient tether means.

In accordance with other embodiments of the invention,  
30 proton donating, acidic media may be employed as reactive functionalities in order to effect acid catalysis of the hydrolysis of the phosphodiester bonds. Such functionalities are depicted as group Y of Scheme 1 and particular embodiments are presented in Example 3 of  
35 Table 1. Any of a large number of acids are believed to be effective in donating protons for the acid catalyzed hydrolysis of phosphodiester bonds in accordance with this invention. Such acid functions include carboxylic acids and heterocyclic acids.

- 53 -

Among the class of heterocyclic acidic functionalities tetrazoles and triazoles are preferred. It is known that several of the nitrogen atoms in the triazoles and tetrazoles possess substantial acidity. It is believed  
5 that they may be used for catalysis of such hydrolysis reactions. The acidic functionalities are attached to the targeting portion of the compositions, as are all of the reactive species in accordance with this invention.

It is believed that the hydrolysis of phosphodiester  
10 bonds in nucleotides proceeds through a pentacoordinate structure on the phosphorus atom having trigonal bipyrimidal geometry and a charge of -2. The presence of an electropositive species which can stabilize this transition state is believed to assist, catalytically, in  
15 the cleavage reaction. Such species may be considered to be an electropositive moiety and is denominated as group Z in Scheme 1. Examples of such functionalities are presented in Example 4 in Table 1. These reactive functionalities may comprise a coordination complex  
20 having two empty sites. The two empty sites are believed to coordinate with and stabilize the phosphorus transition state. Among the reactive functionalities which may be useful for this purpose are those containing metal ions such as:  $\text{Ca}^{+2}$ ,  $\text{Sc}^{+3}$ ,  $\text{Cr}^{+3}$ ,  $\text{Mn}^{+2}$ ,  $\text{Fe}^{+3/+2}$ ,  $\text{Co}^{+2/+1}$ ,  
25  $\text{Zn}^{+2}$ ,  $\text{Al}^{+3}$ ,  $\text{Ga}^{+3}$ ,  $\text{Rh}^{+3}$ ,  $\text{Mg}^{+2}$ . Ligands which are suitable for the formation of such coordination complexes include multidentate carboxylates, aminos, hydroxamic acids catecholates, heterocyclic amines, and peptides.

Radical forming moieties, especially those which form  
30 oxygen radicals, may also be employed as the reactive functionalities of the compositions of this invention. When radical generating species are used as the reactive portions of the compositions of the present invention, it is believed to be necessary that they be delivered into  
35 the minor groove of the complex formed between the compositions and the target RNA. Minor groove delivery is believed to be essential to their effective operation in high yield without adverse, and potentially toxic side effects. It is known that prior employment of radical



- 54 -

generating species has been met by migration of toxic compounds either comprising or related to the radicals formed, with toxicity and damage to other intracellular species resulting. It is believed that minor groove  
5 delivery of these species obviates or at least substantially diminishes these negative effects. A large number of organic radical generating species such as the quinones, coumarins, and other agents which are capable of producing active oxygen may be employed.

10 Coordination complexes may also be employed as radical-generating agents in the minor groove. It is known that certain metals which have more than one accessible oxidation state, when complexed by appropriate ligands and in the presence of reducing agents and  
15 oxygen, will generate oxygen radical species capable of cleaving oligonucleic acid targets. As with the organic radical generating species, it is necessary to deliver the coordination complex into the minor groove of the hybrid formed between the compositions and the target  
20 RNA.

Coordination complexes useful in generating radical species include complexes of iron, copper, and molybdenum with ligands such as carboxylic acids, amines, heterocyclic nitrogen containing species such as phenanthro-  
25 lines, dipyridyl, imidazole, or peptidyl ligands such as gly gly his, gly gly his lys, and gly gly gly gly, and hydroxamic acids. These ligands may be mixed to generate more effective combinations within the scope of this invention.

30 A further group of moieties which may serve as reactive portions of the compositions of the present invention are alkylating agents. Several classes of alkylating agents are known to persons of ordinary skill in the art including the N-mustard types which are  
35 generally  $\beta,\beta$ -haloethylamines. Other alkylating agents such as  $\alpha$ -haloketones, sulfonyl halides, aziridines, allylic halides, epoxides, alkyl/aryl halides, and light activatable alkylating agents such as psoralines may also

- 55 -

be employed. Michael reaction accepting compositions may also find utility in the practice of this invention.

A 2'-hydroxyl group of the targeted RNA, exposed in the minor groove or side, serves as a nucleophilic center for bond formation with tethered electrophilic functionality that also resides in the minor groove of the heteroduplex. Classes of electrophiles for attachment appropriately to nucleotides are listed above. Targeted RNA is inactivated by formation of covalent links between a modified oligonucleotide and the RNA 2'-hydroxyl group.

It has now been discovered from structure activity relationships studies that a significant increase in binding ( $T_m$ )s of certain 2'-sugar modified oligonucleotides to its RNA target (complement) is correlated with an increased "A" type conformation of the heteroduplex. Furthermore, absolute fidelity of the modified oligonucleotides is maintained. The increased binding of our 2'-sugar modified sequence-specific oligonucleotides provides superior potency and specificity compared to phosphorus modified antisense oligonucleotides such as methyl phosphonates, phosphorothioates, phosphate triesters and phosphoramidites as known in the literature.

The only structural difference between DNA and RNA duplexes is an hydrogen atom in the 2'-position of the DNA ribofuranosyl moieties versus a hydroxyl group in the 2'-position of the RNA ribofuranosyl moieties (assuming that the presence or absence of a methyl group in the uracil ring system has no effect). However, gross conformational differences exist between DNA and RNA duplexes.

It is known from X-ray diffraction analysis of nucleic acid fibers, Arnott and Hukins, Biochemical and Biophysical Research Communication, Vol. 47, pp. 1504-1510 (1970), and analysis of crystals of double-stranded nucleic acids that DNA takes a "B" form structure and that RNA only takes the much more rigid "A" form structure. The difference between the sugar puckering

- 56 -

(C2' endo for "B"- form DNA and C3' endo for "A"-form RNA) of the nucleoside monomeric units of DNA and RNA is the major conformational difference between double-stranded nucleic acids.

5 The primary contributor to the pentofuranosyl moiety conformation is the nature of the substituent in the 2'-position. Thus, the population of the C3'-endo form increases with respect to the C2'-endo as the electro-  
negativity of the 2'-substituent increases. For example,  
10 among 2'-deoxy-2'-halo-adenine nucleosides, the 2'-fluoro derivative exhibits the largest population (65%) of C3'-endo, and the 2'-iodo shows the lowest (7%). Those of the adenosine (2'-OH) and deoxyadenosine (2'-H) are 36% and 19%, respectively. Furthermore, the effect of the  
15 2'-fluoro group of adenine dinucleotides (2'-deoxy-2'-fluoroadenosine-2'-deoxy-2'-fluoroadenosine or uridine) is further correlated to the stabilization of the stacked conformations more than ribo or deoxyribo modified dimers. Research indicates that the dinucleosides  
20 phosphates have a stacked conformation with a geometry similar to that of A-A but with a greater extent of base-base overlapping than A-A. It was assumed that the highly polar nature of the C2'-F bond and the extreme preference for C3'-endo puckering may stabilize the  
25 stacked conformation in an "A" structure.

Data from UV hypochromicity, circular dichromism, and  
'H NMR also indicate that the degree of stacking decreases as the electronegativeness of halogen decreases. Furthermore, a sterical bulkiness in the  
30 2'-position is better accommodated in an "A" form duplex than a "B" form duplex.

Thus, a 2'-substituent on the 3'-nucleotidyl unit of a dinucleoside monophosphate is thought to exert a number of effects on the stacking conformation: steric  
35 repulsion, furanose puckering preference, electrostatic repulsion, hydrophobic attraction, and hydrogen bonding capabilities. These substituent effects are thought to be determined by the molecular size, electronegativity, and hydrophobicity of the substituent.

- 57 -

The 2'-iodo substituted nucleosides possess the lowest C3'-endo population (7%) of the halogen series. Thus, on steric effects alone, one would predict an 2'-iodo or similar groups would contribute stacking destabilizing properties and thus reduced binding ( $T_m$ )s for antisense oligonucleotides. However, the lower electronegativeness and high hydrophobic attractive forces of the iodine atom and similar groups complicates the ability to predict stacking stabilities and binding strengths.

10 Studies with the 2'-OMe modification of 2'-deoxy guanosine, cytidine, and uridine dinucleoside phosphates exhibit enhanced stacking effects with respect to the corresponding unmethylated species (2'-OH). In this case, the hydrophobic attractive forces of the methyl  
15 group tend to overcome the destabilizing effects of its sterical bulkiness (hindrance).

2'-Fluoro-2'-deoxyadenosine has been determined to have an unusually high population of 3'-endo puckering among nucleosides. Adenosine, 2'-deoxyadenosine, and  
20 other derivatives typically have population below 40% in the 3'-endo conformer. It is known that a nucleoside residue in well-stacked oligonucleotides favors 3'-endo ribofuranose puckering.

Melting temperatures (complementary binding) are  
25 increased with the 2'-substituted adenosine diphosphates. It is not clear whether the 3'-endo preference of the conformation or the presence of the substituent is responsible for the increased binding. However, as noted, greater overlap of adjacent bases (stacking) can be  
30 achieved with the 3'-endo conformations.

The present novel approach to obtaining stronger binding is to prepare antisense RNA mimics to bind to the targeted RNA. Therefore, a random structure-activity relationship approach was undertaken to discover nuclease  
35 resistant antisense oligonucleotides that maintained appropriate hybridization properties.

A series of 2'-deoxy-2'-modified nucleosides of adenine, guanine, cytosine, thymidine and certain analogs of these bases have been prepared and have been inserted

- 58 -

as the modified nucleosides into sequence-specific oligonucleotides via solid phase nucleic acid synthesis. The novel antisense oligonucleotides were assayed for their ability to resist degradation by nucleases and to  
5 possess hybridization properties comparable to the unmodified parent oligonucleotide. Initially, small electronegative atoms or groups were selected because these type are not likely to sterically interfere with required Watson-Crick base pair hydrogen bonding  
10 (hybridization). However, electronic changes due to the electronegativeness of the atom or group in the 2'-position may profoundly effect the sugar conformation. During our structure activity relationship studies we discovered that the sugar modified oligonucleotides  
15 hybridized to the targeted RNA stronger than the unmodified (2'-deoxyribosyl type).

In further embodiment of this invention, a linking moiety has been devised to allow the direct attachment of a modified unit to the terminal position of the 3'-end of  
20 the modified oligonucleotides. Thus, an ester, or more preferably a bromomethylketo group, is attached to the 3'-hydroxyl of a modified 2'-modified nucleoside having its 5'-hydroxyl protected with a dimethoxytriphenylmethyl group and the heterocycle benzoylester of the cytosine  
25 series. If the required targeting sequence has a terminal 3'-thymine or cytosine base, the desired modified thymine or cytosine base containing the bromomethylketo linker is utilized as the first monomer to attach to the control pore glass (CPG) solid support  
30 which contains a normal nucleoside attached via its 3'-hydroxyl group. The base sensitive ester linkage attaching the 2'-modified nucleoside to the nucleoside attached to the CPG is cleaved under the usual concentrated ammonium hydroxide conditions that are  
35 utilized to remove the oligonucleotide from the CPG support. This will allow the modified oligonucleotide to have a 2'-modified unit at its terminal, 3'-end.

Cleavage of oligonucleotides by nucleolytic enzymes require the formation of an enzyme-substrate complex, or

- 59 -

in particular a nuclease-oligonucleotide complex. The nuclease enzymes will generally require specific binding sites located on the oligonucleotides for appropriate attachment. If the oligonucleotide binding sites are removed or hindered such that the nucleases will not attach to the oligonucleotides, the nuclease resistant oligonucleotides result. In the case of restriction endonucleases that cleave sequence-specific palindromic double-stranded DNA, certain binding sites such as the ring nitrogen in the 3- and 7-positions have been identified as required binding sites. Removal of one or more of these sites or hindering the nuclease approach to these particular positions within the recognition sequence has provided various levels of resistance to the specific nucleases.

The oligonucleotides or oligonucleotide analogs of this invention can be used in diagnostics, therapeutics, and as research reagents and kits. For therapeutic use the oligonucleotide is administered to an animal suffering from a disease affected by some protein. It is preferred to administer to patients suspected of suffering from such a disease with amounts of oligonucleotide which are effective to reduce the symptomology of that disease. It is within the scope of a person's skill in the art to determine optimum dosages and treatment schedules for such treatment regimens.

It is generally preferred to apply the therapeutic agents in accordance with this invention internally such as orally, intravenously, or intramuscularly. Other forms of administration, such as transdermally, topically, or intralesionally may also be useful. Inclusion in suppositories may also be useful. Use of pharmacologically acceptable carriers is also preferred for some embodiments.

The following procedures and examples illustrate the practice of this invention. These procedures and examples are not to be construed as limiting the invention.

- 60 -

Once nucleotides of the invention have been prepared, they can then subsequently be incorporated into oligonucleotides of the invention.

Oligonucleotides of the invention are synthesized by  
5 the standard solid phase, automated nucleic acid synthesizer such as the Applied Biosystems, Incorporated 380B or MilliGen/Bioscience 7500 or 8800. Triester, phosphoramidite, or hydrogen phosphonate coupling chemistries, M. Caruthers, Oligonucleotides. Antisense  
10 Inhibitors of Gene Expression., pp. 7-24, J. S. Cohen, ed. (CRC Press, Inc. Boca Raton, Florida, 1989), are used in with these synthesizers to provide the desired oligonucleotides. The Beaucage reagent, Journal of  
15 American Chemical Society, Vol. 112, pp. 1253-1255 (1990) or elemental sulfur, S. Beaucage et al., Tetrahedron Letters, Vol. 22, pp. 1859-1862 (1981), is used with phosphoramidite or hydrogen phosphonate chemistries to provide substituted phosphorothioate oligonucleotides.

In preparing certain of the compounds of the invention  
20 fugitive masking groups have been used. Such masking groups allow for ease of synthesis of the compounds. The masking groups are subsequently converted to the desired functionality. Such conversion preferably occurs during a standard deblocking step for a later reaction. As an  
25 example of use of this procedure is the use of phthalimide group for the introduction of an amino functionality. Alkyl phthalimides are attached at the proper position in a compound of interest, as for example a nucleoside, via a suitable intermediate such as an N-  
30 (haloalkyl)phthalimide. Upon completion of the synthesis of the compound of interest it is then used as a structural nucleotide for oligonucleotide synthesis utilizing standard oligonucleotide synthetic techniques on a nucleotide synthesizer. After the desired  
35 oligonucleotide is completed, it is cleaved from the synthesizer support and in doing so the cleaving reagent also converts the alkylphthalimide to the desired alkylamine. The above procedure can be expanded to attach longer chain polyamino functionalities to th

- 61 -

oligonucleotides of the invention. Nucleotides or oligonucleotides having a first alkylamino functionality are treated with a further N-(haloalkyl) phthalimide. The extended functionality is then treated to yield the terminal amine group. This can be repeated to further extend the polyamino functionality as desired. Alternately, the extended polyamino functionality is first synthesized and reacted with the first alkylamino functionality to form the polyamino functionality.

10 **PROCEDURE 1 - Hybridization Analysis.**

**A. Evaluation of the thermodynamics of hybridization of modified oligonucleotides.**

The ability of the modified oligonucleotides of the invention to hybridize to their complementary RNA or DNA sequences was determined by thermal melting analysis. The RNA complement was synthesized from T7 RNA polymerase and a template-promoter of DNA synthesized with an Applied Biosystems, Inc. 380B nucleic acid synthesizer. The RNA species was purified by ion exchange using FPLC (LKB Pharmacia, Inc.). Natural antisense oligonucleotides or those containing the modifications at specific locations were added to either the RNA or DNA complement at stoichiometric concentrations and the absorbance (260 nm) hyperchromicity upon duplex to random coil transition was monitored using a Gilford Response II spectrophotometer. These measurements were performed in a buffer of 10 mM Na-phosphate, pH 7.4, 0.1 mM EDTA, and NaCl to yield an ionic strength of 10 either 0.1 M or 1.0 M. Data was analyzed by a graphic representation of  $1/T_m$  vs  $\ln[Ct]$ , where  $[Ct]$  was the total oligonucleotide concentration. From this analysis the thermodynamic parameters were determined. Based upon the information gained concerning the stability of the duplex of heteroduplex formed, the placement of modified pyrimidine into oligonucleotides were assessed for their effects on helix stability. Modifications that drastically alter the stability of the hybrid exhibit reductions in the free



- 62 -

energy ( $\Delta G$ ) and decisions concerning their usefulness as antisense oligonucleotides were made.

**B. Fidelity of hybridization of modified oligonucleotides**

5 The ability of the modified antisense oligonucleotides of the invention to hybridize with absolute specificity to the targeted mRNA was shown by Northern blot analysis of purified target mRNA in the presence of total cellular RNA. Target mRNA was synthesized from a vector containing  
10 the cDNA for the target mRNA located downstream from a T7 RNA polymerase promoter. Synthesized mRNA was electrophoresed in an agarose gel and transferred to a suitable support membrane (i.e. nitrocellulose). The support membrane was blocked and probed using [ $^{32}\text{P}$ ]-labeled  
15 antisense oligonucleotides. The stringency was determined by replicate blots and washing in either elevated temperatures or decreased ionic strength of the wash buffer. Autoradiography was performed to assess the presence of heteroduplex formation and the autoradiogram  
20 quantitated by laser densitometry (LKB Pharmacia, Inc.). The specificity of hybrid formation was determined by isolation of total cellular RNA by standard techniques and its analysis by agarose electrophoresis, membrane transfer and probing with the labeled modified oligo-  
25 nucleotides. Stringency was predetermined for the unmodified antisense oligonucleotides and the conditions used such that only the specifically targeted mRNA was capable of forming a heteroduplex with the 2'-modified oligonucleotide.

- 63 -

## PROCEDURE 2 - Nuclease Resistance

## A. Evaluation of the resistance of modified oligonucleotides to serum and cytoplasmic nucleases.

5 Natural phosphorothioate, and modified oligonucleotides of the invention were assessed for their resistance to serum nucleases by incubation of the oligonucleotides in media containing various concentrations of fetal calf serum or adult human serum.

10 Labeled oligonucleotides were incubated for various times, treated with protease K and then analyzed by gel electrophoresis on 20% polyacrylamide-urea denaturing gels and subsequent autoradiography. Autoradiograms were quantitated by laser densitometry. Based upon the

15 location of the modifications and the known length of the oligonucleotide it was possible to determine the effect on nuclease degradation by the particular modification. For the cytoplasmic nucleases, a HL60 cell line was used. A post-mitochondrial supernatant was prepared by

20 differential centrifugation and the labeled oligonucleotides were incubated in this supernatant for various times. Following the incubation, oligonucleotides were assessed for degradation as outlined above for serum nucleolytic degradation.

25 Autoradiography results were quantitated for comparison of the unmodified, the phosphorothioates, and the modified oligonucleotides.

## B. Evaluation of the resistance of modified oligonucleotides to specific endo- and

30 exo-nucleases.

Evaluation of the resistance of natural and 2'-modified oligonucleotides to specific nucleases (ie, endonucleases, 3',5'-exo-, and 5',3'-exonucleases) was done to determine the exact effect of the modifications

35 on degradation. Modified oligonucleotides were incubated in defined reaction buffers specific for various selected nucleases. Following treatment of the products with proteinase K, urea was added and analysis on 20% polyacrylamide gels containing urea was done. Gel products

- 64 -

were visualized by staining using Stains All (Sigma Chemical Co.). Laser densitometry was used to quantitate the extend of degradation. The effects of the modifications were determined for specific nucleases and compared with the results obtained from the serum and cytoplasmic systems.

### PROCEDURE 3 - 5-Lipoxygenase Anaylsis, Therapeutics and Assays

#### A. Therapeutics

For therapeutic use, an animal suspected of having a disease characterized by excessive or abnormal supply of 5-lipoxygenase is treated by administering oligonucleotide in accordance with this invention. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Such treatment is generally continued until either a cure is effected or a diminution in the diseased state is achieved. Long term treatment is likely for some diseases.

#### B. Research Reagents

The oligonucleotide compounds of this invention are also useful as research reagents when used to cleave or otherwise modulate 5-lipoxygenase mRNA in crude cell lysates or in partially purified or wholly purified RNA preparations. This application of the invention is accomplished, for example, by lysing cells by standard methods, optimally extracting the RNA and then treating it with a composition as described in the following Examples at concentrations ranging as for instances from about 100 to about 500 ng per 10 Kg of total RNA in a buffer consisting, for example, of 50 mM phosphate, pH ranging from about 4-10 at a temperature from about 30 to about 50C. The cleaved 5-lipoxygenase RNA can be analyzed by agarose gel electrophor sis and hybridization with radiolabeled DNA probes or by other standard methods.

- 65 -

### C. Diagnostics

The compounds of this invention will also be useful in diagnostic applications, particularly for the determination of the expression of specific mRNA species in various tissues or the expression of abnormal or mutant RNA species. In this example, the oligonucleotide compound targets a hypothetical abnormal mRNA by being designed complementary to the abnormal sequence, but would not hybridize to or cleave the normal mRNA.

10 Tissue samples can be homogenized, and RNA optionally extracted by standard methods. The crude homogenate or extract can be treated for example to effect cleavage of the target RNA. The product can then be hybridized to a solid support which contains a bound oligonucleotide

15 complementary to a region on the 5' side of the cleavage site. Both the normal and abnormal 5' region of the mRNA would bind to the solid support. The 3' region of the abnormal RNA, which was cleaved by the invention compound, would not be bound to the support and therefore

20 would be separated from the normal mRNA.

Several preferred embodiments of this invention are exemplified in accordance with the following examples. The target mRNA species for modulation relates to 5-lipoxygenase. Persons of ordinary skill in the art will

25 appreciate that the present invention is not so limited, however, and that it is generally applicable. The inhibition or modulation of production of the enzyme 5-lipoxygenase is expected to have significant therapeutic benefits in the treatment of disease. In order to assess

30 the effectiveness of the compositions, an assay or series of assays is required.

### D. In Vitro Assays

The cellular assays for 5-lipoxygenase preferably use the human promyelocytic leukemia cell line HL-60. These

35 cells can be induced to differentiate into either a monocytic-like cell or a neutrophil-like cell by various known agents. Treatment of the cells with 1.3% dimethyl sulfoxide, DMSO, is known to promote differentiation of the cells into neutrophils. It has now been found that

- 66 -

basal HL-60 cells do not synthesize detectable levels of 5-lipoxygenase protein or secrete leukotrienes (a downstream product of 5-lipoxygenase). Differentiation of the cells with DMSO causes an appearance of 5-lipoxygenase protein and leukotriene biosynthesis 48 hours after addition of DMSO. Thus induction of 5-lipoxygenase protein synthesis can be utilized as a test system for analysis of antisense oligonucleotides which interfere with 5-lipoxygenase synthesis in these cells.

10 A second test system for antisense oligonucleotides makes use of the fact that 5-lipoxygenase is a "suicide" enzyme in that it inactivates itself upon reacting with substrate. Treatment of differentiated HL-60 or other cells expressing 5 lipoxygenase, with 10  $\mu$ M A23187, a

15 calcium ionophore, promotes translocation of 5-lipoxygenase from the cytosol to the membrane with subsequent activation of the enzyme. Following activation and several rounds of catalysis, the enzyme becomes catalytically inactive. Thus, treatment of the

20 cells with calcium ionophore inactivates endogenous 5-lipoxygenase. It takes the cells approximately 24 hours to recover from A23187 treatment as measured by their ability to synthesize leukotriene B<sub>4</sub>. Antisense oligonucleotides directed against 5-lipoxygenase can be

25 tested for activity in two HL-60 model systems using the following quantitative assays. The assays are described from the most direct measurement of inhibition of 5-lipoxygenase protein synthesis in intact cells to more downstream events such as measurement of 5-lipoxygenase

30 activity in intact cells.

The most direct effect which antisense oligonucleotides exert on intact cells which can be easily be quantitated is specific inhibition of 5-lipoxygenase protein synthesis. To perform this

35 technique, cells can be labelled with <sup>35</sup>S-methionine (50  $\mu$ Ci/mL) for 2 hours at 37° C to label newly-synthesized protein. Cells are extracted to solubilize total cellular proteins and 5-lipoxygenase is immunoprecipitated with 5-lipoxygenase antibody followed by lution

- 67 -

from protein A Sepharose beads. The immunoprecipitated proteins are resolved by SDS-polyacrylamide gel electrophoresis and exposed for autoradiography. The amount of immunoprecipitated 5-lipoxygenase is  
5 quantitated by scanning densitometry.

A predicted result from these experiments would be as follows. The amount of 5-lipoxygenase protein immunoprecipitated from control cells would be normalized to 100%. Treatment of the cells with 1  $\mu$ M, 10  $\mu$ M, and 30  $\mu$ M  
10 of effective antisense oligonucleotide for 48 hours would reduce immunoprecipitated 5-lipoxygenase to 5%, 25% and 75% of control, respectively.

Measurement of 5-lipoxygenase enzyme activity in cellular homogenates could also be used to quantitate the  
15 amount of enzyme present which is capable of synthesizing leukotrienes. A radiometric assay has now been developed for quantitating 5-lipoxygenase enzyme activity in cell homogenates using reverse phase HPLC. Cells are broken by sonication in a buffer containing protease inhibitors  
20 and EDTA. The cell homogenate is centrifuged at 10,000 x g for 30 min and the supernatants analyzed for 5-lipoxygenase activity. Cytosolic proteins are incubated with 10  $\mu$ M  $^{14}$ C-arachidonic acid, 2mM ATP, 50  $\mu$ M free calcium, 100  $\mu$ g/ml phosphatidylcholine, and 50 mM bis-  
25 Tris buffer, pH 7.0, for 5 min at 37° C. The reactions are quenched by the addition of an equal volume of acetone and the fatty acids extracted with ethyl acetate. The substrate and reaction products are separated by reverse phase HPLC on a Novapak C18 column (Waters Inc.,  
30 Millford, MA). Radioactive peaks are detected by a Beckman model 171 radiochromatography detector. The amount of arachidonic acid converted into di-HETE's and mono-HETE's is used as a measure of 5-lipoxygenase activity.

35 A predicted result for treatment of DMSO differentiated HL-60 cells for 72 hours with an effective antisense oligonucleotide at 1  $\mu$ M, 10  $\mu$ M, and 30  $\mu$ M would be as follows. Control cells oxidize 200 pmol arachidonic acid/ 5 min/  $10^6$  cells. Cells treated with

- 68 -

1  $\mu$ M, 10  $\mu$ M, and 30  $\mu$ M of an effective oligonucleotide would oxidize 195 pmol, 140 pmol, and 60 pmol of arachidonic acid/ 5 min/  $10^6$  cells respectively.

A quantitative competitive enzyme linked immunosorbant assay (ELISA) for the measurement of total 5-lipoxygenase protein in cells has been developed. Human 5-lipoxygenase expressed in *E. coli* and purified by extraction, Q-Sepharose, hydroxyapatite, and reverse phase HPLC is used as a standard and as the primary antigen to coat microtiter plates. 25 ng of purified 5-lipoxygenase is bound to the microtiter plates overnight at 4° C. The wells are blocked for 90 min with 5% goat serum diluted in 20 mM Tris-HCL buffer, pH 7.4, in the presence of 150 mM NaCl (TBS). Cell extracts (0.2% Triton X-100, 12,000 x g for 30 min.) or purified 5-lipoxygenase were incubated with a 1:4000 dilution of 5-lipoxygenase polyclonal antibody in a total volume of 100  $\mu$ L in the microtiter wells for 90 min. The antibodies were prepared by immunizing rabbits with purified human recombinant 5-lipoxygenase. The wells were washed with TBS containing 0.05% tween 20 (TBST), then incubated with 100  $\mu$ L of a 1:1000 dilution of peroxidase conjugated goat anti-rabbit IgG (Cappel Laboratories, Malvern, PA) for 60 min at 25° C. The wells are washed with TBST and the amount of peroxidase labelled second antibody determined by development with tetramethylbenzidine.

Predicted results from such an assay using a 30 mer antisense oligonucleotide at 1  $\mu$ M, 10  $\mu$ M, and 30  $\mu$ M would be 30 ng, 18 ng and 5 ng of 5-lipoxygenase per  $10^6$  cells, respectively with untreated cells containing about 34 ng 5-lipoxygenase.

A net effect of inhibition of 5-lipoxygenase biosynthesis is a diminution in the quantities of leukotrienes released from stimulated cells. DMSO-differentiated HL-60 cells release leukotriene B4 upon stimulation with the calcium ionophore A23187. Leukotriene B4 released into the cell medium can be quantitated by radioimmunoassay using commercially available diagnostic kits (New England Nuclear, Boston,

- 69 -

MA). Leukotriene B<sub>4</sub> production can be detected in HL-60 cells 48 hours following addition of DMSO to differentiate the cells into a neutrophil-like cell. Cells ( $2 \times 10^5$  cells/mL) will be treated with increasing concentrations of antisense oligonucleotides for 48-72 hours in the presence of 1.3 % DMSO. The cells are washed and resuspended at a concentration of  $2 \times 10^6$  cell/mL in Dulbecco's phosphate buffered saline containing 1% delipidated bovine serum albumin. Cells are stimulated with  $10 \mu\text{M}$  calcium ionophore A23187 for 15 min and the quantity of LTB<sub>4</sub> produced from  $5 \times 10^5$  cell determined by radioimmunoassay as described by the manufacturer.

Using this assay the following results would likely be obtained with a 15-mer antisense oligonucleotide (GCAAGGTCACTGAAG) directed to the 5-LO mRNA. Cells will be treated for 72 hours with either  $1 \mu\text{M}$ ,  $10 \mu\text{M}$  or  $30 \mu\text{M}$  oligonucleotide in the presence of 1.3% DMSO. The quantity of LTB<sub>4</sub> produced from  $5 \times 10^5$  cells would be expected to be about 75 pg, 50 pg, and 35 pg, respectively with untreated differentiated cells producing 75 pg LTB<sub>4</sub>.

#### E. In Vivo Assay

Inhibition of the production of 5-lipoxygenase in the rat can be demonstrated in accordance with the following protocol. Topical application of arachidonic acid results in the rapid production of leukotriene B<sub>4</sub>, leukotriene C<sub>4</sub> and prostaglandin E<sub>2</sub> in the skin followed by edema and cellular infiltration. Certain inhibitors of 5-lipoxygenase have been known to exhibit activity in this assay. For the assay, 2 mg of arachidonic acid is applied to a rat ear with the contralateral ear serving as a control. The polymorphonuclear cell infiltrate is assayed by myeloperoxidase activity in homogenates taken from a biopsy 1 hour following the administration of arachidonic acid. The edematous response is quantitated by measurement of ear thickness and wet weight of a punch biopsy. Measurement of leukotriene B<sub>4</sub> produced in biopsy specimens is performed as a direct measurement of 5-



- 70 -

lipoygenase activity in the tissue. Antisense oligonucleotides will be applied topically to both ears 12 to 24 hours prior to administration of arachidonic acid to allow optimal activity of the compounds.

- 5 An expected result using a 15-mer antisense oligonucleotide directed against rat 5-lipoygenase corresponding to nucleotides 80-97 is given below. Both ears are pretreated for 24 hours with either 0.1  $\mu\text{mol}$ , 0.3  $\mu\text{mol}$ , or 1.0  $\mu\text{mol}$  antisense oligonucleotide prior to
- 10 challenge with arachidonic acid. Values are expressed as the mean for three animals per concentration. Inhibition of polymorphonuclear cell infiltration for 0.1  $\mu\text{mol}$ , 0.3  $\mu\text{mol}$ , and 1  $\mu\text{mol}$  is expected to be about 10 %, 75 % and 92 % of control activity, respectively. Inhibition of
- 15 edema is expected to be about 3 %, 58% and 90%, respectively while inhibition of leukotriene B<sub>4</sub> production would be expected to be about 15 %, 79% and 99%, respectively.

- 71 -

## EXAMPLES

## EXAMPLE 1

## 2'-O-(nonyl) adenosine (1)

To a solution of 10g of adenosine in 400ml of dimethyl  
5 formamide is added 2.25g of 60% sodium hydride (oil).  
After one hour, 8.5 ml of 1-bromo-nonane is added. The  
reaction is stirred for 16 hours. Ice is added and the  
solution evaporated in vacuo. Water and ethyl acetate are  
added. The organic phase is separated, dried, and  
10 evaporated in vacuo to give a white solid, which is  
recrystallized from ethanol to yield 4.8g of the title  
compound, M.P. 143-144°C analysis for:  $C_{19}H_{31}N_5O_4$ .  
Calculated: C, 57.99; H, 7.94; N, 17.79. Found: C, 58.13;  
H, 7.93; N, 17.83.

## 15 EXAMPLE 2

2'-O-(nonyl)-N<sup>6</sup>-benzoyl adenosine (2)

2'-O-(nonyl)adenosine (1) is treated with benzoyl  
chloride in a manner similar to the procedure of B.L.  
Gaffney and R.A. Jones, Tetrahedron Lett., Vol. 23, p.  
20 2257 (1982). After chromatography on silica gel (ethyl  
acetate-methanol), the title compound was obtained.  
Analysis for:  $C_{26}H_{35}N_5O_5$ . Calculated: C, 62.75; H, 7.09; N,  
17.07. Found: C, 62.73; H, 14.07; N, 13.87.

## EXAMPLE 3

25 2'-O-(nonyl)-5'-O-dimethoxytrityl-N<sup>6</sup>-benzoyl adenosine (3)

To a solution of 4.0g of 2'-O-(nonyl) N<sup>6</sup>-  
benzoyl adenosine (2) in 250ml of pyridine is added 3.3g  
of 4,4'-dimethoxy-trityl chloride. The reaction is  
stirred for 16 hours. The reaction is added to  
30 ice/water/ethyl acetate, the organic layer is separated,  
dried, and concentrated in vacuo to a gum. 5.8g of the  
title compound was obtained after chromatography on  
silica gel (ethyl acetate-methanol triethylamine).  
Analysis for:  $C_{47}H_{53}N_5O_7$ . Calculated: C, 70.56; H, 6.68; N,  
35 8.75. Found: C, 70.26; H, 6.70; N, 8.71.

- 72 -

**EXAMPLE 4**

**N<sup>6</sup>-benzoyl-5'-0-dimethoxytrityl-2'-0-(nonyl)adenosine-3'-0,N,N-diisopropyl- $\beta$ -cyanoethyl phosphoramidite (4)**

2'-0-(nonyl)-5'-0-dimethoxytrityl-N-benzoyladenosine  
5 (3) is treated with ( $\beta$ -cyanoethoxy)chloro(N,N-diisopropyl)aminephosphane in a manner similar to the procedure of F. Seela and A. Kehne, Biochemistry, Vol. 26, p. 2233 (1987). After chromatography on silica gel (E=OAC/hexane), the title compound was obtained as a  
10 white foam.

**EXAMPLE 5**

**2'-0-(pentyl) adenosine (5)**

The title compound is prepared as per example 1, using 1-bromopentane instead of 1-bromononane, M.P.-108-109°C.  
15 Analysis for: C<sub>15</sub>H<sub>24</sub>N<sub>5</sub>O<sub>4</sub>. Calculated: C, 53.24; H, 7.15; N, 20.69. Found: C, 53.37; H, 6.86; N, 20.51.

**EXAMPLE 6**

**2'-0-(pentyl)-N<sup>6</sup>-benzoyladenosine (6)**

Benzoylation of 2'-0-(pentyl) adenosine, as per  
20 example 2, gives the title compound. Analysis for: C<sub>22</sub>H<sub>29</sub>N<sub>5</sub>O<sub>5</sub>. Calculated: C, 59.58; H, 6.59; N, 15.79. Found: C, 59.34; H, 6.27; N, 15.53.

**EXAMPLE 7**

**2'-0-(pentyl)-5'-0-dimethoxytrityl-N<sup>6</sup>-benzoyladenosine (7)**

25 The title compound is prepared from 2'-0-(pentyl)-N<sup>6</sup>-benzoyladenosine (6) as per example 3. Analysis for: C<sub>43</sub>H<sub>46</sub>N<sub>5</sub>O<sub>7</sub>. Calculated: C, 69.34; H, 6.22; N, 9.40. Found: C, 68.97; H, 6.07; N, 9.12.

- 73 -

## EXAMPLE 8

N<sup>6</sup>-benzoyl-5'-O-dimethoxytrityl-2'-O-(pentyl)adenosine-3'-O, N,N-diisopropyl-β-cyanoethyl phosphoramidite (8)

The title compound is prepared from 2'-O-(pentyl)-5'-O-dimethoxytrityl-N<sup>6</sup>-benzoyl-adenosine (7) as per example 4.

## EXAMPLE 9

2'-O-(benzyl) adenosine (9)

The title compound is prepared as per example 1, using benzylbromide instead of 1-bromononane. After chromatography on silica gel (E=OAC-meoH) gives a white solid. M.P.-87-88°C. Analysis for: C<sub>17</sub>H<sub>19</sub>N<sub>5</sub>O<sub>6</sub>·25m H<sub>2</sub>O. Calculated: C, 56.42; H, 5.43; N, 19.35. Found: C, 56.46; H, 5.23; N, 19.50.

## 15 EXAMPLE 10

2'-O-(benzyl)-N<sup>6</sup>-benzoyl-adenosine (10)

Benzoylation of 2'-O-(benzyl)adenosine (9), as per example 2, gives the title compound. Analysis for: C<sub>24</sub>H<sub>23</sub>N<sub>5</sub>O<sub>5</sub> · 25m H<sub>2</sub>O. Calculated: C, 61.86; H, 5.08; N, 15.03. Found C, 61.89; H, 5.01; N, 14.88.

## EXAMPLE 11

2'-O-(benzyl)-5'-O-dimethoxytrityl-N<sup>6</sup>-benzoyl-adenosine (11)

The title compound is prepared from 2'-O-(benzyl)-N<sup>6</sup>-benzoyl-adenosine (10) as per example 3. Analysis for: C<sub>45</sub>H<sub>41</sub>N<sub>5</sub>O<sub>7</sub>. Calculated: C, 70.76; H, 5.41; N, 9.17. Found: C, 70.84; H, 5.35; N, 8.90.

- 74 -

**EXAMPLE 12**

**N<sup>6</sup>-benzoyl-5'-0-dimethoxytrityl-2'-0-(benzyl)adenosine-3'-O,N,N-diisopropyl-β-cyanoethyl phosphoramidite (12)**

The title compound is prepared from 2'-0-(benzyl)5'-0-dimethoxytrityl-N<sup>6</sup>-benzoyladenosine (11) as per example 4. The compound is a white foam.

<sup>31</sup>P-NMR(CDCN) S 151.08, 151.25

**EXAMPLE 13**

**2'-0-(butyl) adenosine (13)**

The title compound is prepared as per example 1 using 1-bromobutane instead of 1-bromononane. After chromatography on silica gel (E=OAC-MEOH), a white solid was obtained.

1 H-NMR (DMSO-d<sub>6</sub>): 5.97 (d, 1H, C, H).

**EXAMPLE 14**

**2'-0-(butyl)-N<sup>6</sup>-benzoyladenosine (14)**

Benzoylation of 2'-0-(butyl)adenosine (13), as per example 2, gives the title compound as a white foam.

1 H-NMR(DMSO-d<sub>6</sub>):6.11 (d,1H,C,H).

**EXAMPLE 15**

**2'-0-(butyl)-5'-0-dimethoxytrityl-N<sup>6</sup>-benzoyl-adenosine (15)**

The title compound is prepared from 2'-0-(butyl)-N<sup>6</sup>-benzoyladenosine (14) as per example 3. After chromatography of silica gel (E=OAC-Hexane-TRA) a yellow foam was obtained. Analysis for: C<sub>42</sub>H<sub>43</sub>N<sub>5</sub>O<sub>7</sub>. Calculated: C, 69.12; H, 5.94; N, 9.59. Found: C, 69.21; H, 5.99; N, 9.43.

**EXAMPLE 16**

**N<sup>6</sup>-benzoyl-5'-0-dimethoxytrityl-2'-0-(butyl)adenosine-3'-O,N,N-diisopropyl-β-cyanoethyl phosphoramidite (16)**

The title compound is prepared from 2'-0-(butyl)-5'-0-dimethoxytrityl-N<sup>6</sup>-benzoyladenosine (15) as per example 4. The compound is a white foam.

**EXAMPLE 17****2'-O-(propyl)adenosine (17)**

The title compound is prepared as per example 1 using 1-bromopropane instead of 1-bromononane, M.P. 127-128°C.  
5 Analysis for:  $C_{13}H_{19}N_5O_4$ . Calculated: C, 50.48; H, 6.19; N, 22.64. Found: C, 50.58; H, 6.21; N, 22.56.

**EXAMPLE 18****2'-O-(propyl)-N<sup>6</sup>-benzoyladenosine (18)**

Benzoylation of 2'-O-(propyl)adenosine (17), as per  
10 example 2, gave the title compound. Analysis for:  $C_{20}H_{23}N_5O_5$ . Calculated: C, 58.10; H, 5.61; N, 16.94. Found C, 58.12; H, 5.58; N, 16.79.

**EXAMPLE 19****2'-O-(propyl)-5'-O-dimethoxytrityl-N<sup>6</sup>-benzoyladenosine (19)**  
15

The title compound is prepared from 2'-O-(propyl)-N<sup>6</sup>-benzoyladenosine (18) as per example 3.

1 H-NMR (DMSO-d<sub>6</sub>): 6.13(d, 1H, C, H)

**EXAMPLE 20****N<sup>6</sup>-benzoyl-5'-O-dimethoxytrityl-2'-O-(propyl)adenosine-3'-O,N,N-diisopropyl-β-cyanoethyl phosphoramidite (20)**  
20

The title compound is prepared from 2'-O-(propyl)-5'-O-dimethoxytrityl-N<sup>6</sup>-benzoyladenosine (19) as per example 4. The compound is a white foam.

**EXAMPLE 21**  
25**2'-O-(allyl) adenosine (21)**

The title compound is prepared as per example 1 using allylbromide instead of 1-bromononane. The compound is a white solid, after chromatography on silica gel (E=OAC-methanol).  
30

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): S 6.03 (d, 1H, C, H).

- 76 -

**EXAMPLE 22****2'-O-(allyl)-N<sup>6</sup>-benzoyladenosine (22)**

Benzoylation of 2'-O-(allyl)adenosine (21), as per example 2, gave the title compound as a white foam.

5 <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): S 6.17 (d, 1H, C, H).

**EXAMPLE 23****2'-O-(allyl)-5'-O-dimethoxytrityl-N<sup>6</sup>-benzoyl-adenosine (23)**

The title compound is prepared from 2'-O-(allyl)-N<sup>6</sup>-benzoyladenosine (22) as per example 3. Analysis for:  
10 C<sub>41</sub>H<sub>39</sub>N<sub>5</sub>O<sub>7</sub>. Calculated: C, 68.99; H, 5.51; N, 9.81. Found: C, 68.68; H, 5.43; N, 9.70.

**EXAMPLE 24****N<sup>6</sup>-benzoyl-5'-O-dimethoxytrityl-2'-O-(allyl)adenosine-3'-O,N,N-diisopropyl-β-cyanoethyl phosphoramidite (24)**  
15

The title compound is prepared from 2'-O-(allyl)-5'-O-dimethoxytrityl-N<sup>6</sup>-benzoyladenosine (23) as per example 4. The compound is a white foam.

<sup>32</sup>P NMR (CD<sub>3</sub>CN) S 151.11, 151.32.

**20 EXAMPLE 25****2'-O-(propylphthalimide)adenosine (25)**

The title compound is prepared as per example 1, using N-(3-bromopropyl)phthalimide. Chromatography on silica gel gives a white solid, M.P. 123-124°C. Analysis for:  
25 C<sub>21</sub>H<sub>22</sub>N<sub>6</sub>O<sub>6</sub>. Calculated: C, 55.03; H, 4.88; N, 18.49. Found: C, 55.38; H, 4.85; N, 18.46.

**EXAMPLE 26****2'-O-(propylphthalimide)-N<sup>6</sup>-benzoyladenosine (26)**

Benzoylation of 2'-O-(propylphthalimide)adenosine (25), as per example 2, gives the title compound.  
30 Analysis for: C<sub>28</sub>H<sub>28</sub>N<sub>6</sub>O<sub>7</sub>. Calculated: C, 60.21; H, 4.69; N, 15.05. Found: C, 59.94; H, 4.66; N, 14.76.

- 77 -

**EXAMPLE 27**

2'-O-(propylphthalimide)-5'-O-dimethoxytrityl-N-benzoyladenosine (27)

The title compound is prepared from 2'-O-(propylphthalimide)-N<sup>6</sup>-benzoyladenosine (26) as per example 3. Analysis for: C<sub>49</sub>H<sub>44</sub>N<sub>6</sub>O<sub>9</sub>. Calculated: C, 68.36; H, 5.15; N, 9.76. Found: C, 68.16; H, 5.03; N, 9.43.

**EXAMPLE 28**

10 N<sub>6</sub>-benzoyl-5'-O-dimethoxytrityl-2'-(propyl)adenosine-3'-O,N,N-diisopropyl-β-cyanoethylphosphoramidite (28)

The title compound is prepared from 2'-O-(propylphthalimide)-5'-O-dimethoxytrityl-N<sup>6</sup>-benzoyladenosine (27) as per example 4. A white foam was obtained.

**EXAMPLE 29**

15 2'-O-(butylphthalimide)adenosine (29)

The title compound is prepared as per example 1, using n-(4-bromobutyl)phthalimide instead of 1-bromononane. Chromatography on silica gel (E=OAC-MeOH) gives a white solid. M.P. 199-200°C. Analysis for: C<sub>22</sub>H<sub>24</sub>N<sub>6</sub>O<sub>6</sub>. Calculated: C, 56.42; H, 5.16; N, 17.94. Found: C, 56.31; H, 5.04; N, 17.95.

**EXAMPLE 30**

2'-O-(butylphthalimide)-N<sup>6</sup>-benzoyladenosine (30)

25 Benzoylation of 2'-O-(butylphthalimide)adenosine (29) as per example 2, gives the title compound. Analysis for: C<sub>29</sub>H<sub>24</sub>N<sub>6</sub>O<sub>7</sub>. Calculated: C, 60.83; H, 4.93; N, 14.68. Found: C, 60.48; N, 14.41.

**EXAMPLE 31**

30 2'-O-(butylphthalimide)-5'-O-dimethoxytrityl-N-benzoyladenosine (31)

The title compound is prepared from 2'-O-(butylphthalimide)-N<sup>6</sup>-benzoyladenosine (30) as per example 3. Analysis for: C<sub>50</sub>H<sub>46</sub>N<sub>6</sub>O<sub>9</sub>. Calculated: C, 68.64; H, 5.29; N, 9.60. Found: C, 68.47; H, 5.12; N, 9.37.



## EXAMPLE 32

N<sup>6</sup>-benzoyl-5'-O-dimethoxytrityl-2'-O-(butylphthalimide)adenosine-3'-O,N,N-diisopropyl-β-cyanoethylphosphoramidine (32)

- 5 The title compound was prepared from compound (31) as per example 4.

<sup>31</sup>P NMR (CD<sub>3</sub>CN) δ 150.88, 151.22.

## EXAMPLE 33

2'-O-(pentylphthalimide)adenosine (33)

- 10 The title compound is prepared as per example 1, using N-(5-bromopentyl)phthalimide instead of 1-bromononane. M.P. 159-160°C. Analysis for: C, 57.26; H, 5.43; N, 17.42. Found: C, 57.03; H, 5.46; N, 17.33.

## EXAMPLE 34

- 15 2'-O-(pentylphthalimide)-N<sup>6</sup>-benzoyladenosine (34)

Benzoylation of 2'-O-(pentylphthalimide)adenosine (33) as per example 2 gives the title compound. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 6.10 (d, 1H, C<sub>1</sub>, H). Analysis for: C<sub>30</sub>H<sub>30</sub>N<sub>6</sub>O<sub>7</sub>. Calculated: C, 61.43; H, 5.16; N, 14.33. Found: C, 61.51;

- 20 H, 4.97; N, 14.10.

## EXAMPLE 35

2'-O-(pentylphthalimide)-5'-dimethoxytrityl-N<sup>6</sup>-benzoyladenosine (35)

- 25 The title compound is prepared from 2'-O-(pentylphthalimide)-N<sup>6</sup>-benzoyladenosine (34) as per example 3. After chromatography on silica gel (ethylacetate, hexane, triethylamine), the title compound was obtained. Analysis for: C<sub>51</sub>H<sub>48</sub>N<sub>6</sub>O<sub>9</sub>. Calculated: C, 68.91; H, 5.44; N, 9.45. Found: C, 68.65; H, 5.45; N, 9.61.

**EXAMPLE 36**

N<sup>6</sup>-benzoyl-5'-O-dimethoxytrityl-2'-O-(pentylphthalimide) adenosine-3'-O,N,N-diisopropyl-β-cyanoethylphosphoramidite (36)

- 5 The title compound is prepared from 2'-O-(pentylphthalimide)-5'-O-dimethoxytrityl-N<sup>6</sup>-benzoyladenosine (35) as per example 4. The compound is a white foam.

**EXAMPLE 37**

2'-O-(ethyl)adenosine (37)

- 10 The title compound is prepared as per example 1 using ethyl bromide instead of 1-bromononane. The compound is a white solid. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): S 6.00 (d, 1H, C<sub>1</sub>, H)

**EXAMPLE 38**

2'-O-(ethyl)-N-benzoyladenosine (38)

- 15 Benzoylation of 2'-O-(ethyl)adenosine (37), as per example 2, gives the title compound. The compound is a white foam.

**EXAMPLE 39**

2'-O-(ethyl)-5'-O-dimethoxytrityl-N<sup>6</sup>-benzoyladenosine (39)

- 20 The title compound is prepared from 2'-O-(ethyl)-N<sup>6</sup>-benzoyladenosine (38) as per example 3. The compound is a white foam.

**EXAMPLE 40**

N<sup>6</sup>-benzoyl-5'-O-dimethoxytrityl-2'-O-(ethyl)adenosine-3'-O,N,N-diisopropyl-β-cyanoethylphosphoramidite (40)

- 25 The title compound is prepared from 2'-O-(ethyl)-5'-O-dimethoxytrityl-N<sup>6</sup>-benzoyladenosine (39) as per example 4. The compound is a white foam.

**EXAMPLE 41**

- 30 2'-O-[imidazo-1-yl-(propyl)]adenosine (41)

The title compound can be prepared as per example 1 using 1-(3-bromopropyl)imidazole in place of 1-bromononane.

- 80 -

**EXAMPLE 42****2'-O-[imidazo-1-yl-(propyl)]-N<sup>6</sup>-benzoyladenosine (42)**

Benoylation of 2'-O-[imidazo-1-yl-(propyl)]adenosine (41) as per example 2 will give the title compound.

**5 EXAMPLE 43****2'-O-[imidazo-1-yl-(propyl)]-5'-O-dimethoxytrityl-N<sup>6</sup>-benzoyladenosine (43)**

The title compound can be prepared from 2'-O-[imidazo-1-yl-(propyl)]adenosine (42) as per example 3.

**10 EXAMPLE 44****N<sup>6</sup>-benzoyl-5'-O-dimethoxytrityl-2'-O-[imidazo-1-yl-(propyl)]-adenosine-3'-O,N,N-diisopropyl-β-cyanoethyl phosphoramidite (44)**

The title compound can be prepared from 2'-O-[imidazo-1-yl-(propyl)]-5'-O-dimethoxytrityl-N<sup>6</sup>-benzoyladenosine (43) as per example 4.

**EXAMPLE 45****2'-O-(phenylpropyl) adenosine (45)**

The title compound can be prepared as per example 1 using 3-bromopropylbenzene in place of 1-bromononane.

**EXAMPLE 46****2'-O-(phenylpropyl)-N<sup>6</sup>-benzoyladenosine (46)**

Benoylation of 2'-O-(phenylpropyl)adenosine (45) as per example 2 will give the title compound.

**25 EXAMPLE 47****2'-O-(phenylpropyl)-5'-O-dimethoxytrityl-N<sup>6</sup>-benzoyl-adenosine (47)**

The title compound can be prepared from 2'-O-(phenylpropyl) adenosine (46) as per example 3.

**EXAMPLE 48**

**N<sup>6</sup>-benzoyl-5'-O-dimethoxytrityl-2'-O-(phenylpropyl)-adenosine-3'-O,N,N-diisopropyl-β-cyanoethylphosphoramidite (48)**

- 5     The title compound can be prepared from 2'-O-(phenylpropyl)-5'-O-dimethoxytrityl-N<sup>6</sup>-benzoyl adenosine (47) as per example 4.

**EXAMPLE 49**

**2'-O-[naphth-1-yl-(propyl)] adenosine (49)**

- 10     The title compound can be prepared as per example 1 using 1-(3-bromopropyl)naphthalene in place of 1-bromononane.

**EXAMPLE 50**

**2'-O-[naphth-1-yl-(propyl)]-N<sup>6</sup>-benzoyl-adenosine (50)**

- 15     Benoylation of 2'-O-[naphth-1-yl-(propyl)]adenosine (49) as per example 2 will give the title compound.

**EXAMPLE 51**

**2'-O-[naphth-1-yl-(propyl)]-5'-O-dimethoxytrityl-N<sup>6</sup>-benzoyl-adenosine (51)**

- 20     The title compound can be prepared from 2'-O-[naphth-1-yl-(propyl)]-N<sup>6</sup>-benzoyl adenosine (50) as per example 3.

**EXAMPLE 52**

**N<sup>6</sup>-benzoyl-5'-O-dimethoxytrityl-2'-O-[naphth-1-yl-(propyl)] adenosine-3'-O,N,N-diisopropyl-β-cyanoethyl phosphoramidite (52)**

- 25     The title compound can be prepared from 2'-O-[naphth-1-yl-(propyl)]-5'-O-dimethoxytrityl-N<sup>6</sup>-benzoyl adenosine (51) as per example 4.

**EXAMPLE 53**

- 30     **2'-O-[anthrac n-2-yl-(propyl)]adenosine (53)**

Th title compound can be prepared as per xample 1 using 2-(3-bromopropyl)anthracene in place of 1-bromononane.

**EXAMPLE 54**

2'-O-[anthracen-2-yl-(propyl)]-N<sup>6</sup>-benzoyl-adenosine (54)  
Benoylation of 2'-O-[anthracen-2-yl-(propyl)]adenosine  
(53) as per example 2 will give the title compound.

**5 EXAMPLE 55**

2'-O-[anthracen-2-yl-(propyl)]-5'-O-dimethoxytrityl-N<sup>6</sup>-  
benzoyl- adenosine (55)

The title compound can be prepared from 2'-O-[anthra-  
cen-2-yl-(propyl)]-N<sup>6</sup>-benzoyl adenosine (54) as per example  
10 3.

**EXAMPLE 56**

N<sup>6</sup>-benzoyl-5'-O-dimethoxytrityl-2'-O-[anthracen-2-yl-  
(propyl)] adenosine-3'-O,N,N-diisopropyl-β-cyanoethyl  
phosphoramidite (56)

15 The title compound can be prepared from 2'-O-[anthra-  
cen-2-yl-(propyl)]-5'-O-dimethoxytrityl-N<sup>6</sup>-benzoyl-  
adenosine (55) as per example 4.

**EXAMPLE 57**

2'-O-[bis(phthalimidobutyl)-(aminobutyl)]-N<sup>6</sup>-benzoyl-  
20 adenosine (57)

Compound 30 is treated with TIPS reagent (tetra-  
isopropyl-disilyldichloride) to block the 3',5'-hydroxyl  
groups with the TIPS blocking group. Treatment with  
ammonium hydroxide cleaves the phthalimide group yielding  
25 a 2'-O-amino butyl blocked adenosine. This compound can  
be further treated as per example 29 with n-(4-bromo-  
butyl)phthalimide to yield the 2'-O-polyalkylamino  
blocked adenosine. Deblocking of the TIPS blocking group  
with tetrabutyl ammonium fluoride will yield the title  
30 compound. The title compound can be used as per  
examples 31 and 32 to form suitable blocked nucleotides  
suitable for incorporation into oligonucleotides.

- 83 -

## EXAMPLE 58

2'-O-[bis(aminobutyl)-(aminobutyl)]-N<sup>6</sup>-benzoyladenine  
(58)

Treatment of compound 57 with ammonium hydroxide  
5 cleaves the phthalimide group to yield the title  
compound. This compound can be further treated in the  
manner of example 57 to further elongate the polyalkyl-  
amino chain.

## EXAMPLE 59

10 N<sup>6</sup>-Benzoyl-[2'-deoxy-2'-fluoro-5'-O-(4,4'-di-methoxy  
trityl)] adenosine-3'-O,N,N-diisopropyl-β-cyanoethyl  
phosphoramidite (59)

N<sup>6</sup>-Benzoyl-9-(2'-fluoro-β-D-ribofuranosyl)adenine was  
prepared from 9-β-D-arabinofuranosyladenine in a  
15 five-step synthesis using a modification of a procedure  
reported by M. Ikehara et al., Nucleosides and  
Nucleotides Vol. 2, pp. 373-385 (1983). Thus, the  
N<sup>6</sup>-benzoyl derivative was obtained in good yield utilizing  
the method of transient protection with chlorotri-  
20 methylsilane. R. A. Jones, J. Am. Chem. Soc. Vol. 104,  
pg. 1316 (1982). Selective protection of the 3' and  
5'-hydroxyl groups of N<sup>6</sup>-Benzoyl-9-β-D-arabinofuranosyl-  
adenine with tetrahydropyranyl (THP) was accomplished by  
modification of a literature procedure G. Butke, et al.,  
25 in Nucleic Acid Chemistry, Part 3, pp. 149-152, Townsend,  
L.B. and Tipson, R.S. eds., (J. Wiley and Sons, New York  
1986) to yield N<sup>6</sup>-Benzoyl-9-[3',5'-di-O-tetrahydropyran-  
2-yl)-β-D-arabino furanosyl] adenine in good yield.  
Treatment of N<sup>6</sup>-Benzoyl-9-[3',5'-di-O-tetrahydropyran-  
30 2-yl)-β-D-arabinofuranosyl] adenine with trifluoro-  
methanesulfonic anhydride in dichloromethane gave the  
2'-triflate derivative N<sup>6</sup>-Benzoyl-9-[2'-O-trifluoro-  
methylsulfonyl-3',5'-di-O-tetrahydropyran-2-yl)-β-D-  
arabinofuranosyl] adenine which was not isolated due to  
35 its lability. Displacement of the 2'-triflate group was  
effected by reaction with tetrabutyl ammonium fluoride in

- 84 -

tetrahydrofuran to obtain a moderate yield of the 2'-fluoro derivative N<sup>6</sup>-Benzoyl-9-[2'-fluoro-3',5'-di-O-tetra-hydropyran-2-yl)- $\beta$ -D-arabino furanosyl] adenine. Deprotection of the THP groups of N<sup>6</sup>-Benzoyl-9-[2'-fluoro-3',5'-di-O-tetrahydro- pyran-2-yl)- $\beta$ -D-arabinofuranosyl] adenine was accomplished by treatment with Dowex-50W in methanol to yield N<sup>6</sup>-benzoyl-9- (2'-deoxy-2'-fluoro- $\beta$ -D-ribofuranosyl) adenine in moderate yield. The <sup>1</sup>H-NMR spectrum was in agreement with the literature values. M. Ikehara and H. Miki, Chem. Pharm. Bull., Vol. 26, pp. 2449-2453 (1978). Standard methodologies such as those of foregoing examples 3 and 4 were employed to obtain the 5'-dimethoxytrityl-3'-phos-phoramidite intermediates N<sup>6</sup>-Benzoyl-9-[2'-fluoro-5'-O-(4,4'-di-methoxytrityl)- $\beta$ -D-ribofuranosyl] adenine and N<sup>6</sup>-Benzoyl-[2'- deoxy-2'-fluoro-5'-O-(4,4'-dimethoxy trityl)] adenosine-3'-O-(N,N- diisopropyl- $\beta$ -cyanoethyl phosphoramidite). K. K. Ogilvie, Can J. Chem. Vol. 67, pp. 831-839 (1989).

**EXAMPLE 60****20 N<sup>6</sup>-Benzoyl-9- $\beta$ -D-arabinofuranosyladenine (60)**

9- $\beta$ -D-arabinofuranosyladenine (1.07 g, 4.00 m.mol) was dissolved in anhydrous pyridine (20 mL) and anhydrous dimethyl formamide (20 mL) under an argon atmosphere. The solution was cooled to ice temperature and chloro-  
25 trimethyl-silane (3.88 mL, 30.6 m.mol) was added slowly to the reaction mixture via syringe. After stirring the reaction mixture at ice temperature for 30 minutes, benzoyl chloride (2.32 mL, 20 m.mol) was added slowly. The reac-tion mixture was allowed to warm to 20° C and  
30 stirred for 2 hours. After cooling the reaction mixture to ice temperature, cold water (8 mL) was added and the mixture was stirred for 15 minutes. Concentrated ammonium hydroxide (8 mL) was slowly added to the reac-tion mixture to give a final concentration of 2M of  
35 ammonia. After stirring the cold reaction mixture for 30 minutes, the solvent was evaporated in vacuo (60 torr) at 20° C followed by evaporati n in vacuo (1 torr) at 40°

- 85 -

C to give an oil. This oil was triturated with diethyl ether (50 ml) to give a solid which was filtered and washed with diethyl ether three times. This crude solid was tri-turated in methanol (100 ml) at reflux temperature three times and the solvent was evaporated to yield N<sup>6</sup>-Benzoyl-9-β-D-arabino-furanosyl adenine (60) as a solid (1.50 g, 100%).

**EXAMPLE 61**

N<sup>6</sup>-Benzoyl-9-[3',5'-di-O-tetrahydropyran-2-yl)-

10 β-D-arabino-furanosyl] adenine (61)

N<sup>6</sup>-Benzoyl-9-β-D-arabinofuranosyladenine (60) (2.62 g, 7.06 m.mol) was dissolved in anhydrous dimethylformamide (150 ml) under an argon atmosphere and p-toluenesulfonic acid monohydrate (1.32 g, 6.92 m.mol) was added. This solution was cooled to ice temperature and dihydropyran (1.26 ml, 13.8 m.mol) was added via syringe. The reaction mixture was allowed to warm to 20° C. Over a period of 5 hours a total of 10 equivalents of dihydropyran were added in 2 equivalent amounts in the fashion described.

20 The reaction mixture was cooled to ice temperature and saturated aqueous sodium bicarbonate was added slowly to a pH of 8, then water was added to a volume of 750 ml. The aqueous mixture was extracted with methylene chloride four times (4 x 200 ml), and the organic phases were

25 combined and dried over magnesium sulfate. The solids were filtered and the solvent was evaporated in vacuo (60 torr) at 30° C to give a small volume of liquid which was evaporated in vacuo (1 torr) at 40° C to give an oil. This oil was coevaporated with p-xylene in vacuo at 40°

30 to give an oil which was dissolved in methylene chloride (100 ml). Hexane (200 ml) was added to the solution and the lower-boiling solvent was evaporated in vacuo at 30° C to leave a white solid suspended in hexane. This solid was filtered and washed with hexane three times (3 x 10

35 ml) then purified by column chromatography using silica and methylene chloride-methanol (93:7, v/v) as eluent. The first fraction yielded the title compound (61) as a



- 86 -

white foam (3.19 g, 83 %) and a second fraction gave a white foam (0.81 g) which was characterized as the 5'-mono-tetrahydropyranyl derivative of N<sup>6</sup>-Benzoyl-9-β-D-arabino-furanosyl adenine.

5 EXAMPLE 62

N<sup>6</sup>-Benzoyl-9-[2'-O-trifluoromethylsulfonyl-3',5'-di-O-tetrahydro-pyran-2-yl)-β-D-arabinofuranosyl]adenine (62)

N<sup>6</sup>-Benzoyl-9-[3',5'-di-O-tetrahydropyran-2-yl)-β-D-arabinofuranosyl] adenine (61) (2.65 g, 4.91 m.mol) was  
10 dissolved in anhydrous pyridine (20 ml) and the solvent was evaporated in vacuo (1 mm Hg) at 40° C. The resulting oil was dissolved in anhydrous methylene chloride (130 ml) under an argon atmosphere and anhydrous pyridine (3.34 ml, 41.3 m.mol) and N,N-dimethylamino- pyridine  
15 (1.95 g, 16.0 mmol) were added. The reaction mixture was cooled to ice temperature and trifluoromethanesulfonic anhydride (1.36 ml, 8.05 mmol) was added slowly via syringe. After stirring the reaction mixture at ice temperature for 1 h, it was poured into cold saturated  
20 aqueous sodium bicarbonate (140 ml). The mixture was shaken and the organic phase was separated and kept at ice temperature. The aqueous phase was extracted with methylene chloride two more times (2 x 140 ml). The organic extracts which were diligently kept cold were  
25 combined and dried over magnesium sulfate. The solvent was evaporated in vacuo (60 torr) at 20° C then evaporated in vacuo (1 torr) at 20° C to give N<sup>6</sup>-Benzoyl-9- [2'-O-trifluoromethylsulfonyl-3',5'-di-O-tetrahydro-pyran-2-yl)- β-D-arabinofuranosyl] adenine (62) as a crude oil  
30 which was not purified further.

## EXAMPLE 63

N<sup>6</sup>-Benzoyl-9-[2'-fluoro-3',5'-di-O-tetrahydro-  
pyran-2-yl)-β-D-arabinofuranosyl]adenine (63)

N<sup>6</sup>-Benzoyl-9-[2'-O-trifluoromethylsulfonyl-3',5'-di-O-  
5 tetrahydropyran-2-yl)-β-D-arabinofuranosyl]adenine (62),  
(4.9 mmol) as a crude oil was dissolved in anhydrous  
tetrahydrofuran (120 ml) and this solution was cooled to  
ice temperature under an argon atmosphere. Tetrabutyl-  
ammonium fluoride as the hydrate (12.8 g, 49.1 mmol) was  
10 dissolved in anhydrous tetrahydrofuran (50 ml) and half  
of this volume was slowly added via syringe to the cold  
reaction mixture. After stirring at ice temperature for  
1 hour, the remainder of the reagent was added slowly.  
The reaction mixture was stirred at ice temperature for  
15 an additional 1 hour, then the solvent was evaporated in  
vacuo (60 torr) at 20° C to give an oil. This oil was  
dissolved in methylene chloride (250 ml) and washed with  
brine three times. The organic phase was separated and  
dried over magnesium sulfate. The solids were filtered  
20 and the solvent was evaporated to give an oil. The crude  
product was purified by column chromatography using  
silica in a sintered-glass funnel (600 ml) and ethyl  
acetate was used as eluent. N<sup>6</sup>-Benzoyl-9-[2'-fluoro-3',5'-  
-di-O- tetrahydropyran- 2-yl)-β-D-arabino-furanosyl]  
25 adenine (63) was obtained as an oil (2.03 g, 76%).

## EXAMPLE 64

N<sup>6</sup>-Benzoyl-9-(2'-fluoro-β-D-ribofuranosyl) adenine (64)

N<sup>6</sup>-Benzoyl-9-[2'-fluoro-3',5'-di-O-tetrahydropyran-  
2-yl) -β-D-arabinofuranosyl]adenine (63) (1.31 g, 2.42  
30 mmol) was dis- solved in methanol (60 ml), and Dowex 50W  
x 2-100 (4 cm<sup>3</sup>, 2.4 m.eq) was added to the reaction  
mixture. The reaction mixture was stir-red at 20° C for  
1 hour then cooled to ice temperature. Triethyl-amine (5  
ml) was then slowly added to th cold reaction mixture to  
35 a pH of 12. The resin was filtered and washed with 30%  
triethyl-amine in methanol until the wash no l ngr r  
contained UV abs rbing mat rial. Toluene (50 ml) was

- 88 -

added to the washes and the solvent was evaporated at 24° C in vacuo (60 torr then 1 torr) to give a residue. This residue was partially dissolved in methylene chloride (30 ml) and the solvent was transferred to a separatory funnel. The remainder of the residue was dissolved in hot (60° C) water and after cooling the solvent it was also added to the separatory funnel. The biphasic system was extracted, and the organic phase was separated and extracted three times with water (3 x 100 ml). The combined aqueous extracts were evaporated in vacuo (60 torr then 1 torr Hg) at 40° C to give an oil which was evaporated with anhydrous pyridine (50 ml). This oil was further dried in vacuo (1 torr Hg) at 20° C in the presence of phosphorous pentoxide overnight to give N<sup>6</sup>-Benzoyl 9-(2'-fluoro-β-D-ribo-furanosyl) adenine (64) as a yellow foam (1.08 g, 100 %) which contained minor impurities.

**EXAMPLE 65**

N<sup>6</sup>-Benzoyl-9-[2'-fluoro-5'-O-(4,4'-dimethoxy-trityl)-β-D-ribofuranosyl]adenine (65)

N<sup>6</sup>-Benzoyl-9-(2'-fluoro-β-D-ribofuranosyl)adenine (64) (1.08 g, 2.89 mmol) which contained minor impurities was dissolved in anhydrous pyridine (20 ml) under an argon atmosphere, and dry triethylamine (0.52 ml, 3.76 mmol) was added followed by addition of 4,4'-dimethoxytrityl chloride (1.13 g, 3.32 mmol). After 4 hours of stirring at 20° C the reaction mixture was transferred to a separatory funnel and diethyl ether (40 ml) was added to give a white suspension. This mixture was washed with water three times (3 x 10 ml), the organic phase was separated and dried over magnesium sulfate. Triethylamine (1 ml) was added to the solution and the solvent was evaporated in vacuo (60 torr Hg) at 20° C to give an oil which was evaporated with toluene (20 ml) containing triethylamine (1 ml). This crude product was purified by column chromatography using silica and ethyl acetat - triethylamine (99:1, v/v) followed by ethyl acetat -

- 89 -

methanol-triethylamine (80:19:1) to give the product in two fractions. The fractions were evaporated in vacuo (60 torr then 1 torr Hg) at 20° C to give a foam which was further dried in vacuo (1 torr Hg) at 20° C in the presence of sodium hydroxide to give N<sup>6</sup>-Benzoyl-9-[2'-fluoro-5'-O-(4,4'-dimethoxytrityl)-β-D-ribofuranosyl] adenine (65) as a foam (1.02 g, 52 %).

**EXAMPLE 66**

N<sup>6</sup>-Benzoyl-[2'-deoxy-2'-fluoro-5'-O-(4,4'-dimethoxytrityl)] adenosine-3'-O,N,N-diisopropyl-β-cyanoethyl phosphoramidite (66)

N<sup>6</sup>-Benzoyl-9-[2'-fluoro-5'-O-(4,4'-dimethoxytrityl)-β-D-ribofuranosyl] adenine (65) (1.26 g, 1.89 mmol) was dissolved in anhydrous dichloromethane (13 ml) under an argon atmosphere, diisopropylethylamine (0.82 ml, 4.66 mmol) was added, and the reaction mixture was cooled to ice temperature. Chloro(diisopropylamino)-β-cyanoethoxyphosphine (0.88 ml, 4.03 mmol) was added to the reaction mixture which was allowed to warm to 20° C and stirred for 3 hours. Ethylacetate (80 ml) and triethylamine (1 ml) were added and this solution was washed with brine solution three times (3 x 25 ml). The organic phase was separated and dried over magnesium sulfate. After filtration of the solids the solvent was evaporated in vacuo at 20° C to give an oil which was purified by column chromatography using silica and hexane-ethyl acetate-triethylamine (50:49:1) as eluent. Evaporation of the fractions in vacuo at 20° C gave a foam which was evaporated with anhydrous pyridine (20 ml) in vacuo (1 torr) at 26° C and further dried in vacuo (1 torr Hg) at 20° C in the presence of sodium hydroxide for 24 h to give N<sup>6</sup>-Benzoyl-[2'-deoxy-2'-fluoro-5'-O-(4,4'-dimethoxytrityl)] adenosine-3'-O,N,N-diisopropyl-β-cyanoethyl phosphoramidite (66) as a foam (1.05 g, 63 %).

- 90 -

## EXAMPLE 67

2'-Deoxy-2'-fluoro-5'-O-(4,4'-dimethoxytrityl)-uridine-3'-O-(N,N-diisopropyl- $\beta$ -cyanoethylphosphoramidite). (67)

The title compound is prepared via a modification of a literature procedure, J. Fox, et al., J. Org. Chem., Vol. 29, pp. 558-564 (1964). Thus, 2,2'-cyclouridine (5.65 g, 25 mmol) is treated with a solution of 70% hydrogen fluoride/pyridine (65 mL) in anhydrous dioxane (500 mL) at 120° C in a sealed steel vessel for twenty hours. The reaction mixture is cooled to ice temperature, poured into ice (400 mL), and the pH is adjusted to 7 by addition of solid CaCO<sub>3</sub>. After the solvent is evaporated in vacuo (1 torr), the residue is dissolved in methanol, adsorbed onto silica gel, and purified by column chromatography using silica and ethyl acetate-methanol (v/v, 20:1) as eluent to give 2'-deoxy-2'-fluoro-uridine as a white solid (4.81 g, 79%). The 5'-DMT and 3'-cyanoethoxy-diisopropyl phosphoramidite derivatized nucleoside are obtained in accordance with the procedures of examples 65 and 66.

## EXAMPLE 68

2'-Deoxy-2'-fluoro-5'-O-(4,4'-dimethoxytrityl)-cytidine-3'-O-(N,N-diisopropyl- $\beta$ -cyanoethylphosphoramidite). (68)

2'-deoxy-2'-fluorouridine (2.51 g, 10.3 mmol) is converted to corresponding cytidine analog via a literature procedure, C.B. Reese, et al., J. Chem. Soc. Perkin Trans I, pp. 1171-1176 (1982), by acetylation with acetic anhydride (3.1 mL, 32.7 mmol) in anhydrous pyridine (26 mL) at room temperature. The reaction is quenched with methanol, the solvent is evaporated in vacuo (1 torr) to give an oil which is coevaporated with ethanol and toluene. 3',5'-O-diacetyl-2'-deoxy-2'-fluorouridine is crystallized from ethanol to afford colorless crystals (2.38 g, 81 %). N-4-(1,2,4-triazol-1-yl)-3',5'-O-diacetyl-2'-deoxy-2'-fluoro-uridine is obtained in a 70% yield (2.37 g) by reaction of 3',5'-O-diacetyl-2'-deoxy-2'-fluoruridine (2.75 g, 9.61 mmol)

- 91 -

with 1,2,4-triazole (5.97 g, 86.5 mmol), phosphorus oxychloride (1.73 mL, 18.4 mmol), and triethylamine (11.5 mL, 82.7 mmol) in anhydrous acetonitrile at room temperature. After 90 min the reaction mixture is cooled to ice temperature and triethylamine (7.98 ml, 56.9 mmol) is added followed by addition of water (4.0 ml). The solvent is evaporated in vacuo (1 torr) to give an oil which is dissolved in methylene chloride and washed with saturated aqueous sodium bicarbonate. The aqueous phase is extracted with methylene chloride twice (2 x 100 mL) and the organic extracts are dried with magnesium sulfate. Evaporation of the solvent afforded an oil from which the product N-4-(1,2,4-triazol-1-yl)-3',5'-O-diacetyl-2'-deoxy-2'-fluorouridine is obtained by crystallization from ethanol. 2'-deoxy-2'-fluorocytidine is afforded by treatment of protected triazol-1-yl derivative with concentrated ammonium hydroxide (4.26 mL, 81.2 mmol) in dioxane at room temperature for 6 hours. After evaporation of the solvent the oil is stirred in half-saturated (at ice temperature) ammonia in methanol for 16 hours. The solvent is evaporated and 2'-deoxy-2'-fluorocytidine is crystallized from ethylacetate-methanol (v/v, 75:25) to give colorless crystals (1.24 g, 75 %). N-4-benzoyl-2'-deoxy-2'-fluorocytidine is prepared by selective benzylation with benzoic anhydride in anhydrous dimethylformamide, V. Bhat, et al. Nucleosides Nucleotides, Vol. 8, pp. 179-183 (1989). The 5'-O-(4,4'-dimethoxytrityl)-3'-O-(N,N-diisopropyl-b-cyanoethylphosphoramidite) can be prepared in accordance with Examples 65 and 66.

**EXAMPLE 69**

N-2-Phenoxyacetyl-2'-Deoxy-2'-fluoro-5'-O-(4,4'-dimethoxytrityl)-guanosine-3'-O-(N,N-diisopropyl- $\beta$ -cyanoethylphosphoramidite). (69)

2'-Deoxy-2'-fluoroguanosine is prepared according to the procedure of M. Ikehara, et al., Chemical and Pharmaceutical Bulletin Vol. 29, pp. 1034-1038 and pp.

- 92 -

3281-3285 (1981). 9- $\beta$ -D- arabinofuranosylguanine is utilized as starting material, M.J. Robins, Tetrahedron, Vol. 40, pp. 125-135 (1984). After 2'-deoxy- 2'- fluoro-guanosine is obtained, it is protected as the N-2-phen-  
oxyacetyl derivative, K.K. Ogilvie, Nucleic Acids Research, Vol. 17, pp. 3501-3517, can then be converted to the 5'-DMT-3'-phosphor-amidite as described in Examples 65 and 66.

**EXAMPLE 70**

10 N<sup>6</sup>-Benzoyl-[2'-deoxy-2'-cyano-5'-O-(4,4'-dimethoxytrityl)] adenosine-3'-O-(N,N-diisopropyl- $\beta$ -cyanoethylphosphoramidite) (70)

2'-Deoxy-2'-cyanoadenosine is prepared by the free radical replacement of the 2'-iodo group of 2'-deoxy-  
15 2'-iodo- 3',5'-O-(disiloxytetraisopropyl)-N<sup>6</sup>-benzoyl-adenosine according to a similar procedure described by K.E.B. Parkes and K. Taylor, Tetrahedron Letters, Vol. 29, pp. 2995-2996 (1988). 2'-Deoxy-2'- iodoadenosine was prepared by R. Ranganathan as described in Tetrahedron  
20 Letters Vol. 15, pp. 1291-1294 (1977), and disilylated as described by W.T. Markiewicz and M. Wiewiorowski in Nucleic Acid Chemistry, Part 3, pp. 222-231, Townsend, L.B.; Tipson, R.S. eds. (J. Wiley and Sons, New York, 1986). This material is treated with hexamethylditin,  
25 AIBN, and t-butylnisocyanate in toluene to provide protected 2'-deoxy-2'-cyanoadenosine. This material, after selective deprotection, is converted to its 5'-DMT-3'-phosphor- amidite as described in Examples 65 and 66.

**EXAMPLE 71**

2'-Deoxy-2'-cyano-5'-O-(4,4'-dimethoxytrityl)-uridine-3'-O-(N,N-diisopropyl- $\beta$ -cyanoethyl phosphoramidite). (71)

2'-Deoxy-2'-iodouridine (or 5-methyluridine), 3',5'-disilylated as described above, is converted to the  
35 2'-iodo derivative by triphenylphosphonium methyl iodide treatment as described by K.E.B. Parkes and K. Taylor,

- 93 -

Tetrahedron Letters Vol. 29, pp. 2995-2996 (1988). Application of free radical reaction conditions as described by K.E.B. Parkes and K. Taylor, Tetrahedron Letters, Vol. 29, pp. 2995-2996 (1988), provides the 2'-cyano group of the protected nucleoside. Deprotection of this material and subsequent conversion to the protected monomer as described in examples 65 and 66 above provides the requisite nucleic acid synthesizer material.

10 **EXAMPLE 72**

2'-Deoxy-2'-cyano-5'-O-(4,4'-dimethoxytrityl)-cytidine-3'-O-(N,N-diisopropyl- $\beta$ -cyanoethylphosphoramidite). (72)

2'-Deoxy-2'-iodocytidine is obtained via a conventional keto to amino conversion.

15 **EXAMPLE 73**

2'-Deoxy-2'-cyano-5'-O-(4,4'-dimethoxytrityl)-guanosine-3'-O-(N,N-diisopropyl- $\beta$ -cyanoethylphosphoramidite). (73)

2'-Deoxy-2'-cyanoguanosine is obtained by the displacement of the triflate group in the 2'-up position (arabino sugar) of 3',5'-disilylated N2-isobutryl-guanosine. Standard deprotection and subsequent reprotection such as through the methodologies of Examples 65 and 66 provides the title monomer.

25 **EXAMPLE 74**

**Preparation of 2'-Deoxy-2'-(trifluoromethyl) Modified Oligonucleotides.**

The requisite 2'-deoxy-2'-trifluoromethyribosides of nucleic acid bases A, G, U(T), and C are prepared by modifications of a literature procedure described by Q.-Y. Chen and S.W. Wu in the Journal of Chemical Society Perkin Transactions, pp. 2385-2387 (1989). Standard procedures, as described in Examples 65 and 66 are employed to prepare the 5'-DMT and 3'-phosphoramidites as listed below.



- 94 -

- A. N<sup>6</sup>-Benzoyl-[2'-deoxy-2'-(trifluoromethyl-5'-O-(4,4'-dimethoxytrityl)] adenosine-3'-O-(N,N-diisopropyl-β-cyanoethylphosphoramidite).
- 5 B. 2'-Deoxy-2'-(trifluoromethyl-5'-O-(4,4'-dimethoxy-trityl)- uridine-3'-O-(N,N-diisopropyl-β-cyano-ethylphosphoramidite).
- C. 2'-Deoxy-2'-(trifluoromethyl-5'-O-(4,4'-dimethoxy-trityl)-cytidine-3'-O-(N,N-diisopropyl-β-cyano-ethylphosphoramidite).
- 10 D. 2'-Deoxy-2'-(trifluoromethyl-5'-O-(4,4'-dimethoxy-trityl)-guanosine-3'-O-(N,N-diisopropyl-β-cyano-ethylphosphoramidite).

## EXAMPLE 75

## 15 Preparation of 2'-Deoxy-2'-(trifluoromethoxy) Modified Oligonucleotides.

The requisite 2'-deoxy-2'-O-trifluoromethoxyribosides of nucleic acid bases A, G, U(T), and C are prepared by modifications of literature procedures described by B.S. Sproat, et al., Nucleic Acids Research, Vol. 18, pp. 41-49 (1990) and H. Inoue, et al., Nucleic Acids Research, Vol. 15, pp. 6131-6148 (1987). Standard procedures, as described in Example 1A, are employed to prepare the 5'-DMT and 3'-phosphoramidites as listed below.

- A. N<sup>6</sup>-Benzoyl-[2'-deoxy-2'-(trifluoromethoxy)-5'-O-(4,4'-dimethoxytrityl)] adenosine-3'-O-(N,N-diisopropyl-β-cyanoethylphosphoramidite).
- 30 B. 2'-Deoxy-2'-(trifluoromethoxy)-5'-O-(4,4'-dimethoxytrityl)-uridine-3'-O-(N,N-diisopropyl-β-cyano-ethylphosphoramidite).
- C. 2'-Deoxy-2'-(trifluoromethoxy)-5'-O-(4,4'-dimethoxytrityl)-cytidine-3'-O-(N,N-diisopropyl-β-cyanoethylphosphoramidite).
- 35 D. 2'-Deoxy-2'-(trifluoromethoxy)-5'-O-(4,4'-dimethoxytrityl)-guanosine-3'-O-(N,N-diisopropyl-β-cyano-ethylphosphoramidite).

**EXAMPLE 76**

**Preparation of 2'-Deoxy-2'-(1-propoxy) Modified G, U (T) and C Oligonucleotides.**

The requisite 2'-deoxy-2'-O-propyl ribosides of nucleic acid bases G, U(T), and C are prepared by routine modification of examples 17 and 18. Also see, B.S. Sproat, et al., Nucleic Acids Research, Vol. 18, pp. 41-49 (1990) and H. Inoue, et al., Nucleic Acids Research Vol. 15, pp. 6131-6148 (1987). The procedures, as described in Examples 65 and 66 or examples 19 and 20 are employed to prepare the 5'-DMT and 3'-phosphoramidites as listed below.

- A. 2'-Deoxy-2'-(1-propoxy)-5'-O-(4,4'-dimethoxytrityl)-uridine-3'-O-(N,N-diisopropyl- $\beta$ -cyanoethylphosphoramidite).
- B. 2'-Deoxy-2'-(1-propoxy)-5'-O-(4,4'-dimethoxytrityl)-cytidine-3'-O-(N,N-diisopropyl- $\beta$ -cyanoethylphosphoramidite).
- C. 2'-Deoxy-2'-(1-propoxy)-5'-O-(4,4'-dimethoxytrityl)-guanosine-3'-O-(N,N-diisopropyl- $\beta$ -cyanoethylphosphoramidite).

**EXAMPLE 77**

**Preparation of 2'-Deoxy-2'-(vinylxy) Modified Oligonucleotides.**

The requisite 2'-deoxy-2'-O-vinyl ribosides of nucleic acid bases A, G, U(T), and C are prepared by modifications of the procedures of example 1. In this case 1,2-dibromoethane is coupled to the 2'-hydroxyl and subsequent dehydrobromination affords the desired blocked 2'-vinyl nucleoside. Standard procedures, as described in Examples 65 and 66 are employed to prepare the 5'-DMT and 3'-phosphoramidites as listed below.

- A. N<sup>6</sup>-Benzoyl-[2'-deoxy-2'-(vinylxy)-5'-O-(4,4'-dimethoxytrityl)] adenine-3'-O-(N,N-diisopropyl- $\beta$ -cyanoethylphosphoramidite).

- 96 -

- B. 2'-Deoxy-2'-(vinylloxy)-5'-O-(4,4'-dimethoxytrityl)-uridine-3'-O-(N,N-diisopropyl- $\beta$ -cyanoethylphosphoramidite).
- 5 C. 2'-Deoxy-2'-(vinylloxy)-5'-O-(4,4'-dimethoxytrityl)-cytidine-3'-O-(N,N-diisopropyl- $\beta$ -cyanoethylphosphoramidite).
- D. 2'-Deoxy-2'-(vinylloxy)-5'-O-(4,4'-dimethoxytrityl)-guanosine-3'-O-(N,N-diisopropyl- $\beta$ -cyanoethylphosphoramidite).

10 **EXAMPLE 78**

Preparation of 2'-Deoxy-2'-(allyloxy) Modified G, U(T), and C Oligonucleotides.

The requisite 2'-deoxy-2'-O-allyl ribosides of nucleic acid bases G, U(T), and C are prepared by modifications  
15 of the procedures of example 21. Also see B.S. Sproat, et al., Nucleic Acids Research Vol. 18, pp. 41-49 (1990) and H. Inoue, et al., Nucleic Acids Research Vol. 15, pp. 6131-6148 (1987). The procedures, described in Examples 65 and 66 or 23 and 24 are employed to prepare the 5'-[MT  
20 and 3'-phosphoramidites as listed below.

- A. N<sup>6</sup>-Benzoyl-[2'-deoxy-2'-(allyloxy)-5'-O-(4,4'-dimethoxytrityl)] adenosine-3'-O-(N,N-diisopropyl- $\beta$ -cyanoethylphosphoramidite).
- 25 B. 2'-Deoxy-2'-(allyloxy)-5'-O-(4,4'-dimethoxytrityl)-uridine-3'-O-(N,N-diisopropyl- $\beta$ -cyanoethylphosphoramidite).
- C. 2'-Deoxy-2'-(allyloxy)-5'-O-(4,4'-dimethoxytrityl)-cytidine-3'-O-(N,N-diisopropyl- $\beta$ -cyanoethylphosphoramidite).
- 30 D. 2'-Deoxy-2'-(allyloxy)-5'-O-(4,4'-dimethoxytrityl)-guanosine-3'-O-(N,N-diisopropyl- $\beta$ -cyanoethylphosphoramidite).

- 97 -

**EXAMPLE 79**

Chemical conversion of an thymine or cytosine (pyrimidine type base) to its  $\beta$ -D-2'-deoxy-2'-substituted erythropentofuranosyl nucleoside; 2'-substituted ribosylation).

The thymine or cytosine type analogs are trimethylsilylated under standard conditions such as hexamethyldisilazane (HMDS) and an acid catalyst (ie. ammonium chloride) and then treated with 3,5-O-ditoluoyl-2-deoxy-2-substituted- $\alpha$ -D-erythropento-furanosyl chloride in the presence of Lewis acid catalysts (ie. stannic chloride, iodine, boron tetrafluoroborate, etc.). A specific procedure has recently been described by J. N. Freskos, Nucleosides & Nucleotides Vol. 8, pp. 1075-1076 (1989) in which copper (I) iodide is the catalyst employed.

**EXAMPLE 80**

Chemical conversion of an adenine or guanine (purine type base) to its  $\beta$ -D-2'-deoxy-2'-substituted erythropentofuranosyl nucleoside; 2'-substituted ribosylation).

The protected purine type analogs are converted to their sodium salts via sodium hydride in acetonitrile and are then treated with 3,5-O-ditoluoyl-2-deoxy-2-substituted- $\alpha$ -D-erythropentofuranosyl chloride at ambient temperature. A specific procedure has recently been described by R.K. Robins et al., Journal of American Chemical Society Vol. 106, p. 6379 (1984).

**EXAMPLE 81**

Conversion of 2'-deoxy-2-substituted thymidines to the corresponding 2'-deoxy-2'-substituted cytidines (chemical conversion of an pyrimidine type 4-keto group to an 4-amino group).

The 3',5'-sugar hydroxyls of the 2'-modified nucleoside types are protected by acyl groups such as toluoyl, benzoyl, p-nitrobenzoyl, acetyl, isobutryl, trifluoroacetyl, etc. using standard conditions of the acid

- 98 -

chlorides or anhydrides and pyridine/dimethyl-amino pyridine solvent and catalyst. The protected nucleoside is now chlorinated with thionyl chloride or phosphoryl chloride in pyridine or other appropriate basic solvents.

- 5 The pyrimidine type 4-chloro groups are now displaced with ammonium in methanol. Deprotection of the sugar hydroxyls also takes place. The amino group is benzoylated by the standard two-step process of complete benzylation (sugar hydroxyls and amino group) and the
- 10 acyls are selectively removed by aqueous sodium hydroxide solution. Alternatively, the in situ process of first treating the nucleoside with chlorotrimethylsilane and base to protect the sugar hydroxyls from subsequent acylation may be employed. K. K. Ogilvie, Can J. Chem.
- 15 Vol. 67, pp. 831-839 (1989). Another conversion approach is to replace the pyrimidine type 4-chloro group with an 1,2,4-triazolo group which remains intact throughout the oligonucleotide synthesis on the DNA synthesizer and is displaced by ammonium during the ammonium hydroxide step
- 20 which removes the oligonucleotide from the CPG support and deprotection of the heterocycles. Furthermore, in many cases the pyrimidine type 4-chloro group can utilized as just described and replaced at the end of the oligonucleotide synthesis.

25 **EXAMPLE 82**

Methyl-5-(cyanomethyl)-1-(2'-deoxy-3,5-di-O-p-toluoyl-β-D-erythro-pentofuranosyl)imidazole-4-carboxylate (82-A) and 5-cyanomethyl-1-(2'-deoxy-β-D-erythro-pentofuranosyl)imidazole-4-carboxamide. (82-B)

- 30 A large scale synthesis of the methyl-5-(cyanomethyl)-1-(2'-deoxy-3,5-di-O-p-toluoyl-β-D-erythro-pentofuranosyl)imidazole-4-carboxylate (82-A), was undertaken according to the sodium-salt glycosylation procedure published by G.R. Revankar, et al. in J. Med. Chem. Vol.
- 35 27, pp. 1389-1396 (1984). In examples 83 to 85 below, the products of alkylation of this starting material are diastereomeric mixtures differing in configuration at the

- 99 -

alkylated carbon. As such, these mixtures exhibit duplicate resonances for each proton in the <sup>1</sup>H-NMR. The 5-cyanomethyl-1-(2'-deoxy-β-D-erythropentofuranosyl)-imidazole-4-carboxamide (82-B), the product of reaction of 82A with liquid ammonia at 100°C, was prepared according to this same literature reference.

**EXAMPLE 83**

Methyl 5-(cyano[nonyl]methyl)-1-(2'-deoxy-3,5-di-O-p-toluoyl-β-D-erythro-pentofuranosyl)imidazole-4-carboxylate (83).

A solution of 82-A (5.7 g, 11 mmol) in anhydrous acetonitrile (75 mL) was treated with sodium hydride (0.88 g, 60% in oil, washed with hexanes) at room temperature and under an atmosphere of argon. This suspension was stirred for 15 minutes and then treated with iodononane (7.5 mL, 37.4 mmol) using a syringe. The reaction mixture was stirred under these conditions for 6 hr; thin layer chromatography plates (ethyl acetate/hexanes, 3:2, v/v) showed the disappearance of starting material nucleoside and the appearance of two faster migrating products. The reaction was quenched with the addition of glacial acetic acid to pH 5 and then evaporated to dryness in vacuo to afford a yellow syrup. The syrup was dissolved in dichloromethane (150 mL) and the solution was washed with cold 0.1 N HCl, water, and then dried over magnesium sulfate. Filtration of the desiccant and evaporation of the solvent afforded a yellow gum which was flash-chromatographed on silica gel (120 g) using ethyl acetate-hexanes (1:4 then 1:1). Fractions corresponding to the alkylated products were pooled and evaporated to yield 83 as a yellowish foam, 2.98 g (47%). <sup>1</sup>H-NMR (Me<sub>2</sub>SO-d<sub>6</sub>): δ, 8.18 and 8.15 (s,s; C-2 H, 1 H); 6.48 and 6.37 (t,t; H-1', 1 H); 1.18 and 0.90 (2 m, nonyl, 19 H).

- 100 -

**EXAMPLE 84**

Methyl 5-(allyl[cyano]methyl)-1-(2'-deoxy-3,5-di-O-p-toluoyl- $\beta$ -D-erythro-pentofuranosyl)imidazole-4-carboxylate (84).

- 5 A solution of 82-A (5.0 g, 9.7 mmol) in anhydrous acetonitrile (75 mL) was treated with sodium hydride (0.46 g, 11.6 mmol) and then allyl bromide (2.5 mL, 29 mmol) in the manner described for 83. The reaction was worked up and chromatographed on silica gel (75 g) using  
10 the aforementioned solvent system to afford 84 as a yellowish foam, 3.66 g (68%).  $^1\text{H-NMR}$  ( $\text{Me}_2\text{SO-d}_6$ ):  $\delta$ , 8.15 and 8.13 (s,s; C-2 H, 1 H); 6.38 (m, H-1', 1 H); 5.75 and 5.08 (2 m, vinyl, 3 H).

**EXAMPLE 85**

- 15 Methyl 5-(benzyl[cyano]methyl)-1-(2'-deoxy-3,5-di-O-p-toluoyl- $\beta$ -D-erythro-pentofuranosyl)imidazole-4-carboxylate (85).

- A solution of 82-A (5.0 g, 9.6 mmol) in anhydrous acetonitrile (75 mL) was treated with sodium hydride  
20 (0.46 g, 11 mmol) under argon and stirred at room temperature for 15 minutes. The mixture was cooled to 4 °C in an ice bath and a solution of benzyl bromide (1.26 mL, 10.6 mmol) in acetonitrile (15 mL) was added dropwise over 70 min. The ice bath was removed and the reaction  
25 further stirred at room temperature for 2.5 hr. Workup of the reaction and purification of the products on silica gel (100 g) afforded 85 as a white foam, 3.4 g (58%).  $^1\text{H-NMR}$  ( $\text{Me}_2\text{SO-d}_6$ ):  $\delta$ , 8.10 and 8.05 (s,s; C-2 H, 1 H); 7.30-7.10 (m, phenyl, 5 H); 5.33 and 6.01 (t,t;  
30 H-1', 1 H).

**EXAMPLE 86**

5-(Cyano[nonyl]methyl)-1-(2'-deoxy- $\beta$ -D-erythropentofuranosyl)imidazole-4-carboxamid (86).

- The nonyl-imidazole nucleoside 83 (2.98 g, 4.6 mmol)  
35 was dissolved in anhydrous methanol (5 mL) and transferred to a stainless steel bomb. The solution was

- 101 -

cooled in a dry-ice /isopropanol bath and then treated with anhydrous liquid ammonia (45 mL). The bomb was sealed and then heated to 100 °C in an oil bath for 21 hr. TLC (ethyl acetate/methanol, 4:1, v/v) indicated a complete removal of the toluoyl protecting groups. Excess ammonia was evaporated at room temperature and the amber gum which resulted was flash chromatographed on silica gel (80 g) using ethyl acetate/methanol (95:5 then 9:1). Fractions corresponding to deblocked products were pooled and evaporated in vacuo to yield 86 as a white foam, 1.14 g (63%). <sup>1</sup>H-NMR (Me<sub>2</sub>SO-d<sub>6</sub>): δ, 8.09 and 8.05 (s,s; C-2 H, 1 H); 6.14 (t, H-1', 1H); 1.40-1.05 and 0.95 (2 m, nonyl, 19 H).

**EXAMPLE 87**

5-(Allyl[cyano]methyl)-1-(2'-deoxy-β-D-erythropentofuranosyl)imidazole-4-carboxamide (87).

The allyl-imidazole nucleoside 84 (3.95g, 7.03 mmol) was treated with liquid ammonia in a stainless steel bomb and heated to 100 °C for 8 hr. The products of this reaction were worked up and purified on silica gel (80 g) in a manner analogous to that for compound 86. The deblocked compound 87 was isolated as a white foam, 1.29 g (58%). <sup>1</sup>H-NMR (Me<sub>2</sub>SO-d<sub>6</sub>): δ, 8.09 and 8.07 (s,s; C-2 H, 1 H); 6.15 (t, H-1', 1 H); 5.75 and 5.10 (2 m, vinyl, 3 H).

**EXAMPLE 88**

5-(Benzyl[cyano]methyl)-1-(2'-deoxy-β-D-erythropentofuranosyl)imidazole-4-carboxamide (88).

The benzyl-imidazole nucleoside 85 (3.0 g, 4.93 mmol) was treated with liquid ammonia in a stainless steel bomb and heated to 100 °C for 6 hr. The products of this reaction were worked up and purified on silica gel (80 g) in a manner analogous to that for compound 86. The deblocked compound 88 was isolated as a white foam, 1.03 g (59%). <sup>1</sup>H-NMR (Me<sub>2</sub>SO-d<sub>6</sub>): δ, 8.03 and 8.04 (s,s;



- 102 -

C-2 H, 1 H); 7.25 (m, phenyl, 5 H); 6.17 and 6.07 (t,t; H-1', 1 H).

**EXAMPLE 89**

5-**(Cyanomethyl)-1-(2'-deoxy-5'-O- dimethoxytrityl-8-D-erythropentofuranosyl)imidazole-4- carboxamide (89).**

The nucleoside 82-B (1.95 g, 7.3 mmol) was dried by coevaporation with pyridine (30 mL). The gum which resulted was dissolved in anhydrous pyridine under argon and then treated with dimethoxytrityl chloride (2.90 g, 12.4 mmol). The mixture was stirred at room temperature for 2 hr after which TLC (ethyl acetate:methanol, 19:1, v/v) indicated complete conversion of starting material. Tritylated products were visualized as orange spots using H<sub>2</sub>SO<sub>4</sub> fumes. The reaction was quenched with the addition of methanol (2 mL) followed by stirring for 15 min. The mixture was evaporated in vacuo to afford a thick orange syrup which was coevaporated with toluene (3 x 25 mL). The syrup was flash chromatographed on silica gel (100 g) using a stepwise gradient of methanol in 1% Et<sub>3</sub>N/CH<sub>2</sub>Cl<sub>2</sub> (0 to 5 % methanol). The appropriate fractions were pooled and evaporated to yield 89 as a white foam, 3.64 g (87%). <sup>1</sup>H-NMR (Me<sub>2</sub>SO-d<sub>6</sub>): δ, 7.96 (s, C-2 H, 1 H); 6.85-7.35 (m, DMT, 13 H); 6.13 (t, H-1', 1 H).

**EXAMPLE 90**

5-**(Cyano[nonyl]methyl)-1-(2'-deoxy-5'-O-dimethoxytrityl-8-D-erythropentofuranosyl)imidazole-4-carboxamide (90).**

The nucleoside 86 (1.20 g, 3.1 mmol) was thoroughly dried by coevaporation with anhydrous pyridine (30 mL). The syrup which resulted was redissolved in anhydrous pyridine under argon and treated with dimethoxytrityl chloride (1.0 g, 3.1 mMol). The reaction mixture was stirred at room temperature for 3.5 hr, after which time TLC (ethyl acetate) indicated complete conversion of the starting material. The reaction mixture was treated with 2 mL of anhydrous methanol, stirred for 15 minutes and then evaporated in vacuo to afford a bright orange syrup.

- 103 -

This syrup was coevaporated with toluene (2 X 50 mL) and then flash chromatographed on silica (80 g) using a step-wise gradient of methanol in dichloromethane/1% triethylamine (0 to 3% methanol). The appropriate fractions were pooled and evaporated in vacuo to yield the dimethoxytritylated compound 90 as a white foam, 1.46 g (69%).  
1H-NMR (Me<sub>2</sub>SO-d<sub>6</sub>):  $\delta$ , 7.98 and 7.93 (s,s; C-2 H, 1 H); 7.30 and 6.92 (2 m, DMT, 13 H); 6.21 (t, H-1', 1 H); 1.20 and 0.92 (2 m, nonyl, 19 H).

## 10 EXAMPLE 91

5-(Allyl[cyano]methyl)-1-(2'-deoxy-5'-O-dimethoxytrityl- $\beta$ -D-erythropentofuranosyl)imidazole-4-carboxamide (91).

The nucleoside 87 (1.25 g, 4.08 mmol) was dried by coevaporation with pyridine and then redissolved in anhydrous pyridine (50 mL) and treated with dimethoxytrityl chloride (1.38 g, 4.08 mmol) under an atmosphere of argon. The reaction was stirred for 2.5 hr and then worked up and products isolated by flash chromatography on silica gel (90 g) in a manner analogous to compound 89. The appropriate fractions were pooled and evaporated in vacuo to yield dimethoxytritylated compound 91 as a white foam, 1.86 g (75%). 1H-NMR (Me<sub>2</sub>SO-d<sub>6</sub>):  $\delta$ , 7.98 and 7.95 (s,s; C-2 H, 1 H); 7.25 and 6.93 (2 m, DMT, 13 H); 6.21 (m, H-1', 1 H); 5.78 and 5.10 (2 m, vinyl, 3 H).

## 25 EXAMPLE 92

5-(Benzyl[cyano]methyl)-1-(2'-deoxy-5'-O-dimethoxytrityl- $\beta$ -D-erythropentofuranosyl)imidazole-4-carboxamide (92).

The nucleoside 88, 930 mg (2.6 mmol) was dried by coevaporation with pyridine and then redissolved in anhydrous pyridine (50 mL) and treated with dimethoxytrityl chloride (884 mg, 2.6 mmol) under an atmosphere of argon. The reaction was stirred for 4 hr and then worked up and the products isolated by flash chromatography on silica gel (80 g) in a manner analogous to compound 89. The appropriate fractions were pooled and evaporated in

- 104 -

vacuo to yield 1.50 g of 92 as a pinkish foam, (87%).  
1H-NMR (Me<sub>2</sub>SO-d<sub>6</sub>):  $\delta$ , 7.70 (s, C-2 H, 1 H); 7.40 and 6.70  
(2m; DMT, phenyl; 18 H); 6.10 (t, H-1', 1 H).

**EXAMPLE 93**

5 5-(Cyanomethyl)-1-(5'-O-dimethoxytrityl-3'-O-[2-cyano-  
ethyl-N,N-diisopropyl]phosphoramidite-2'-deoxy- $\beta$ -D-  
erythropentofuranosyl)imidazole-4-carboxamide (93).

The tritylated nucleoside 89 (1.82 g, 3.2 mmol) was  
dissolved in anhydrous tetrahydrofuran (50 mL) under  
10 argon and then treated with diisopropyl ethylamine (1  
mL). The solution was cooled to 4 °C in an ice bath and  
then treated with 2-cyanoethyl-N,N-diisopropyl phosphoro-  
chloridate (1.2 g, 5.12 mmol) in one portion. The ice  
15 bath was removed and the reaction mixture was stirred at  
room temperature for 3 hr. At the end of this time, TLC  
(CH<sub>2</sub>Cl<sub>2</sub>/1% MeOH/1% Et<sub>3</sub>N) indicated complete conversion of  
starting material. Reaction products were visualized  
using H<sub>2</sub>SO<sub>4</sub> fumes. The reaction mixture was evaporated  
in vacuo to afford a thick syrup which was immediately  
20 redissolved in dichloromethane (100 mL) and washed with  
cold, saturated sodium bicarbonate (2 x 50 mL) and brine  
(50 mL). The organic layer was dried over magnesium  
sulfate, filtered and evaporated to yield a yellowish  
foam (2.8 g). This foam was flash chromatographed on  
25 silica gel (75 g) using a stepwise gradient of ethyl  
acetate/hexanes (1:4 to 1:1) containing 1% Et<sub>3</sub>N. The  
appropriate fractions were pooled and evaporated to yield  
93 as a white foam, 1.65 g (59%). An aliquot of this  
material was precipitated by dissolving the foam in a  
30 small volume of dichloromethane and adding it to a large  
proportion (1:50) of hexanes. 1H-NMR (CD<sub>3</sub>CN):  $\delta$ , 7.78  
and 7.72 (s,s; C-2 H, 1H); 7.25 and 6.95 (m, DMT, 13 H);  
6.12 (t, H-1', 1H). 31P-NMR (CD<sub>3</sub>CN):  $\delta$ , 150.02, 149.98.

## EXAMPLE 94

5-(Cyano[nonyl]methyl)-1-(5'-O-dimethoxytrityl-3'-O-[2-cyanoethyl-N,N-diisopropyl]phosphoramidite-2'-deoxy-β-D-erythropentofuranosyl)imidazole-4-carboxamide (94).

- 5 The tritylated nucleoside 90, 1.46 g (2.1 mmol) was dissolved in anhydrous tetrahydrofuran (50 mL) under argon and then treated with diisopropyl ethylamine (1.5 mL). This mixture was cooled to 4 °C in an ice bath and then treated with 2-cyanoethyl-N,N-diisopropylamino phosphorochloridite (488 mg, 2.1 mmol) in one portion. The ice bath was removed and the reaction was further stirred at room temperature for a total of 3 hr. Additional phosphorochloridite (48 mg) was added at the end of the first and second hours of reaction. At the end of this time, TLC (CH<sub>2</sub>Cl<sub>2</sub>/3% MeOH/1% Et<sub>3</sub>N) indicated complete conversion of starting material. The reaction mixture was worked up as described for 93 and the products purified on silica gel (80 g) using a stepwise gradient of ethyl acetate/hexanes (2:3 to 3:2) containing 1% Et<sub>3</sub>N.
- 10 20 The appropriate fractions were pooled and evaporated to afford 94 as a colorless foam, 1.37 g (73%). An aliquot of this material was precipitated using a procedure described for compound 93. <sup>1</sup>H-NMR (CD<sub>3</sub>CN): δ, 7.75 (s, C-2 H); 7.25 and 6.85 (2 m, DMT, 13 H), 6.25 and 6.20 (2 m, H-1', 1 H). <sup>31</sup>P-NMR (CD<sub>3</sub>CN): δ, 150.1, 150.0 and 149.9.

## EXAMPLE 95

- 5-(Allyl[cyano]methyl)-1-(5'-O-dimethoxytrityl-3'-O-[2-cyanoethyl-N,N-diisopropyl]phosphoramidite-2'-deoxy-β-D-erythro-pentofuranosyl)imidazole-4-carboxamide (95).

- 30 The tritylated nucleoside 91, 1.86 g (3.1 mmol) was dissolved in anhydrous THF (50 mL) and then treated with diisopropyl ethylamine (1.5 mL) and 2-cyanoethyl N,N-diisopropylamino phosphochloridite (705 mg, 3.1 mmol) in a manner analogous to compound 94. The mixture was stirred at room temperature for a total of 3 hr. Additional phosphochloridite (140 mg) was added at the end

- 106 -

of 2 hr reaction time. The reaction was worked up and the products isolated by flash chromatography on silica gel (80 g) using a stepwise gradient of ethyl acetate/hexanes (2:3 to 4:1, v/v). The appropriate fractions were pooled and evaporated to yield 2.18 g (88%) of 95 as a white solid foam. An aliquot of this material was precipitated using a procedure described for compound 93. <sup>1</sup>H-NMR (CD<sub>3</sub>CN): δ, 7.77 and 7.75 (s,s; C-2 H, 1 H); 7.45 and 6.85 (2m, DMT, 13 H); 6.25 and 6.20 (t,t; H-1', 1 H). <sup>31</sup>P-NMR (CD<sub>3</sub>CN): δ, 150.05, 149.98 and 149.85.

**EXAMPLE 96**

5-(Benzyl[cyano]methyl)-1-(3'-O-dimethoxytrityl-3'-O-[2-cyanoethyl-N,N-diisopropyl]phosphoramidite-2'-deoxy-β-D-erythro-pentofuranosyl)imidazole-4-carboxamide (96).

The tritylated nucleoside 92, 1.50 g (2.3 mmol) was dissolved in anhydrous THF (50 mL) and then treated with diisopropyl ethylamine (1.5 mL) and 2-cyanoethyl-N,N-diisopropylamino phosphochloridite (539 mg, 2.3 mmol) in a manner analogous to compound 93. The mixture was stirred at room temperature for a total of 3 hr. Additional phosphorochloridite (110 mg) was added at the end of 2 hr reaction time. The reaction was worked up and the products isolated by flash chromatography on silica gel (40 g) using a stepwise gradient of ethyl acetate/hexanes (1:4 to 3:2, v/v). The appropriate fractions were pooled and evaporated to yield 96, 1.03 g (55%) as a colorless foam. An aliquot of this material was precipitated using a procedure described for compound 93. <sup>1</sup>H-NMR (CD<sub>3</sub>CN): δ, 7.72 (s,s; C-2 H; 1 H); 7.30 and 6.80 (2 m; DMT, benzyl; 18 H); 6.10 (m, H-1', 1 H). <sup>31</sup>P-NMR (CD<sub>3</sub>CN): δ, 150.00, 149.90 and 149.85.

- 107 -

**EXAMPLE 97****Synthesis of Oligomers.**

Oligomers incorporating the nucleotides of examples 93, 94, 95 and 96 were made using standard phosphoramidite chemistries on an Applied Biosystems 380B synthesizer. Average coupling efficiencies were 94% using a method which left the 5'-dimethoxytrityl group of each oligonucleotide on. After cleavage from the solid support, the oligomers were treated with excess concentrated ammonium hydroxide and then heated in a sealed vessel at 55°C for a minimum of 15 hr. This procedure removed all protecting groups from A, G and C bases and caused cyclization of the 5-cyano[alkyl]methyl imidazole to the 3-deaza-3-alkyl (or aryl) heterocycles. For each nucleotide, verification of this cyclization was made from the analysis of a 1H-NMR spectrum of a TG3C trimer containing a single 3-deaza-3-substituted 2'-deoxyguanosine base. Purification of these trityl-on oligomers was performed using a C-4 Waters Prepak cartridge (10 cm) using a gradient elution of acetonitrile in 50 mM triethyl ammonium acetate (pH 7.0) (4 to 48 % over 60 min). The trityl group was removed by a treatment with 15 % acetic acid and the oligomers were then precipitated from 70 % ethanol.

**EXAMPLE 98**

5-(Cyano[propylphthalimido]methyl)-1-(5'-O-dimethoxytrityl-3'-O-[2-cyanoethyl-N,N-diisopropyl]phosphoramidite-2'-deoxy-β-D-erythro-pentofuranosyl-imidazole-4-carboxamide (98).

In a manner as per example 83 compound 82-A can be treated with N-(3-bromopropyl)phthalimide instead of iodononane. After work-up as per example 83 the product can be treated as per examples 89 and 93 to yield the title compound. In addition to ring closure upon cleavage from the oligonucleoside synthesizer solid support, the phthalimide is also cleaved to amino as per other above examples. Further the alkylamino product can

- 108 -

be chain extended to a polyalkylamine as per examples 57 and 58.

**EXAMPLE 99**

5-(Cyano[imidazo-1-yl(propyl)]methyl)-1-(5'-O-dimethoxy-trityl-3'-O'[2-cyanoethyl-N,N-diisopropyl]phosphor-amidite-2'-deoxy- $\beta$ -D-erythro-pentofuranosyl-imidazole-4-carboxamide (99).

In a manner as per example 83 compound 82-A can be treated with 1-(3-bromopropyl)imidazole instead of iodo-nonane. After work-up as per example 83 the product can be treated as per examples 89 and 93 to yield the title compound.

**EXAMPLE 100**

5-(Cyano[anthracen-2-yl(propyl)]methyl)-1-(5'-O-dimethoxy-trityl-3'-O'[2-cyanoethyl-N,N-diisopropyl]phosphoramidite-2'-deoxy- $\beta$ -D-erythro-pentofuranosyl-imidazole-4-carboxamide (100).

In a manner as per example 83 compound 82-A can be treated with 2-(3-bromopropyl)anthracene instead of iodo-nonane. After work-up as per example 83 the product can be treated as per examples 89 and 93 to yield the title compound.

**EXAMPLE 101**

2-chloro-6-allyloxy-9-2'-deoxypurine (101).

To a stirred suspension of 2,6-dichloro-9-(2'-deoxy-3',5'-di-o-toluy)-purine (10.3g, 19.04 mmol) in allyl alcohol (150ml) was added sodium hydride (0.8g, 20mmol) in small portions over 10 minutes period at room temperature. After the addition of sodium hydride, the reaction mixture was placed in a preheated oil bath at 55°C for 20 minutes with exclusion of moisture. The reaction mixture was cooled, filtered and washed with allyl alcohol (100ml). To the filtrate IRC-50 (weakly acidic) H<sup>+</sup> resin was added until the pH of the solution becomes 4-5. The resin was filtered, washed with methanol (100 ml) and the

- 109 -

filtrate was evaporated to dryness. The residue was adsorbed on silica gel (10g, 60-100 mesh) and evaporated to dryness. The dried silica gel was placed on top of silica column (5x25 cm, 100-250 mesh) packed in dichloromethane. The column was eluted with  $\text{CH}_2\text{Cl}_2 \rightarrow$  acetone. The fractions having the product was pooled together and evaporated to dryness to give 6g (96%) of the title compound as foam.

**EXAMPLE 102**

10 2-chloro-2'-deoxy inosine (102).

A mixture of (101) (6g, 18.4 mmol), Pd/C (10%, 1g) and triethyl amine (1.92g, 19 mmol) in ethyl alcohol (200 ml) was hydrogenated at atmospheric pressure during 30 minutes periods at room temperature. The reaction mixture was filtered, washed with MeOH (50 ml) and the filtrate evaporated to dryness. The product (5.26g, 100%) was found to be moisture sensitive and remains as an oil. The oil was used as such for further reaction without purification.

20 **EXAMPLE 103**

$\text{N}_2$ -[Imidazol-1-yl-(propyl)]-2'-deoxyguanosine (103)

A solution of the nucleoside 102 (10.31g, 36mmol) and 1-(3-amino-propyl) imidazol (9g, 72mmole) in 2-methoxy-ethanol (60ml) was heated in a steel bomb at 120°C (oil bath) for 24 hrs. The bomb was cooled to 0°C, opened carefully and the precipitated solid was filtered. The solid was washed with methanol (50ml) and acetone (50ml), and dried over sodium hydroxide to give 9g (67%) of pure 103. A small amount was recrystallized from DMF for analytical purposes. mp 245-47°.



- 110 -

## EXAMPLE 104

$N_2,3',5'$ -Tn-0-isobutyryl- $N_2$ -[imidazol-1-yl(propyl)]-2'-deoxyguanosine (104)

To a well dried solution of the substrate 103 (1.5g, 4mmole) and triethylamine (1.62g, 16mmol) in dry pyridine (30ml) and dry DMF (30ml) was added isobutyryl chloride (1.69g, 16mmol) at room temperature. The reaction mixture was allowed to stir at room temperature for 12 hrs and evaporated to dryness. The residue was partitioned between dichloromethane and water (100ml) and extracted with  $CH_2Cl_2$  (2x200ml). The organic extract was washed with brine (100ml) and dried over anhydrous magnesium sulfate. The dried organic extract was evaporated to dryness and the residue was purified over flash chromatography using  $CH_2Cl_2 \rightarrow$  MeOH as the eluent. The pure fractions were pooled and evaporated to dryness to give 1.8g (77%) of the title compound as colorless crystals from  $CH_2Cl_2$ /MeOH. mp 210-12°C.

## EXAMPLE 105

6-0-[2-nitrophenyl, ethyl]- $N_2,3',5'$ -Tn-0-isobutyryl- $N_2$ -[imidazol-1-yl(propyl)]-2'-deoxyguanosine (105).

To a stirred solution of 104 (2.0g, 3.42mmol), triphenyl phosphine (2.58g, 10.26mmol) and p-nitrophenyl ethanol (1.72g, 10.26mmol) in dry dioxane was added dichthylazodicarboxylate (1.78g, 10.26mmol) at room temperature. The reaction mixture was stirred at room temperature for 12h and evaporated to dryness. The residue was purified by flash chromatography over silica gel using  $CH_2Cl_2 \rightarrow$  acetone as the eluant. The pure fractions were pooled together and evaporated to dryness to give 2.4g (96%) of 6 as amorphous solid.

- 111 -

## EXAMPLE 106

6'-O-[2-(4-nitrophenyl)-ethyl]-N<sub>2</sub>-isobutyryl-N<sub>2</sub>-[imidazol-1-yl-(propyl)-2'-deoxyguanosine (106).

A stirred solution of 105 (9g, 12.26 mmol) in methanol  
5 (250ml) was treated with ammonium hydroxide (30%, 150ml) at room temperature. The reaction mixture was stirred at room temperature for 12h and evaporated to dryness. The residue was purified by flash chromatography over silica gel using CH<sub>2</sub>Cl<sub>2</sub> → acetone as the eluent. The pure  
10 fractions were pooled together and evaporated to dryness to give 2.4g (96%) of 105 as amorphous solid.

## EXAMPLE 107

5'-O-(4,4'-dimethoxytrityl)-6'-O-[2-(4-nitrophenyl)ethyl]-N<sub>2</sub>-isobutyryl-N<sub>2</sub>-[imidazol-1-yl(propyl)-2'-deoxyguanosine  
15 (107).

The substrate 106 (5.94g, 10mmol) was dissolved in dry pyridine (75ml) and evaporated to dryness. This was repeated three times to remove traces of moisture. To this well dried solution of the substrate in dry pyridine  
20 (100ml) was added dry triethylamine (4.04g, 40mmol), N,N-dimethylamino pyridine (10.14g, 30mmol) at room temperature. The reaction mixture was stirred at room temperature for 12 hrs under argon atmosphere. Methanol (50ml) was added and the stirring was continued for 15 minutes.  
25 The mixture was evaporated to dryness. The residue was purified by flash chromatography over silica gel using CH<sub>2</sub>Cl<sub>2</sub> → acetone containing 1% triethylamine as the eluant. The pure fractions were pooled together and evaporated to dryness to give 7.2g (80%) of (107) as  
30 colorless foam.

- 112 -

## EXAMPLE 108

3'-O-[N,N-diisopropylamino) ( $\beta$ -cyanoethoxy)phosphanyl]-  
5'-O-(4,4'-dimethoxytrityl)-6-O-[2-(4-nitrophenyl)ethyl]-  
N<sub>2</sub>-isobutyryl-N<sub>2</sub>-[imidazol-1-yl(propyl)-2'-deoxyguanosine  
5 (108).

The substrate 107 (2.5g, 2.7mmol) was dissolved in dry  
pyridine (30ml) and evaporated to dryness. This was  
repeated three times to remove remaining traces of water  
and dried over solid sodium hydroxide overnight. This  
10 dried 107 was dissolved in dry dichloromethane (30ml) and  
cooled to 0°C under argon atmosphere. To this cold stirred  
solution was added N,N-diisopropylethylamine (0.72g,  
5.6mmol) followed by ( $\beta$ -cyanoethoxy)chloro(N,N-diiso-  
propylamino) phosphane (1.32g, 5.6mmol) dropwise over a  
15 period of 15 minutes. The reaction mixture was stirred at  
0°C for 1 hr and at room temperature for 2 hr. The reac-  
tion mixture was diluted with dichloromethane (100ml) and  
washed with brine (50ml). The organic extract was dried  
over anhydrous magnesium sulfate and evaporated to dry-  
20 ness. The residue was purified by flash chromatography  
over silica gel using hexane  $\rightarrow$  acetone containing 1%  
triethylamine as the eluant. The main fractions were col-  
lected and evaporated to dryness. The residue was dis-  
solved in dry dichloro-methane (10ml) and added dropwise,  
25 into stirred solution of hexane (15--ml), during 30  
minute period. After the addition, the stirring was  
continued for an additional 1hr at room temperature under  
argon. The precipitated solid was filtered and dried over  
solid NaOH under vacuum overnight to give 2g (65%) of the  
30 title compound as colorless powder.

## EXAMPLE 109

N<sub>2</sub>-[imidazol-1-yl(propyl)]-2'-deoxyadenosine (109)

A suspension of 2-chloro-2'-deoxyadenosine (10.68g,  
35 37.4 mmole) and 1-(3-aminopropyl)imidazole (12.5g, 100  
mmole) in 2-methoxyethanol (80 ml) was heated at 125 °C  
for 45 hrs in a steel bomb. The bomb was cooled to 0 °C,  
opened carefully and evaporated to dryness. The residue

- 113 -

was co-evaporated several times with ethanol and toluene. The residue was dissolved in ethanol, which on cooling gave a precipitate. The precipitate was filtered and dried. The filtrate was evaporated to dryness and the  
5 residue was carried over to the next reaction without characterization.

**EXAMPLE 110**

3',5'-O-(tetraisopropylidisiloxane-1,3-diyl)-N<sub>2</sub>-  
10 [imidazole-1-yl(propyl)]-2'-deoxyadenosine (110).

The crude product (14.03 g) from example 109 was dissolved in dry DMF (100 ml) and dry pyridine (50 ml) and evaporated to dryness. This was repeated three times to remove all the water. The dried substrate was dis-  
15 solved in dry DMF (75 ml) and dry pyridine (75 ml) and allowed to stir at room temperature under an argon atmosphere. To this stirred solution was added dry triethylamine (10.1 g, 100 mmol) and dichloro-1,1,3,3-tetra-isopropylidisiloxane (Tipsicl) (15.7 g, 50 mmol)  
20 during a 15 minute period. After the addition of Tipsicl, the reaction mixture was stirred at room temperature overnight. The reaction mixture was evaporated to dryness. The residue was mixed with toluene (100 ml) and evaporated again. The residue was  
25 purified by flash chromatography over silica gel using methylene chloride ---> methanol as the eluent. The pure fractions were pooled and evaporated to dryness to give 12.5 g (54%) of 110 as an amorphous powder.

**EXAMPLE 111**

3',5'-O-(tetraisopropylidisiloxane-1,3-diyl)-N<sub>2</sub>,N<sub>6</sub>-di-isobutyryl-N<sub>2</sub>-[imidazole-1-yl(propyl)]-2'-deoxyadenosine  
(111).

A solution of compound 110 (12.0 g, 19.42 mmol) was  
35 allowed to stir at room temperature with triethylamine (10.1 g, 100 mmol) under an argon atmosphere. To this stirred solution in pyridine (100 ml) was added isobutyryl chloride (6.36 g, 60 mmol) dropwise during a 15

- 114 -

minutes period. The reaction mixture was stirred under argon for 10 hrs and evaporated to dryness. The residue was partitioned between dichloromethane/water and extracted with dichloromethane (2 X 150 ml). The organic  
5 extract was washed with brine (50 ml) and dried over anhydrous magnesium sulfate. The organic extract was evaporated to dryness and the residue was purified by flash chromatography over silica gel using methylene chloride ---> acetone as the eluent to give pure 111 as  
10 a foam.

## EXAMPLE 112

$N_2, N_6$ -di-isobutyryl- $N_2$ -[imidazol-1-yl(propyl)]-2'-deoxyadenosine (112).

15 The title compound will be prepared by exposure of compound 111 with tetrabutylammonium fluoride (5 equiv.) in dry tetrahydrofuran at room temperature. The reaction mixture on evaporation and purification by silica column chromatography will yield 112.

20

## EXAMPLE 113

5'-O-(4,4'-dimethoxytrityl)- $N_2, N_6$ -di-isobutyryl- $N_2$ -[imidazol-1-yl(propyl)]-2'-deoxyadenosine (113).

25 Compound 113 is prepared from compound 112 as per example 107 above.

## EXAMPLE 114

3'-O-[(N,N-diisopropylamino)( $\beta$ -cyanoethoxy)prospanyl]-  
5'-O-(4,4'-dimethoxytrityl)- $N_2, N_6$ -di-isobutyryl- $N_2$ -[imid  
30 azol-1-yl(propyl)]-2'-deoxyadenosine (113).

The phosphoramidite monomer 113 is prepared from compound 112 as per the procedure of example 109. The crude product is purified by silica gel chromatography.

- 115 -

**EXAMPLE 115****N<sub>2</sub>-nonyl-2'-deoxyguanosine (115).**

The compound 115 is prepared from compound 102 (9.5 g, 22 mmol) and nonylamine (9.58 g, 67 mmole) as per example 103. The crude product was evaporated to dryness and the residue was co-evaporated with a mixture of ethanol and toluene (three times). The resulting residue was dissolved in ethanol and cooled to yield a small amount of product. It was filtered and the filtrate evaporated to dryness. The residue was used without characterization.

**EXAMPLE 116****3',5',N<sub>2</sub>-triisobutyryl-N<sub>2</sub>-nonyl-2'-deoxyguanosine (116)**

The title compound 116 was prepared as per the procedure of example 111 utilizing compound 115 (18 g, 32.9 mmol), TEA (30.3 g, 300 mmole), isobutyryl chloride (21.2 g, 200 mmole) and dry pyridine (150 ml). The crude product was purified by flash chromatography over silica gel using methylene chloride ---> ethyl acetate as the eluent. The pure product was obtained as a foam in 40% yield.

**EXAMPLE 117****N<sup>2</sup>-isobutyryl-N<sub>2</sub>-nonyl-2'-deoxyguanosine (117).**

A stirred solution of compound 116 (5 g, 6.61 mmol) in methanol (100 ml) was treated with concentrated ammonium hydroxide (30%, 100 ml) at room temperature. After 8 hours, the reaction mixture was evaporated to dryness and the residue was purified by flash chromatography using methylene chloride ---> methanol as the eluent. The pure fractions were pooled together and evaporated to a residue. The residue was crystallized from methylene chloride/acetone to give 1.5 g (49%) of 117.

- 116 -

**EXAMPLE 118**

5'-O-(4,4'-dimethoxytrityl)-N<sub>2</sub>-isobutyryl-N<sub>2</sub>-nonyl-2'-deoxyguanosine (118).

Compound 118 was prepared from compound 117 by following the procedure of example 107 utilizing compound 117 (2.2 g, 4.75 mmole), DMTCl (1.69 g, 5 mmole), dry TEA (1.01 g, 10 mmole) and dry pyridine (70 ml). The crude compound was purified using flash chromatography and methylene chloride/methanol/TEA as the eluent. 2 g (55%) of pure product was obtained as a foam.

**EXAMPLE 119**

3'-O-[(N,N-diisopropylamino)( $\beta$ -cyanoethoxy)prospanyl]-5'-O-(4,4'-dimethoxytrityl)-N<sub>2</sub>,N<sub>6</sub>-diisobutyryl-N<sub>2</sub>-nonyl-2'-deoxyadenosine (119).

The phosphoramidite monomer compound 119 is prepared from 118 as per the procedure of example 109. The crude product is purified by silica gel chromatography.

**EXAMPLE 120**

3',5'-O-(tetraisopropylidisiloxane-1,3-diyl)-2-chloro-2'-deoxyadenosine (120).

Compound 120 was prepared from 2-chloro-2'-deoxyadenosine as per the procedure of example 110 utilizing 2-chloro-2'-deoxyadenosine (4.3 g, 15.09 mmol), 1,3-dichloro-1,1,3,3-tetraisopropylidisiloxane (4.74 g, 15.1 mmol), dry TEA (3.05 g, 30.2 mmol) and dry pyridine (100 ml). The pure product was purified by flash chromatography using methylene chloride ---> acetone as the eluent to give 7.3 g (92%) of pure 120 as an amorphous solid.

**EXAMPLE 121**

3',5'-O-(tetraisopropylidisiloxane-1,3-diyl)-2-chloro-N<sub>6</sub>-benzoyl-2'-deoxyadenosine (120).

A well dried solution of compound 120 (8 g, 15 mmol) in dry pyridine was allowed to react with triethylamine (4.55 g, 45 mmol) and benzoyl chloride (6.3 g, 45 mmol)

- 117 -

at room temperature for 12 hrs under an argon atmosphere. The reaction mixture was evaporated to dryness and the residue partitioned between methylene chloride/water and extracted in methylene chloride (2 X 150 ml). The organic extract was washed with brine (60 ml), dried and evaporated to dryness. The residue on purification by silica column gave 8.2 g (86%) of 121 as a foam.

**EXAMPLE 122**

10 **N<sub>6</sub>-benzoyl-2-chloro-2'-deoxyadenosine (122).**

To a stirred solution of compound 121 (7.9 g, 12.5 mmol) in dry THF (100 ml) was added a 1 M solution of tetrabutylammonium fluoride in THF (50 ml), 5 mmol) slowly during a 15 minute period at room temperature. The reaction was stirred for an additional 6 hrs and evaporated to dryness. The residue on purification by silica gel chromatography using methylene chloride ---> acetone gave 3.88 g (80%) of pure 122.

20 **EXAMPLE 123**

**5'-O-(4,4'-dimethoxytrityl)-N<sub>6</sub>-benzoyl-2-chloro-2'-deoxyadenosine (123)**

Compound 123 was prepared from compound 122 as per the procedure of example 107 using compound 122 (2.5 g, 6.43 mmol), DMTC1 (2.37 g, 7 mmol), dry TEA (0.71 g, 7 mmol) and dry pyridine (100 ml). The crude product was purified by flash chromatography using methylene chloride ---> ethyl acetate containing 1% TEA as the eluent to give 3 g (68%) of pure 123 as a foam.

30

**EXAMPLE 124**

**3'-O-[(N,N-diisopropylamino)(β-cyanoethoxy)phosphanyl]-5'-O-(4,4'-dimethoxytrityl)-N<sub>6</sub>-benzoyl-2-chloro-2'-deoxyadenosine (124)**

35 The phosphoramidite monomer 124 is prepared from compound 123 as per the procedure of example 109 using compound 123 (2.4 g, 3.47 mmol), N,N-diisopropyl ethylamine (1.22 ml, 7 mmol), 2-cyanoethyl-N,N-diisopropyl-



- 118 -

chloro-phosphoramidate (1.65 g, 7 mmol) and dry methylene chloride (30 ml). The crude product was purified by flash chromatography using hexane ---> ethyl acetate containing TEA (1%) as the eluent. The pure fractions  
5 were pooled together and evaporated to dryness. The residue was dissolved in 10 ml of dry methylene chloride and added dropwise into a well stirred hexane (1500 ml) under argon atmosphere. After the addition, stirring was continued for an additional 1 hr and the precipitated  
10 solid was filtered, washed with hexane and dried over solid sodium hydroxide for 3 hrs. The dried powder showed no traces of impurity on  $^{31}\text{P}$  spectrum.

**EXAMPLE 125**

15  $\text{N}_2$ -[imidazol-4-yl(ethyl)]-2'-deoxyguanosine (125).

A mixture of 2-chloro-2'-deoxyinosine (6.0 g, 20.97 mmol) and histamine (4.4 g, 40 mmol) in 2-methoxyethanol (60 ml) was heated in a steel bomb at 110 °C for 12 hrs. The bomb was cooled to 0 °C, opened carefully and the  
20 precipitated solid was filtered. The solid was washed with methanol and acetone and crystallized from DMF/water to give 6 g (79%) of pure 125. mp 220-222 °C.

**EXAMPLE 126**

25 3',5'-O-(tetraisopropylidisiloxane-1,3-diyl)- $\text{N}_2$ -[imidazol-4-yl(ethyl)]-2'-deoxyguanosine (126)

To a stirred solution of compound 125 (2.4 g, 6.65 mmol) in dry DMF (50 ml) and dry pyridine (20 ml) was added triethylamine (4.04 g, 40 mmol) followed by 1,3-dichloro-1,1,3,3-tetraisopropylidisiloxane (4.18 g, 13.3  
30 mmol) at room temperature. The reaction mixture was stirred at room temperature overnight and evaporated to dryness. The residue on purification silica gel chromatography using methylene chloride ---> methanol as the  
35 eluent gave 3.2 g (80%) of 126 as an amorphous solid.

## EXAMPLE 127

3',5'-O-(tetraisopropylidisiloxane-1,3-diyl)-6-O-diphenylcarbamoyl-N<sub>2</sub>-[1-diphenylcarbamoylimidazol-4-yl-(ethyl)]-2'-deoxyguanosine (127).

- 5 To a solution of compound 126 (12.34 g, 20 mmol) in dry pyridine (50 ml) and dry DMF (200 ml) was added N,N-diisopropylethylamine (7.74 g, 60 mmol) and diphenylcarbamoyl chloride (11.55 g, 50 mmol) at room temperature. The reaction mixture was stirred at room temperature for 12 hrs under an argon atmosphere. It was evaporated to dryness and the residue was dissolved in methylene chloride (400 ml). The organic extract was washed with water (100 ml) and brine (50 ml), dried and evaporated to dryness. The residue was purified by flash chromatography over silica gel using methylene chloride  
10 ---> acetone as the eluent. The pure fractions were pooled together and evaporated to dryness to give 15 g (75%) of 127 as a foam.

## 20 EXAMPLE 128

6-O-diphenylcarbamoyl-N<sub>2</sub>-[1-diphenylcarbamoylimidazol-4-yl-(ethyl)]-2'-deoxyguanosine (128).

- Compound 128 can be prepared from compound 127 by deblocking the "TIP" group selectively with tetrabutylammonium fluoride/pyridine/THF at room temperature.  
25

## EXAMPLE 129

5'-O-(4,4'-dimethoxytrityl)-6-O-diphenylcarbamoyl-N<sub>2</sub>-[1-diphenylcarbamoylimidazol-4-yl-(ethyl)]-2'-deoxyguanosine (129).  
30

Compound 129 can be prepared from compound 128 as per the procedure of example 107.

- 120 -

## EXAMPLE 130

3'-O-[(N,N-diisopropylamino)( $\beta$ -cyanoethoxy)prosphanyl]-  
5'-O-(4,4'-dimethoxytrityl)-6-O-diphenylcarbamoyl-N<sub>2</sub>-[1-  
diphenylcarbamoylimidazol-4-yl-(ethyl)]-2'-deoxyguanosine  
5 (130).

Compound 130 can be prepared from compound 129 as per the procedure of example 108.

## EXAMPLE 131

Procedure for the attachment of modified 5'-dimethoxytri-  
10 phenylmethyl ribonucleosides of the invention to the 5'-hydroxyl of nucleosides bound to CPG support.

The modified nucleosides that will reside in the terminal 3'- position of certain antisense oligonucleotides are protected as their 5'-DMT (the cytosine and adenine exocyclic amino groups are benzoylated and the guanine amino is isobutrylated) and treated with trifluoroacetic acid/ bromoacetic acid mixed anhydride in pyridine and dimethylaminopyridine at 50° C for five  
15 hours. The solution is evaporated under reduced pressure to a thin syrup which is dissolved in ethyl acetate and passed through a column of silica gel. The homogenous fractions were collected and evaporated to dryness. A solution of 10 ml of acetonitrile, 10 micromols of the 3'-O-bromomethyl-ester modified pyrimidine nucleoside,  
20 and one ml of pyridine/ dimethylaminopyridine (1:1) is syringed slowly (60 to 90 sec) through a one micromole column of CPG thymidine (Applied Biosystems, INC.) that had previously been treated with acid according to standard conditions to afford the free 5'-hydroxyl group.  
25 Other nucleoside bound CPG columns could be employed. The eluent is collected and syringed again through the column. This process is repeated three times. The CPG column is washed slowly with 10 ml of acetonitrile and then attached to an ABI 380B nucleic acid synthesizer.  
30 Oligonucleotide synthesis is now initiated. The standard conditions of concentrated ammonium hydroxide deprotection that cleaves the thymidine ester linkage from the  
35

- 121 -

CPG support also cleaves the 3',5' ester linkage connecting the pyrimidine modified nucleoside to the thymidine that was initially bound to the CPG nucleoside. In this manner, any modified nucleoside or generally any nucleoside with modifications in the heterocycle and/or sugar can be attached at the very 3'-end of an oligonucleotide sequence.

**EXAMPLE 132**

Procedure for the conversion of modified nucleoside-5'-DMT-3'-phosphoramidites into oligonucleotides.

The polyribonucleotide solid phase synthesis procedure of B. S. Sproat, et al., Nucleic Acids Research Vol. 17, pp. 3373-3386 (1989) is utilized to prepare the modified oligonucleotides.

**15 EXAMPLE 133**

Preparation of Modified Phosphorothioates Oligonucleotides.

Substituted 5'-DMT nucleoside 3'-phosphoramidites prepared as per the above described Examples are inserted into sequence-specific oligonucleotide phosphorothioates as described by S. Beaucage et al., Journal of American Chemical Society Vol. 112, pp. 1253-1255 (1990) and B. S. Sproat, et al., Nucleic Acids Research Vol. 17, pp. 3373-3386 (1989).

**25 EXAMPLE 134**

Preparation of Modified Phosphate Methylated Oligonucleotides.

The protection, tosyl chloride mediated methanolysis, and mild deprotection described by L. H. Koole et al., in the Journal of Organic Chemistry Vol. 54, pp. 1657-1664 (1989), is applied to substituted oligonucleotides to afford phosphate-methylated substituted oligonucleotides.

- 122 -

## EXAMPLE 135

The ability of modified oligonucleotides of the invention to hybridize to their complementary RNA or DNA sequences is determined by thermal melting analysis.

5 Natural antisense oligonucleotides or those containing modifications at specific locations were added to either the DNA or the RNA complement at stoichiometric concentrations. The helix to coil transition was monitored by measuring absorbance at 260 nm vs. temperature. Measure-

10 ments were made in 100mM Na<sup>+</sup>, 10 mM phosphate, pH 7 and 0.1 mM EDTA; oligonucleotide concentrations were 4  $\mu$ M. The melting temperature or T<sub>m</sub> is the temperature at which half the molecules are in the duplex state and half are single stranded. Reported T<sub>m</sub>'s are the temperature at

15 which dAbsorbance/dTemperature is maximum. Thermodynamic parameters of duplex formation were obtained from a non-linear least squares fit for the absorbance vs. temperature curves to a two state model with linear sloping baselines.

20 Each modification was incorporated into several oligonucleotide sequences ranging in length from 15 to 21 residues. The sequences studied were various 15 to 21 mer of target organism including papiloma virus, herpes virus, 5-LO, HIV and other testing sequences.

25 Oligonucleotides containing from one to five modifications were tested in addition to fully modified sequences. For each modified oligonucleotide, the T<sub>m</sub> was compared to that of the unmodified DNA analog. In addition comparisons were made to oligonucleotides having

30 their phosphodiester linkages modified via phosphorothioate substitution.

In this example the thermodynamic stability of certain of the novel modified oligonucleotides of the invention were determined by melting studies. Certain of these

35 studies as shown in Table 2 below reveal that up to five pendant groups specifically placed within oligonucleotide sequences of 15 to 21 bases in length generally resulted in only a modest effect on T<sub>m</sub>s compared with the unmodi-

- 123 -

fied parent oligonucleotides. The effects on  $T_m$ s were dependent on location in the sequence in that modifications placed near the 3'-end had less destabilizing effects than 5'-end placements.

5

Table 2

## THERMODYNAMIC STABILITY

Sequence studied A'CC GA'C G'AT C'AT GTC GTA' CGC  
(21 mer, positions 1, 5, 8, 11 & 18 modified)

	<u>Modification</u>	<u><math>T_m</math> (delta <math>T_m</math>, C)</u>
10	natural	67
	P=S (1-20)	57 (-10)
	2'-O-nonyl	51 (-16)
	2'-O-aminopropyl	61 (-6)
	2'-O-fluoro	67 (0)
15	2'-O-allyl	58 (-9)

Sequence studied CGA CTA TGC AAG TA'C  
(15 mer, position 14 modified)

	<u>Modification</u>	<u><math>T_m</math> (delta <math>T_m</math>, C)</u>
	natural	53
20	P=S (1-14)	46 (-7)
	2'-O-nonyl	54 (+1)
	2'-t-butyl dimethylsilyl	52 (-1)
	2'-O-aminopropyl	53 (0)
	2'-O-aminobutyl	53 (0)
25	RNA	46
	2'-O-allyl	45 (-1)
	2'-O-benzyl	45 (-1)

Sequence studied CGA CTA TGC AAA' A'A'C  
(15 mer, positions 12, 13, 14 modified)

	<u>Modification</u>	<u><math>T_m</math> (delta <math>T_m</math>, C)</u>
30	natural	52
	P=S (1-14)	45 (-17)
	2'-O-nonyl	49 (-3)
	2'-t-butyl dimethylsilyl	45 (-17)
35	2'-O-aminopropyl	54 (+2)
	2'-O-allyl	51 (-1)

## EXAMPLE 136

In this test specificity of hybridization was determined by comparing stability of a duplex containing  
40 an oligonucleotide modified at a single residue and its complement to the stability of a duplex containing the same singly modified oligonucleotide and a complementary

- 124 -

- sequence containing a not-complementary base opposite the modification. For example, specificity of an unmodified adenosine was determined by measuring the  $T_m$  of the sequence CTC GTA CCA TCC CCG TCC (X strand) hybridized to
- 5 the sequence GGA CCG GAA YGG TAC GAG (Y strand) where Y = A, C, G, T or none. The difference in  $T_m$  for the matched duplex (Y = T) compared to the duplex containing a single mismatch or bulge (Y = A, C, G or none) is a measure of the specificity of the unmodified A-T pair.
- 10 Similar experiments were performed with a modified adenosine replacing the A in position 9 of X strand. Comparison of the delta  $T_m$ 's (mismatch vs. match) for the modified sequence to those for the unmodified analog yields a measure of specificity of that modification.
- 15 In this test the specificity of the modified adenosines to Watson-Crick base pair with thymidine was determined by preparing as a Y stand, an 18 mer with a T (Watson-Crick complement), A, G, C (mismatch), or the deletion of the nucleotide at the 9 position (bulge loop)
- 20 The X complement strand contains a 2'-modified adenosine at the 9th position. The data is shown in Table 3. As per Table 3, specificity of the modified adenosines were retained, i.e. their  $T_m$ 's were greater when paired with T rather than the mismatch situations of A, G, C or bulge
- 25 loop.

- 125 -

Table 3

## SPECIFICITY OF MODIFICATION

	X STRAND	5'-CTC GTA CCA'	TTC CGG TCC		
	Y STRAND	3'-GAG CAU GGY'	AAG GCC AGG (18 mer, 9-position modified)		
5	Y = T (Watson-Crick COMPLEMENT)				
	Y = A, C OR G (MISMATCH)				
	Y = NONE (BULGE LOOP, "Bu")				
	Modification	T <sub>m</sub> (delta T <sub>m</sub> , C)			
10		A-T, A-A, A-G,	A-C, A-Bu		
	natural	63	-9,	-6,	-14, -12
	2'-O-nonyl	58	-3,	-3,	-6, -3
	2'-t-butyldimethylsilyl	58	-4,	-4,	-2, -4
15	2'-O-aminopropyl	60	-6,	-6,	-12, -7
	2'-fluoro	61	-10,	-7,	-14, -10
	2'-O-allyl	60	-8,	-6,	-13, -10
	2'-O-benzyl	59	-6,	-6,	-12, -7
	2'-O-methyl	60	-8,	-6,	-13, -10
20	2'-chloro	60	-4,	-4,	-13, -7
		G-C, G-A, G-G, G-T, G-Bu			
	natural	63	-7,	-6,	-4, -11
25	3-deazaguanosine	59	-2,	+1,	-1, -7

## EXAMPLE 137

In this example resistance to nuclease degradation is measured. Aliquots of 10% fetal calf serum in DMEM media were incubated at 37°C containing 50 μM oligonucleotide. At various times, 15 μl aliquots were removed and mixed with 15 μl 9 M urea in 1x TBE. Time points were frozen until analysis via 20% polyacrylamide / 7M urea electrophoresis. Gels were stained with "Stains All" (Sigma Chem Co., St. Louis, MO) and following destaining, scanned using a LKB Laser Densitometer. Scans were analyzed by integration of bank peaks and calculations of % degraded from the full length oligonucleotide (n) to the loss of two base residues (n-2). Data was plotted as % degraded vs time and compared to curves generated for phosphodiester (PO) and phosphorothioate (PS) oligonucleotides.



- 126 -

For this example the data is shown in Table 4. As per this data the modified oligonucleotides of the invention are at least as resistant as unmodified sequences and in certain instances more resistant than phosphorothioate modified sequences.

Table 4  
NUCLEASE RESISTANCE

Sequence studied CGA CTA TGC AAG TA'C  
(15 mer, position 14 modified)

	<u>Modification</u>	<u>T<sub>1/2</sub> (hr)</u>
10	natural	1.2
	P=S (1-14)	2.3
	2'-O-nonyl	5.8
	2'-t-butyldimethylsilyl	7
15	2'-O-aminopropyl	7

Sequence studied CGA CTA TGC AAA' A'A'C  
(15 mer, positions 12, 13, 14 modified)

	<u>Modification</u>	<u>T<sub>1/2</sub> (hr)</u>
20	natural	1.0
	P=S (1-14)	20
	2'-O-nonyl	> 64
	2'-t-butyldimethylsilyl	8.0
	2'-O-aminopropyl	17.7
	2'-O-allyl	10

25 Sequence studied CGA CAA TGC AAG' G'G'T  
(15 mer, positions 12, 13, 14 modified)

	<u>Modification</u>	<u>T<sub>1/2</sub> (hr)</u>
30	natural	2.5
	3-deaza deoxy G	1.0
	3-deaza-3-nonyl deoxy G	4.4
	3-deaza-3-allyl deoxy G	20+
	3-deaza-3-benzyl deoxy G	20

#### EXAMPLE 138

In general the delta T<sub>m</sub> between the modified oligonucleotide and its unmodified analog is proportional to the number of modified residues. In a further test delta T<sub>m</sub> per modification were noted. Values of delta T<sub>m</sub> per modifications were averaged over all modified sequences studied and are listed in Table 5. Data is shown as average delta T<sub>m</sub> per modification compared to wild type DNA for both DNA targets and RNA targets.

- 127 -

Further, specificity verse DNA targets is noted as an arbitrary score wherein 1 is specific and 4 is non-specific.

Table 5

5	Specificity	BIOPHYSICAL EVALUATION OF OLIGONUCLEOTIDES	
		Stability	Stability
		vs DNA	vs RNA
		vs DNA	vs DNA
	<u>Modification</u>	<u>Targets</u>	<u>Targets</u>
10	Natural DNA	0	1
	Natural RNA	+0.1	1
	phosphorothioate	-0.5	1+
	2'-Chloro-2'-deoxy A	-3.2	3
15	3-deaza-2'-deoxy G	-3.3	5
	2'-O-nonyl A	-2.5	4
	2'-t-butyl-dimethyl-silyl A	-4.5	3
20	2'-O-allyl A	-1.6	2
	2'-O-benzyl A	-2.7	2
	2'-O-propylamine	-0.9	3
	2'-O-butylamine	-1.2	-0.6

## EXAMPLE 139

25 In further tests nucleases resistance was compared relative to that of phosphorothioate and phosphodiester. Using the protocols of example 137, the degradation curve for phosphodiester oligonucleotides were determined and rated as 5. For phosphodiester oligonucleotides degradation was about 50% at about 1 hour and complete at about 30 4 hours. Phosphorothioate oligonucleotides were rated as 1. For phosphorothioate oligonucleotides degradation was about 20% at about four hours and about 50% at about 20 hours. Oligonucleotides containing modifications were 35 assigned values based upon visual comparison to the phosphodiester and phosphorothioate curves. These values are reported in Table 6.

- 128 -

Table 6

## RELATIVE RESISTANCE TO FETAL CALF SERUM NUCELASES

	<u>Modification</u>	<u>Position (#)</u>	<u>Relative Rate</u>
	Natural DNA	all	5
5	phosphorothioate	all	1
	phosphodiester	all	5
	2'-Chloro-2'-deoxy A		5
	2'-O-nonyl A	3' (1)	3
10		3' (3)	1
	2'-t-butyldimethylsilyl A	3' (3)	1
	2'-O-allyl A	3' (1)	3
	2'-O-benzyl A	3' (1)	3
15	2'-O-propylamine	3' (1)	2
	2'-O-Fluoro-2'-deoxy A	3' (3)	5
		internal	5
		all	5
20	2'-O-methyl A	all	1

## EXAMPLE 140

Utilizing the protocols of examples 135 and 138 oligonucleotides of the invention having novel nucleotides as prepared per certain of the above examples were tested for their DNA-DNA duplex stability or DNA-RNA duplex stability against DNA or RNA sequences. Hybridization analysis based upon the  $T_m$ , the  $\Delta T_m$ , and the  $\Delta T_m$  per modification were determined. The results of these tests are shown in Table 6 below.

The test sequences represent various sequences chosen to demonstrate specific activity in known pathogens or test sequences. The JM-12 is a synthetic sequence that is self complementary. Since it is self complementary each modification is expressed twice, once for each occurrence in the complementary sequences. The HSV-21 is a 21 mer sequence from herpes virus. The G capped PAP is sequence from papiloma virus having a C nucleotide at the 3' end that is the linking nucleotide during synthesis. This is followed by three capping G nucleotides. This capping is specifically added to enhance the nuclease resistance of the sequence. The XY sequence is particularly useful for the determination of steric hindrance of

- 129 -

the Watson-Crick binding between complementary strands. The 5L0 is a 16 mer from the mRNA of human 5-lipoxygenase.

For these tests the "2 Chloro dA" nucleotide was prepared as per the synthetic chemical sequence terminating with example 124 above and the "2 propyl imidazolyl dG" nucleotide as per the synthetic chemical sequence terminating with Example 107 above.

Table 7

10

**DUPLEX STABILITY****DNA-RNA DUPLEX STABILITY**

Complementary strand - YX: GGA CCG GAA GGT ACG AG

Test nucleotide - 2 Chloro dA

15	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
	natural	58.8	-	-
	3 8 9 13 16	35.0	-24	-5.2

Complementary strand- HSV-21: ACC GAG GAT CAT GTC GTA CGC

Test nucleotide - 2 Chloro dA

20	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
	natural	64.3	-	-
	1 5 8 11 18	54.5	-9.8	-2.0

**DNA-RNA DUPLEX STABILITY**

25 Complementary strand - HSV-21 GCC GAG GTC CAT GTC GTA CGC

Test nucleotide - 2 propyl imidazolyl dG

30	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
	natural	69.1	-	-
	13	69.0	0.4	0.4
	4 13 20	70.3	+1.3	+0.2
	1 4 6 7 13 16 20	70.4	+1.3	+0.2

- 130 -

Complementary strand- HSV-21: ACC GAG GTC CAT GTC GTA CGC

Test nucleotide - 2'-fluor-2'-deoxy A

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
5	DNA	67.5	-	-
	(terminal A C mismatch)			
	1 5 11 18	70.0	+2.5	+0.6
	(terminal A C mismatch)			

Complementary strand- HSV-21: ACC GAG GAT CAT GTC GTA CGC

10 Test nucleotide - 2'-fluoro-2'-deoxy A

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
	DNA	63.1	-	-
	1 5 8 11 18	67.7	+4.5	+0.9

15 Complementary strand - PAP: CGA CTA TGC AAG TAC

Test nucleotide - 2'-fluor-2'-deoxy A

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
	natural	45.1	-	-
20	3 6 10 11 14	53.0	+7.9	+1.6
	3 5 6 7 10 11 13 14	58.9	+13.8	+1.7
	1 3 4 5 6 7 9 10 13 14	65.2	+20.1	+1.8

Complementary strand - PAP: CGA CTA TGC AAG TAC

Test nucleotide - 2'-fluor-2'-deoxy A

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
25	all PS	33.9	-	-
	all PS with 2'-fluor in			
	1 3 4 5 6 7 9 10 11 13 14	60.9	+27.0	+2.5

- 131 -

Complementary strand - XY: GGA CCG GAA GGT ACG AG

Test nucleotide - 2'-fluor-2'-deoxy A

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
5	natural	56.9	-	-
	3 8 9 13 17	65.9	+9.0	+1.8

Complementary strand - XY: CTC GTA CCT TCC GGTC C

Test nucleotide - 2'-fluor-2'-deoxy A

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
10	natural	65.0	-	-
	2 5 9 10 15	68.5	+3.5	+0.7
	1 2 3 5 6 7 8 9 10			
	12 15 16	75.5	+10.4	+0.8

15 Complementary strand - 5LO: TCC AGG TGT CCG CAT C

Test nucleotide - 2'-fluor-2'-deoxy A

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
	natural	62.0	-	-
20	1 7 9 15	64.7	+2.8	+0.7

**DNA-DNA DUPLEX STABILITY**

Complementary strand- HSV-21: GCC GAG GTC CAT GTC GTA CGC

Test nucleotide - 2 propyl imidazol dG

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
25	natural	69.7	-	-
	13	72.3	2.6	2.6
	4 13 20	75.8	6.1	2.0
	1 4 6 7 13 16 20	83.3	+13.6	+1.9

- 132 -

Complementary strand - JM-12: GCC TGA TCA GGC

Test nucleotide - 2 propyl imidazol dG

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
5	natural	50.8	-	-
	5	63.4	+12.6	+6.3

Complementary strand - G capped PAP: CGA CTA TGC AAG GGC

Test nucleotide - 2 propyl imidazol dG

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
10	natural	56.5	-	-
	12 13 14	64.9	+8.4	+2.8

Complementary strand - XY: CTC GTA CCA TTC CGG TCC

Test nucleotide - 2'-fluor-2'-deoxy A

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
15	natural	62.6	-	-
	9	60.6	-2.0	-2.0

Complementary strand - XY: GGA CCG GAA GGT ACG AG

20 Test nucleotide - 2'-fluor-2'-deoxy A

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
	natural	60.6	-	-
	3 8 9 13 16	61.0	+0.4	+0.1

25 Complementary strand - XY: CTC GTA CCT TTC GGTC C

Test nucleotide - 2'-fluor-2'-deoxy A

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
30	natural	60.6	-	-
	2 5 9 10 15	55.9	-3.9	-0.8

- 133 -

Complementary strand - XY: CTC GTA CCT TTC CGG TCC

Test nucleotide - 2'-fluor-2'-deoxy A

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
5	natural	61.7	-	-
	9	60.7	-1.0	-1.0

Complementary strand- HSV-21: ACC GAG GTC CAT GTC GTA CGC

Test nucleotide - 2'-fluor-2'-deoxy A

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
10	natural	68.9	-	-
	1 5 11 18	67.0	-1.8	-0.5

Complementary strand- HSV-21: ACC GAG GAT CAT GTC GTA CGC

Test nucleotide - 2'-fluor-2'-deoxy A

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
15	natural	66.1	-	-
	1 5 8 11 18	66.4	+0.3	+0.1

Complementary strand - PAP: CGA CTA TGC AAG TAC

20 Test nucleotide - 2'-fluor-2'-deoxy A

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
	natural	53.2	-	-
	3 6 10 11 14	53.0	-0.2	0.0

25 Complementary strand - PAP: CGA CTA TGC AAG TAC

Test nucleotide - 2'-fluor-2'-deoxy A

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
	natural	52.3	-	-
30	5 7 13	52.1	-0.2	-0.1
	3 5 6 7 10 11 13 14	55.4	+3.2	+0.5



- 134 -

Complementary strand - A capped PAP: CGA CTA TGC AAG TAC

Test nucleotide - 2'-fluor-2'-deoxy A

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
5	natural	50.4	-	-
	12 13 14	43.6	-6.8	-2.3

Complementary strand - A capped PAP: CGA CTA TGC AAG TAC

Test nucleotide - 2'-fluor-2'-deoxy A

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
10	all PS	43.2	-	-
	all Ps with 2'-fluoro A			
	in 12 13 14	47.3	+4.1	+1.4

Complementary strand- PAP-mod: CGA CTA TGC AAG TAC AAA T

15 Test nucleotide - 2'-fluor-2'-deoxy A

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
	natural	56.5	-	-
	3 6 10 11 14 16 17 18	56.0	-0.5	-0.1

20 Complementary strand - 5LO: TCC AGG TGT CCG CAT C

Test nucleotide - 2'-fluor-2'-deoxy A

	<u>positions of modification</u>	<u>T<sub>m</sub></u>	<u>delta T<sub>m</sub></u>	<u>delta T<sub>m</sub>/mod</u>
	natural	62.5	-	-
25	1 7 9 15	61.2	-1.3	-0.3

**EXAMPLE 141**

Compositions for modulating the activity of messenger RNA coding for production of the protein 5-lipoxygenase are constructed as per this example and others that follow. The sequence CGUCCAGUGACUUC appears in the 5-lipoxygenase mRNA beginning at nucleotide 1065, defined by counting the initiation of translation "A" as zero. Other regions may also be selected which are likely to

- 135 -

lead to effective antisense agents. For the above described region, a target antisense oligonucleotide (GCAAGGTCACCTGAAG) directed to the 5-lipoxygenase mRNA may be constructed by standard solid state techniques leading to an effective composition in accordance with this invention. The 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing specific adenines replaced by 2-imidazolylethyl amino-2-deoxyadenine can be prepared as per examples 109 to 114 from 2-chloro-2'-deoxyadenosine (5.72 g, 20 mmol), Journal of the American Chemical Society Vol. 106, p. 6379 (1984). Appropriate amounts of this material are dissolved in dry THF and employed in the automated, solid phase synthesis of the 5LO sequence 5'- GAA GTC ACT GGA ACG -3'.

Modified sequences synthesized will be:

- GaA GTC ACT GGA ACG
  - GAA GTC ACT GGA aCG
  - GaA GTC ACT GGA aCG
  - GAA GTC aCT GGA ACG
  - GAA GTC ACT GGa ACG
  - GaA GTC aCT GGA aCG
  - Gaa GTC ACT GGA ACG
  - GAA GTC ACT GGa aCG and
  - Gaa GTC aCT GGA aCG
- Where a = 2-(imidazol-4-yl-ethylamino)adenine

#### EXAMPLE 142

Synthesis of the 5LO - sequence 5'- GAA GTC ACT GGA ACG -3' containing N2-imidazolylethylguanine (2-imidazolylamino hypoxanthine) at specific guanine positions is performed as per examples 101 to 108 substituting 1-(2-aminoethyl)-imidazol for 1-(3-aminopropyl)-imidazol. Appropriate amounts of this material are dissolved in dry THF and employed in the automated, synthesis of the 5LO 5'-sequence 5'- GAA GTC ACT GGA ACG -3'.

Modified sequences synthesized will be:

- gAA GTC ACT GGA ACG
- GAA GTC ACT GGA ACg

- 136 -

GAA gTC ACT GGA ACG

GAA GTC ACT gGA ACG

GAA GTC ACT GgA ACG

gAA GTC ACT GGA ACg

5 gAA gTC ACT GGA ACg and

gAA gTC ACT ggA ACg

Where g = N2-(imidazolylethyl)guanine ((2-imidazol-4-ylethylamino) hypoxanthine)

**EXAMPLE 143**

10 Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 2'-deoxy-2'-(imidazol-4-yl-propyl)adenine at specific positions is performed. A solution of 3',5'-O-(tetraisopropyl disiloxane-1,3-diyl)-9- $\beta$ -D-arabinofuranosyladenine (10.2 g, 20 mmol) in 200 ml of  
15 dry pyridine is cooled in an ice bath and treated with trimethylchlorosilane (6 ml). After 0.5 hr, benzoyl chloride (6 ml) is added and the ice bath is removed. The reaction is chilled after two hr at ambient temperature and treated with 20 ml of cold water followed after 15  
20 min with 20 ml of concentrated ammonia hydroxide. The reaction is evaporated under reduced pressure to an oil which is purified by silica gel chromatography to afford nine g of the 3',5',N6 blocked nucleoside. This material is dissolved in dry acetonitrile (200 ml) and dry DMF (50  
25 ml), chilled in an ice bath, and treated with sodium hydride (400 mg) followed 30 min later with trifluoromethylsulfonic acid anhydride (5 g) in dry acetonitrile (50 ml) added dropwise. The reaction is evaporated under reduced pressure. The residue is dissolved in ethyl  
30 acetate, washed with water, and dried with MgSO<sub>4</sub>. Removal of the solvent under reduced pressure provides the 2'-trifluoro sulfonate as a hard foam. A sample of this material (4 g, 5.5 mmol) is dissolved in dry ethyl acetate (100 ml) and diisopropylethylamine (800 mg),  
35 cooled in an ice bath and treated with 1-(4-imidazolyl)-3-hydroxy-n-propane (750 mg). After two hr at ambient temperature, the reaction is washed with 50 ml of cold

- 137 -

- water and dried with  $\text{MgSO}_4$ . Removal of the solvent under reduced pressure provides the 2'-(imidazolylpropoxy) adenosine blocked in the 3',5' and N6-positions. This material is dissolved in 50 ml of dry THF and treated with tetrabutylammonium fluoride (10 ml of 1 M in THF) at room temperature for two hr. The solvent is removed under reduced pressure and the residue, absorbed on silica gel, is chromatographed over silica gel, eluting with chloroform/methanol mixtures. The 5'-DMT and 3'-cyanoethoxyphosphoramidite of this material is prepared as per examples 107 and 108 above and appropriate quantities of this material is dissolved in dry acetonitrile and utilized in the automated, solid phase synthesis of the 5LO sequence 5'- GAA GTC ACT GGA ACG.
- Modified sequences synthesized will be:
- GAA GTC ACT GGA aCG
  - GaA GTC ACT GGA aCG
  - GaA GTC aCT GGA aCG
  - GAA GTC aCT GGA ACG
  - GAA GTC ACT GGA aCG
  - GaA GTC ACT GGA ACG
  - GAA GTC ACT GGA ACG
  - GAA GTC ACT GGA ACG and
  - Gaa GTC aCT GGA aCG
- Where a = 2'-O-(imidazol-4-yl-n-propyl)adenine

**EXAMPLE 144**

- Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG - 3' containing 2'-(imidazol-4-yl-propyl)thymidine at specific positions is described. A solution of 3',5'-O-(tetra-isopropylidisiloxane -1,3-diyl)-9- $\beta$ -D-arabinofuranosyl-thymine (9.7 g, 20 mmol) is dissolved in dry acetonitrile (200 ml) and dry DMF (50 ml), chilled in an ice bath, and treated with sodium hydride (400 mg) followed 30 min later with trifluoromethylsulfonic acid anhydride (5 g) in dry acetonitrile (50 ml) added dropwise. The reaction is evaporated under reduced pressure. The residue is dissolved in ethyl acetate, washed with water, and dried

- 138 -

with  $\text{MgSO}_4$ . Removal of the solvent under reduced pressure provides the 2'-trifluoro sulfonate as a hard foam. A sample of this material (4 g, 5.5 mmol) is dissolved in dry ethyl acetate (100 ml) and diisopropyl ethylamine (800 mg), cooled in an ice bath and treated with 1-(imidazol-4-yl)-3-hydroxy-n-propane (750 mg). After two hr at ambient temperature, the reaction is washed with 50 ml of cold water and dried with  $\text{MgSO}_4$ . The material is further reacted as per examples 107 and 108 above to provide the 5'-DMT-3'cyanoethoxyphosphoramidite. Appropriate quantities of this material is dissolved in dry acetonitrile and utilized in the automated, solid phase synthesis of the 5LO sequence 5'- GAA GTC ACT GGA ACG.

Modified sequences synthesized will be:

GAA GTC ACT GGA ACG

GAA GtC ACT GGA ACG and

GAA GtC ACT GGA ACG

Where t = 2'-O-(imidazol-4-yl-n-propyl)thymine

**EXAMPLE 145**

Synthesis of the 5LO - sequence 5'- GAA GTC ACT GGA ACG containing 3-deaza-3-(imidazol-4-yl-propyl)guanine at specific positions is described. A solution of 5.2 g, 10 mmol, of 5-(cyanomethyl)-1-(2-deoxy-3,5-di-O-p-toluoyl- $\beta$ -D-erythro-pentofuranosyl)imidazole-4-carboxylate, Journal of Medicinal Chemistry, Vol. 27, p. 1389 (1984), is dissolved in dry THF (100 ml) and treated with sodium hydride (840 mg) at room temperature. After evolution of hydrogen ceases, 4-(2-bromoethyl)imidazole hydrobromide (2.6g), Rec. Trav. Chim., Vol. 89, p. 1189 (1970), is added and the solution stirred at ambient temperature for five hr. The reaction is evaporated to dryness. The residue is dissolved in ethyl acetate, washed with water and dried with  $\text{MgSO}_4$ . The residue obtained after filtration and solvent removal is dissolved in liquid ammonia and kept in a pressure vessel at ambient temperature for 24 hr and then heated to 100°C for 1 hr. The

- 139 -

ammonia is vented and the residue coevaporated with methanol two times. This material is dissolved in methanol, absorbed on silica gel and placed on a silica gel column. Elution with chloroform/methanol provides 2.0 g of 1-(2-deoxy- $\beta$ -D-erythro-pentofuranosyl)-5-(1-cyano-3-[imidazol-4-yl]prop-1-yl)-4-carboxamide. This material is converted to the 5'-DMT-3'-cyanoethoxy isopropylphosphoramidite as per examples 107 and 108 above, affording a hard foam. Appropriate amounts of this material is dissolved in dry acetonitrile and utilized in the automated, solid phase synthesis of the 5'LO sequence 5'-GAA GTC ACT GGA ACG.

Modified sequences synthesized will be:

gAA GTC ACT GGA ACg  
15 gAA GTC ACT gGA ACg  
gAA gTC ACT GGA ACg  
GAA GTC ACT GGA ACg  
GAA GTC ACT ggA ACG  
GAA gTC ACT GGA ACG  
20 gAA GTC ACT GGA ACG  
GAA GTC ACT GgA ACG  
GAA gTC ACT GGA ACG and  
gAA gTC ACT gGA ACg

Where g = 3-deaza-3-(4-imidazolylethyl)guanine

#### 25 EXAMPLE 146

Synthesis of the 5'LO sequence 5'- GAA GTC ACT GGA ACG containing 7-(imidazol-4-yl-methyl)-7,8-dihydro-imidazo [4,5-c]pyrrolo[3,2-e]pyridin-4(5H)-one at specific sites is performed. A solution of methyl 1-(3,5-di-O-toluoyl- $\beta$ -D-erythro-pentofuranosyl)-5-cyanomethyl-imidazole-4-carboxylate (10.34 g, 20 mmol), Journal of Medicinal Chemistry, Vol. 27, p. 1389 (1984), and dry acetonitrile is treated with sodium hydride (500 mg). After stirring at ambient for 30 min, the solution is treated with 1-bromo-3-(1-t-Boc-imidazol-4-yl)-2-oxo ethylene ketal (7.24 g) and stirring is continued for 16 hr. The reaction is evaporated under reduced pressure and the

- 140 -

residue is distributed between ethyl acetate and water. The solvent is dried and removed under reduced pressure. The residue is placed in a Parr pressure vessel with liquid ammonia. The reaction is carried out for eight hr  
5 at 80 C cooled and vented. The residue is co-evaporated with methanol several times, dissolved in hot aqueous ethanol and adjusted to pH 4 with hydrochloric acid. After one hour the solution is selectively hydrogenated in the presence of Raney nickel. After one equivalent of  
10 hydrogen is taken up, the reaction is filtered and adjusted to pH 7. The solution is evaporated to dryness under reduced pressure. This material is absorbed on silica gel with the aid of methanol and chromatographed over silica gel, eluting with methanol/chloroform to  
15 afford 3.2 g of 3-(2-deoxy- $\beta$ -D-erythro-pentofuranosyl)-7-(imidazol-4-yl-methyl)-7,8-dihydro-imidazo [4,5-c]-pyrrolo[3,2-e]pyridin-4(5H)-one. This material is converted to the 5'-DMT-3'-cyanoethoxyphosphoramidite-5,6-diisobutryl according to the procedures of examples  
20 107 and 108 above and inserted into the 5LO-sequence 5'-GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

Modified sequences synthesized will be:

gAA GTC ACT GGA ACG  
25 gAA GTC ACT gGA ACG  
gAA gTC ACT GGA ACG  
GAA GTC ACT GGA ACG  
GAA GTC ACT ggA ACG  
GAA gTC ACT GGA ACG  
30 gAA GTC ACT GGA ACG  
GAA GTC ACT GgA ACG  
GAA gTC ACT GGA ACG and  
gAA gTC ACT ggA ACG

Where g will be 7-(imidazol-4-yl-methyl)-7,8-dihydro-  
35 imidazo [4,5-c]pyrrolo [3,2-e] pyridin-4(5H)-one

- 141 -

**EXAMPLE 147**

Synthesis of the 5LO sequence 5'- GAA GTC ACT GGA ACG containing 8-(imidazol-4-yl-methyl)-7,8-dihydroimidazo [4,5-c] pyrrolo [3,2-e]pyridin-4(5H)-one within the oligonucleotide is performed. A solution of methyl 1-(3,5-di-O-toluoyl- $\beta$ -D- erythro-pentofuranosyl)-5-cyanomethylimidazole-4-carboxylate (10.34 g, 20 mmol) in dry acetonitrile is treated with sodium hydride (500 mg). After stirring at ambient temperature for 30 min, the solution is treated with 2-bromo-3-(1-t-Boc-imidazol-4-yl)-propanone ethylene ketal (7.24 g) and stirring is continued for 16 hr. The reaction is worked up and further reacted as described above to provide 1-(2-deoxy- $\beta$ -D-erythropento-furanosyl)-8-(imidazol-4-yl-methyl)-7,8-dihydroimidazo-[4,5-c]pyrrolo[3,2-e]pyridin-4(5H)-one. This material is converted to the 5'-DMT-3'-cyanoethoxy-phosphoramidite -5,6-di isobutryl according to examples 107 and 108 above and inserted into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

Modified sequences synthesized will be:

gAA GTC ACT GGA ACg  
 gAA GTC ACT gGA ACg  
 gAA gTC ACT GGA ACg  
 25 GAA GTC ACT GGA ACg  
 GAA GTC ACT ggA ACG  
 GAA gTC ACT GGA ACG  
 gAA GTC ACT GGA ACG  
 GAA GTC ACT GgA ACG  
 30 GAA gTC ACT GGA ACG and  
 gAA gTC ACT ggA ACG

Where g = 8-(imidazol-4-yl-methyl)-7,8-dihydro-imidazo [4,5-c]pyrrolo[3,2-e]pyridin-4(5H)-one

**EXAMPLE 148**

35 Synthesis of the 5LO sequence 5'- GAA GTC ACT GGA ACG containing 5-(imidazol-4-yl-ethoxymethyl) imidazole-2(H)one 4-carboxamide within the sequence is performed.



- 142 -

A mixture of ethyl 5-bromomethyl -2-oxo-imidazole 4-carboxylate ( 5.4 g, 20 mmol), imidazol-4-yl- 2-ethanol (2.3 g), dry acetonitrile (100 ml) is treated with sodium hydride (750 mg) and stirred for 5 hr at ambient temperature. The solution is adjusted to approximately pH 7 with acetic acid and evaporated to dryness. Purification by chromatography provides the desired material in 30% yield. This material dissolved in dry pyridine (100 ml) and N-ethyl-diisopropyl amine (8 ml) is treated with diphenylaminocarbonyl chloride (5 ml). Stirring is continued for 5 hr. The solution is evaporated to dryness under reduced pressure and the residue is distributed between ethyl acetate and water. The organic layer is dried with magnesium sulfate and evaporated to dryness. Purification by silica gel chromatography afforded the bis(diphenylcarbonyl) protected imidazole (3.5 g, 5 mmol). This is dissolved in dry acetonitrile (50 ml) and treated with sodium hydride (130 mg) after stirring at ambient temperature for 30 min, 3,5-di-O-(p-toluoyl)- $\alpha$ -D-erythro- pentofuransyl chloride is added. The mixture is stirred for 8 hr and filtered. The filtrate is evaporated to dryness and the filtered residue washed with water. The remaining residue is dissolved in ethyl acetate and combined with the filtrate residue. This material is purified by chromatography over silica gel to yield 4 g of blocked nucleoside. This is dissolved in methanol saturated at 20°C with ammonia. The solution is allowed to stir at ambient temperature overnight. Evaporation under reduced pressure and subsequent silica gel chromatography provides the sugar deprotected nucleoside. This material is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'- $\beta$ -cyanoethylphosphoramidite which is inserted into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

Modified sequences synthesized will be:

gAA GTC ACT GGA ACG  
gAA GTC ACT gGA ACG  
gAA gTC ACT GGA ACG

- 143 -

GAA GTC ACT GGA ACg  
GAA GTC ACT ggA ACG  
GAA gTC ACT GGA ACG  
gAA GTC ACT GGA ACG  
5 GAA GTC ACT GgA ACG  
GAA gTC ACT GGA ACG and  
gAA gTC ACT ggA ACg  
Where g = 5-(imidazol-4-yl-ethoxymethyl)-imidazole-  
2(H)-one 4-carboxamide.

10 **EXAMPLE 149**

Synthesis of the 5LO sequence 5'- GAA GTC ACT GGA ACG  
containing 4-amino-5-(imidazol-4-yl-ethoxymethyl)-  
imidazole-2 (3H)one at specific sites is described. A  
mixture of 2,4-dibromo -5-nitroimidazole (5.4 g, 20  
15 mmol), imidazol-4-yl-2-ethanol (2.3 g), dry acetonitrile  
(100 ml) is treated with sodium hydride (750 mg) and  
stirred for 5 hr at ambient temperature. The solution is  
adjusted to approximately pH 7 with acetic acid and  
evaporated to dryness. Purification by chromatography  
20 provides the desired material in 30% yield. This  
material dissolved in dry pyridine (100 ml) and N-ethyl-  
diisopropyl amine (8 ml) is treated with diphenylamino-  
carbonyl chloride (5 ml). Stirring is continued for 5  
hr. The solution is evaporated to dryness under reduced  
25 pressure and the residue distributed between ethyl  
acetate and water. The organic layer is dried with  
magnesium sulfate and evaporated to dryness. Purification  
by silica gel chromatography afforded the bis(diphenyl-  
carbamoyl) protected imidazole (3.5 g, 5 mmol). This is  
30 dissolved in dry acetonitrile (50 ml) and treated with  
sodium hydride (130 mg) after stirring at ambient  
temperature for 30 min, 3,5-di-O-(p-toluoyl)- $\alpha$ -D-erythro-  
pentofuransyl chloride is added. The mixture is stirred  
for 8 hr and filtered. The filtrate is evaporated to  
35 dryness. The filtered residue is washed with water and  
the remaining residue is dissolved in ethyl acetate and  
combined with the filtrate residue. This material is

- 144 -

purified by chromatography over silica gel to yield 4 g of blocked nucleoside. This is dissolved in methanol saturated at 20 °C with ammonia. The solution is allowed to stir at ambient temperature overnight. Evaporation under reduced pressure and subsequent silica gel chromatography provides the sugar deprotected nucleoside. This material is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'- $\beta$ -cyanoethylphosphoramidite which is inserted into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis. Modified sequences synthesized will be:

gAA GTC ACT GGA ACg  
gAA GTC ACT gGA ACg  
gAA gTC ACT GGA ACg  
15 GAA GTC ACT GGA ACg  
GAA GTC ACT ggA ACG  
GAA gTC ACT GGA ACG  
gAA GTC ACT GGA ACG  
GAA GTC ACT GgA ACG  
20 GAA gTC ACT GGA ACG and  
gAA gTC ACT ggA ACg

where g is 4-amino-5-(imidazol-4-yl-ethoxymethyl)-imidazole- 2(H)-one.

#### EXAMPLE 150

25 Synthesis of the 5LO sequence 5'- GAA GTC ACT GGA ACG containing 4-amino-5-(imidazol-4-yl-ethoxymethyl)-imidazole at specific sites is described. A mixture of 2,4-dibromo-5-nitro imidazole ( 5.4 g, 20 mmol), 2-(1-benzyl-imidazol-4-yl)-ethanol, dry acetonitrile (100 ml) is  
30 treated with sodium hydride (750 mg) and stirred for 5 hr at ambient temperature. The solution is adjusted to approximately pH 7 with acetic acid and evaporated to dryness. Purification by chromatography provides the desired material in 30% yield. This is dissolved in dry  
35 acetonitrile (50 ml) and treated with sodium hydride (130 mg) after stirring at ambient temperature for 30 min, 3,5-di-O-(p-toluoyl)- $\alpha$ - D-erythro-pentofuransyl chloride

- 145 -

is added. The mixture is stirred for 8 hr and filtered. The filtrate is evaporated to dryness. The filtered residue is washed with water and the remaining residue is dissolved in ethyl acetate and combined with the filtrate residue. This material is purified by chromatography over silica gel to yield 4 g of blocked nucleoside. This is dissolved in methanol saturated at 20°C with ammonia. The solution is allowed to stir at ambient temperature overnight. Evaporation under reduced pressure and subsequent silica gel chromatography provides the sugar deprotected nucleoside. Catalytic reduction of this material with hydrogen provides 2-amino-1-(2-deoxy-β-D-erythropentofuranosyl) -5-(imidazol-4-yl-ethoxymethyl)imidazole. Transient benzylation provides the 4-benzoylamino-5-(1-benzoylimidazol-4-yl-ethoxymethyl) derivative. This material is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'-b-cyanoethylphosphoramidite which is inserted into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

Modified sequences synthesized will be:

gAA GTC ACT GGA ACg  
 gAA GTC ACT gGA ACg  
 gAA gTC ACT GGA ACg  
 GAA GTC ACT GGA ACg  
 GAA GTC ACT ggA ACG  
 GAA gTC ACT GGA ACG  
 gAA GTC ACT GGA ACG  
 GAA GTC ACT GgA ACG  
 GAA gTC ACT GGA ACG and  
 gAA gTC ACT ggA ACg

where g is 4-amino-5-(imidazol-4-yl-ethoxymethyl)-imidazole.

**EXAMPLE 151**

Synthesis of the 5LO sequence 5'- GAA GTC ACT GGA ACG containing 5-(imidazol-4-yl-ethoxymethyl)imidazole-2(H)one at specific sites will be carried out. A mixture of 4-bromoimidazole-2(H)one (20 mmol), and diphenyl-

- 146 -

aminocarbonyl chloride is treated with sodium hydride (750 mg) and stirred for 5 hr at ambient temperature. The solution is adjusted to approximately pH 7 with acetic acid and evaporated to dryness. Purification by chromatography provides the desired material in 30% yield. This is dissolved in dry acetonitrile (50 ml) and treated with sodium hydride (130 mg). After stirring at ambient temperature for 30 min, 3,5-di-O-(p-toluoyl)- $\alpha$ -D-erythro-pentofuransyl chloride is added. The mixture is stirred for 8 hr and filtered. The filtrate is then evaporated to dryness. The filtered residue is washed with water and the remaining residue dissolved in ethyl acetate and combined with the filtrate residue. This material is purified by chromatography over silica gel to yield 4 g of blocked nucleoside. This is dissolved in methanol saturated at 20°C with ammonia. The solution is allowed to stir at ambient temperature overnight. Evaporation under reduced pressure and subsequent silica gel chromatography provides the sugar deprotected nucleoside. This material is then hydroxymethylated in the 5-position. Hydrogenolysis of this material with hydrogen provides 1-(2-deoxy - $\beta$ -D-erythropentofuranosyl)-5-hydroxymethylimidazole. Transient benzylation provides the 4-benzoylamino-5-(1-benzoylimidazo -4-yl-ethoxymethyl) derivative. This material is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'- $\beta$ -cyanoethyl phosphoramidite which is inserted into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

Modified sequences synthesized will be:

- gAA GTC ACT GGA ACg
- gAA GTC ACT gGA ACg
- gAA gTC ACT GGA ACg
- GAA GTC ACT GGA ACg
- GAA GTC ACT ggA ACG
- GAA gTC ACT GGA ACG
- gAA GTC ACT GGg ACG
- GAA GTC ACT GgA ACG

- 147 -

GAA gTC ACT GGA ACG and

gAA gTC ACT gga ACg

where g = 4-amino-5-(imidazol-4-yl-ethoxymethyl)-imidazole.

## 5 EXAMPLE 152

Synthesis of the 5LO sequence 5'- GAA GTC ACT GGA ACG containing 2-amino-5-(imidazol-4-yl-ethoxymethyl)-imidazole at specific sites is described. A mixture of 4-bromo-2-nitroimidazole (20 mmol), acetonitrile (100 ml), and sodium hydride is stirred at 50°C for 30 min, cooled and treated with 3,5-di-O-p-methyl benzoyl- $\alpha$ -D-erythropentofuranosyl chloride. The solution is stirred at ambient temperature for eight hours and filtered to remove sodium chloride. The filtrate is evaporated to dryness and purified by silica gel chromatography. This material is chloromethylated in the 5-position. Subsequent nucleophilic displacement with 2-(imidazol-4-yl) ethanol under basic catalysis provides the 5-(imidazol-4-yl-ethoxymethyl) derivative. Hydrogenolysis of the bromo atom and reduction of the group is accomplished by hydrogen and palladium on charcoal conditions. Transient benzoylation is used to protect the 5-position imidazole derivative. This material is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'-b-cyano-ethylphosphoramidite which is inserted into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

Modified sequences synthesized will be:

Gaa GTC ACT GGA ACG

30 Gaa GTC ACT GGA aCG

Gaa GTC ACT GGA ACG

GAA GTC ACT GGa aCG

GAA GTC aCT GGA ACG

GAA GTC ACT GGa ACG

35 GAA GTC ACT GGA ACG

Gaa GTC aCT GGA ACG and

Gaa GTC aCT GGa aCG

- 148 -

where a is 2-amino-5-(imidazol-4-yl-ethoxymethyl) imidazole.

**EXAMPLE 153**

Synthesis of the 5LO sequence 5'- GAA GTC ACT GGA ACG  
5 containing 1-(imidazol-4-yl-ethyl)thymine-6-yl at specific  
sites is described. A mixture of 3,5-di-O-p-methyl-  
benzoyl- $\alpha$ -D- erythropentofuranosyl chloride in aceto-  
nitrile is cooled to 0°C and treated with 6-lithio-1-(1-  
[benzoyl]-imidazol-4-yl)ethyl- 3-benzylthymine. The  
10 solution is allowed to warm to room temperature and after  
three hours at ambient temperature the mixture is eva-  
porated to dryness under reduced pressure. Silica gel  
chromatography with chloroform and methanol) afforded and  
material which is then debenzoylated with methanolic  
15 ammonia at room temperature overnight. Recrystallization  
from ethanol provides 6'-(2-deoxy- $\beta$ -D-erythropentofura-  
nosyl)-1-(imidazol- 4-yl-ethyl)thymine. This material is  
converted by procedures of examples 107 and 108 above  
into its 5'-DMT-3'- $\beta$ -cyanoethylisopropyl phosphoramidite  
20 which is inserted into the 5LO-sequence 5'- GAA GTC ACT  
GGA ACG via automated, solid phase DNA synthesis.  
Modified sequences synthesized will be:

GAA GtC Act GGA ACG

GAA GTC Act GGA ACG and

25 GAA GtC Act GGA ACG

where t = 1-(imidazol-4-yl-ethyl)thymine-6-yl.

**EXAMPLE 154**

Synthesis of the 5LO sequence 5'- GAA GTC ACT GGA ACG  
containing 2-amino-4-oxo(3H)-8-(imidazol-4-yl[propyl]  
30 guanazolin-7-oxy at specific sites is described. A  
solution of 2-isobutryamide-4-diphenylaminocarbonyl-  
oxy-8-[1-(benzoyl)imidazol-4-ylpropylguanazolin-7-ol is  
treated with sodium hydride at room temperature for 1  
hour and then heated to 50°C for 10 min. This solution  
35 is treated with 3,5-di-O-p-methylbenzoyl- $\alpha$ -D-erythro-  
pentofuranosyl chloride. The solution is stirred at

- 149 -

ambient temperature for three hours and evaporated to dryness under reduced pressure. Silica gel chromatography (chloroform/methanol) is followed by selectively debenzoylation with methanolic ammonia at room temperature overnight. Recrystallization from ethanol provides 7-(2-deoxy- $\beta$ -D-erythro-pentofuranosyl)-8-(1-[benzoyl]imidazol-4-yl-propyl)-2-isobutryamide-4-diphenylamino carbonyloxypropylquanazolin-7-yl. This material is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'-b-cyanoethylphosphoramidite which is inserted into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

Modified sequences synthesized will be:

gAA GTC ACT GGA ACg  
15 gAA GTC ACT gGA ACg  
gAA gTC ACT GGA ACg  
GAA GTC ACT GGA ACg  
GAA GTC ACT ggA ACg  
GAA gTC ACT GGA ACg  
20 gAA GTC ACT GGA ACg  
GAA GTC ACT GgA ACg  
GAA gTC ACT GGA ACg and  
gAA gTC ACT ggA ACg  
where g is 2-amino-4-oxo(3H)-8-(imidazol-4-yl-[propyl])  
25 quanazolin-7-oxy.

#### EXAMPLE 135

Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 4-amino-7-(imidazol-4-yl)-ethoxyimidazo-[4,5-d]pyridazine at specific positions is described. A solution of 4,7-dichloroimidazo [4,5-d]pyridazine, Journal of Organic Chemistry, Vol. 23, p. 1534 (1958), in acetonitrile (100 ml) is treated with sodium hydride for one hour and then heated to 50°C for 10 min. This solution is treated with 3,5-di-O-(methylbenzoyl)- $\alpha$ -D-erythro-pentofuranosyl chloride. After stirring at ambient temperature for three hours the mixture is filtered. The filtrate is evaporated to dryness under



- 150 -

reduced pressure providing a foam which is chromatographed over silica gel. The pure product is aminated with liquid ammonia to provide 4-amino-7-chloro-1-(2-deoxy- $\beta$ -D-erythro-pentofuranosyl) imidazo [4,5-d]-pyridazine. Recrystallization of this material from ethanol/water provides pure deblocked nucleoside. This material is dissolved in DMF and treated with the disodium salt of 2-(imidazol-4-yl)ethanol. The solution is heated at 100°C for one hour and then evaporated to dryness under reduced pressure. The residue is recrystallized from water. After drying at 100°C (0.1 torr) for 2 hours, the material is transiently benzoylated on the 4-amino and 7-position imidazole ring as previously described. M. J. Gait, Oligonucleotide Synthesis, (IRL Press 1985). This material is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'- $\beta$ -cyanoethyl phosphoramidite which is inserted into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

Modified sequences synthesized will be:

GAA GTC ACT GGA aCG  
GaA GTC ACT GGA aCG  
GaA GTC aCT GGA aCG  
GAA GTC aCT GGA ACG  
25 GAA GTC ACT GGA aCG  
GaA GTC ACT GGA ACG  
GAA GTC ACT GGa ACG  
GAa GTC ACT GGA ACG and  
Gaa GTC aCT GGa aCG

30 where a is 4-amino-7-(imidazol-4-yl)-ethoxyimidazo [4,5-d] pyridazine.

#### EXAMPLE 156

Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 2'-deoxy-2'-(imidazol-4-yl)-ethoxyadenosine at specific positions is described. A solution of 3',5'-(1,1,3,3-tetraisopropylidisiloxanyl)-9-( $\beta$ -D-arabino-furanosyl) ad nine in pyridine is transiently benzoylated

- 151 -

by subsequent treatment with chlorotrimethylsilane, benzoyl chloride, and methanolic ammonia. The resulting solution is evaporated to dryness and recrystallized from ethanol. The pure product, dissolved in dry acetonitrile, is

5 treated with trifluoromethylsulfonic acid anhydride and sodium hydride at 0 °C. The solution is stirred at 0°C for one hour and then is allowed to warm to ambient temperature (one hour). The solution is evaporated to dryness and purified by silica gel chromatography

10 (methylene chloride/methanol). This material, N6-benzoyl-3',5'-(1,1,3,3-tetraisopropylidisiloxanyl)-2'-trifluoromethylsulfonyl-9-(β-D-arabinofuranosyl)adenine, serves as the starting material for the synthesis of adenosines substituted in the 2'-position with a variety

15 of functional groups and tethered functional groups. The β-D-arabinofuranosyl derivatives of guanine, cytosine, and thymine (Aldrich Chemical Co.) (as well as analog bases) protected such as by the N2-isobutryl, and N6-benzoyl, respectively, by conventional procedures can

20 be 2'-modified as just described. The 2'-modified ribonucleotides can be inserted into sequence-specific oligonucleotides as of examples 107 and 108 above. A solution of N6-benzoyl-3',5'-(1,1,3,3-tetraisopropyl disiloxanyl)-2'-trifluoromethylsulfonyl-9-(β-D-arabino-

25 furanosyl) adenine and acetonitrile is cooled to 0°C and treated with the disodium salt of 2-(imidazol-4-yl) ethanol in acetonitrile. The suspension is allowed to warm to room temperature and stirred at ambient temperature for two hours. The suspension is neutralized with

30 acetic acid, treated with tetrabutylammonium fluoride for 5 hours at room temperature, and evaporated to dryness under reduced pressure. The residue is dissolved in pyridine and transiently benzoylated as described above. After removal of the solvents, the residue is

35 purified by silica gel chromatography. The resulting N6-benzoyl-2'-deoxy-2'-(1-[benzoyl]imidazol-4-ylethoxy) adenosine is converted by procedures described above into its 5'-DMT-3'-β- cyanoethylphosphoramidite which is

- 152 -

inserted into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

Modified sequences synthesized will be:

- GAA GTC ACT GGA aCG  
 5 GAA GTC ACT GGA aCG  
 GAA GTC aCT GGA aCG  
 GAA GTC aCT GGA ACG  
 GAA GTC ACT GGA aCG  
 GAA GTC ACT GGA ACG  
 10 GAA GTC ACT GGa ACG  
 GAa GTC ACT GGA ACG and  
 Gaa GTC aCT GGa aCG

where a is 2'-deoxy-2'-(imidazo-4-yl)ethoxyadenosine.

#### EXAMPLE 157

- 15 Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 2'-deoxy-2'-(9-[3-chloro-6-methoxyacridinyl] amino-n-pentyloxy)adenosine at specific positions is described. A solution of N6-benzoyl-3',5'-(1,1,3,3-tetraisopropylidisiloxanyl)-2'-trifluoromethylsulfonyl-9-  
 20 ( $\beta$ -D-arabinofuranosyl)adenine, 3-chloro-6-methoxyacridinyl-9-amino-n-pentanol, and acetonitrile is cooled to 0 °C and treated with the sodium hydride. The solution is allowed to warm to room temperature, then stirred at ambient temperature for two hours, and treated with  
 25 tetrabutylammonium fluoride for two hours. The solution is evaporated to dryness and the residue purified by silica gel chromatography. This material is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'- $\beta$ -cyanoethyldiisopropylphosphoramidite derivative  
 30 which is inserted into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis. Modified sequences synthesized will be:
- GAA GTC ACT GGA aCG  
 GAA GTC ACT GGA aCG  
 35 GAA GTC aCT GGA aCG  
 GAA GTC aCT GGA ACG  
 GAA GTC ACT GGA aCG

- 153 -

Gaa GTC ACT GGA ACG

GAA GTC ACT GGa ACG

GAa GTC ACT GGA ACG and

Gaa GTC aCT GGa aCG

- 5 where a is 2'-deoxy-2'-(9-[3-chloro-6-methoxyacridinyl]  
amino-n-pentyloxy)adenosine.

**EXAMPLE 158**

- Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 2'-deoxy-2'-(N1,N1,N2-trimethoxycarbonyl-N2-methylcarbamoylethoxy)ethylenediamine adenosine at specific positions is described. A solution of N6-benzoyl-3',5'-(1,1,3,3-tetraisopropylidisiloxanyl)-2'-trifluoromethylsulfonyl-9-( $\beta$ -D-arabinofuranosyl)adenine and acetonitrile is cooled to 0°C and treated with the sodium salt of (N1,N1,N2-trimethoxycarbonyl-N2-methylcarbamoyl-ethanol) ethylenediamine in acetonitrile. The solution is allowed to warm to room temperature, then stirred at ambient temperature for two hours, and treated with tetrabutyl ammonium fluoride for two hours. The solution is evaporated to dryness and the residue purified by silica gel chromatography. This material is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'- $\beta$ -cyanoethylphosphoramidite derivative which is inserted into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis. Modified sequences synthesized will be:

GAa GTC ACT GGA aCG

GaA GTC ACT GGA aCG

GaA GTC aCT GGA aCG

- 30 GAA GTC aCT GGA ACG

GAA GTC ACT GGA aCG

GaA GTC ACT GGA ACG

GAA GTC ACT GGa ACG

GAa GTC ACT GGA ACG and

- 35 Gaa GTC aCT GGa aCG

where a is 2'-deoxy-2'-(N1,N1,N2-trimethoxycarbonyl-N2 methyl carbamoylethoxy)ethylenediamine adenosine.

**EXAMPLE 159**

Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 2'-deoxy-2'-nonanoxadenosine at specific positions is performed. A solution of N6-benzoyl-3',5'-(1,1,3,3-tetraisopropylidisiloxanyl)-2'-trifluoromethylsulfonyl-9-( $\beta$ -D-arabinofuranosyl)adenine and acetonitrile is cooled to 0°C and treated with the sodium salt of nananol in acetonitrile. The solution is allowed to warm to room temperature, then stirred at ambient temperature for two hours, and treated with tetrabutylammonium fluoride for two hours. The solution is evaporated to dryness and the residue purified by silica gel chromatography. This material is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'-b-cyanoethylidiisopropyl phosphoramidite derivative which is inserted into the 5LO- sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

Modified sequences synthesized will be:

GAA GTC ACT GGA aCG  
20 GaA GTC ACT GGA aCG  
GaA GTC aCT GGA aCG  
GAA GTC aCT GGA ACG  
GAA GTC ACT GGA aCG  
GaA GTC ACT GGA ACG  
25 GAA GTC ACT GGa ACG  
GAa GTC ACT GGA ACG and  
Gaa GTC aCT GGa aCG

where a is 2'-deoxy-2'-nonanoxadenosine.

**EXAMPLE 160**

30 Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 2'-deoxy-2'-dimethylamino-tris(ethylamino) adenosine at specific positions is described. A 0°C solution of N6-benzoyl-3',5'-(1,1,3,3-tetraisopropylidisiloxanyl)-2'-trifluoromethylsulfonyl-9-( $\beta$ -D-arabinofuranosyl)adenine, dimethylaminotris (ethylamine), ethyl isopropylamine, and acetonitrile is allowed to warm to

- 155 -

room temperature and stirred at ambient temperature for 3 hours and then treated with tetrabutylammonium fluoride for two hours. The solution is evaporated to dryness and the residue purified by silica gel chromatography. This material is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'- $\beta$ -cyanoethyl phosphoramidite derivative which is inserted into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

10 Modified sequences synthesized will be:

GAA GTC ACT GGA aCG  
GAA GTC ACT GGA aCG  
GAA GTC aCT GGA aCG  
GAA GTC aCT GGA ACG  
15 GAA GTC ACT GGA aCG  
GAA GTC ACT GGA ACG  
GAA GTC ACT GGA ACG  
GAA GTC ACT GGA ACG and  
Gaa GTC aCT GGA aCG

20 where a is 2'-deoxy-2'-dimethylamino-tris(ethylamino) adenosine.

#### EXAMPLE 161

Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 2'-deoxy-2'-(methoxytris-ethoxy)adenosine at specific position is described. A 0°C solution of N6-benzoyl-3',5'-(1,1,3,3-tetraisopropylidisiloxanyl)-2'-trifluoromethylsulfonyl-9-( $\beta$ -D-arabinofuranosyl)adenine, methoxybisethoxyethanol and acetonitrile is treated with sodium hydride and allowed to warm to room temperature, stirred at ambient temperature for 3 hours, and then treated with tetrabutylammonium fluoride for two hours. The solution is evaporated to dryness and the residue purified by silica gel chromatography. This material is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'- $\beta$ -cyanoethyldiisopropylphosphoramidite derivative which is inserted into the 5LO-sequence 5'-

- 156 -

GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

Modified sequences synthesized will be:

- GAA GTC ACT GGA aCG  
5 GAA GTC ACT GGA aCG  
GAA GTC aCT GGA aCG  
GAA GTC aCT GGA ACG  
GAA GTC ACT GGA aCG  
GAA GTC ACT GGA ACG  
10 GAA GTC ACT GGA ACG  
GAA GTC ACT GGA ACG and  
Gaa GTC aCT GGA aCG

where a is 2'-deoxy-2'-(methoxytris-ethoxy)adenosine.

#### EXAMPLE 162

- 15 Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 2'-deoxy-2'-(N1,N1,N2-trimethoxycarbonyl-N2-methyl carbamoylethoxy)ethylenediamine adenosine at specific positions is described. A solution of 2-chloro-2'-deoxyadenosine, Journal of the American Chemical  
20 Society, Vol. 106, p. 6379 (1984), DMF, and potassium thiomethylate is heated at 100°C for one hour, cooled to room temperature and treated with meta-chloroperbenzoate. After stirring at room temperature for three hours the solution is evaporated to dryness under reduced pressure.  
25 The residue is subjected to silica gel chromatography purification. The resulting 2-methylsulfonyl-2'-deoxyadenosine is benzoylated via the transient method as of examples 107 and 108 above. The N6-benzoyl-2'-deoxy-2-methylsulfonyladenosine thus prepared serves as the  
30 starting material for the preparation of a variety of 2'-deoxyadenosine and 2'-deoxyguanosine substituted in the 2-position. A solution of N6-benzoyl-2'-deoxy-2-methylsulfonyladenosine in DMF is treated with a solution of the sodium salt of (N1,N1,N2-trimethoxycarbonyl-N2-  
35 methylcarbamoylethanol)ethylenediamine in acetonitrile. The solution is heated to 100°C for 30 min and evaporated to dryness under reduced pressure. Recrystallization of

- 157 -

the residue affords N6-benzoyl-2'-deoxyadenosine with the ethylenediamine derivative in the 2-position. This material is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'- $\beta$ -cyanoethylphosphoramidite derivative which is inserted into the 5LO-sequence 5'-  
5 GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

Modified sequences synthesized will be:

GAA GTC ACT GGA aCG  
10 GaA GTC ACT GGA aCG  
GaA GTC aCT GGA aCG  
GAA GTC aCT GGA ACG  
GAA GTC ACT GGA aCG  
GaA GTC ACT GGA ACG  
15 GAA GTC ACT GGa ACG  
GAa GTC ACT GGA ACG and  
Gaa GTC aCT GGa aCG

where a is 2'-deoxy-2'-(N1,N1,N2-trimethoxycarbonyl-N2 methyl carbamoylethoxy)ethylenediamine adenosine.

#### 20 EXAMPLE 163

Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 2'-deoxy-2'-(9-[3-chloro-6-methoxyacridinyl] amino-n-pentylamino)adenosine at specific positions is performed. A solution of N6-benzoyl-2'-deoxy-2-methyl-sulfonyl-  
25 sulfonadenosine, ethyl diisopropylamine, 9-[3-chloro-6-methoxyacridinyl]amino-n-pentylamine, and DMF is heated to 100°C for 30 min and evaporated to dryness under reduced pressure. Recrystallization of the residue affords N6-benzoyl-2'-doxyadenosine with the amino-  
30 acridine derivative in the 2-position. This material is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'- $\beta$ -cyanoethylphosphor-amidite derivative which is inserted into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.  
35 Modified sequences synthesized will be:  
GAA GTC ACT GGA aCG  
GaA GTC ACT GGA ACG



- 158 -

GaA GTC aCT GGA aCG

GAA GTC aCT GGA ACG

GAA GTC ACT GGA aCG

GaA GTC ACT GGA ACG

5 GAA GTC ACT GGa ACG

GAa GTC ACT GGA ACG

Gaa GTC aCT GGa aCG

where a is 2'-deoxy-2-(9-[3-chloro-6-methoxyacridinyl] amino-n-pentylamino)adenosine.

10 **EXAMPLE 164**

Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 2-dimethylamino-tris(ethylamino)adenosine at specific positions is described. A solution of N6-benzoyl-2'-deoxy-2-methyl sulfonyladenosine, ethyl diisopropylamine, 2-dimethylamino- tris(ethylamine, and DMF is heated to 100°C for 30 min and evaporated to dryness under reduced pressure. Recrystallization of the residue affords N6-benzoyl-2'-doxyadenosine with the polyamine derivative in the 2-position. This material is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'-β-cyanoethyl diisopropylphosphoramidite derivative which is inserted into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis. Modified sequences synthesized will be:

25 GAa GTC ACT GGA aCG

GaA GTC ACT GGA aCG

GaA GTC aCT GGA aCG

GAA GTC aCT GGA ACG

GAA GTC ACT GGA aCG

30 GaA GTC ACT GGA ACG

GAA GTC ACT GGa ACG

GAa GTC ACT GGA ACG and

Gaa GTC aCT GGa aCG

where a is 2-dimethylamino-tris(ethylamino)adenine.

**EXAMPLE 165**

Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 2-(methoxybisethoxyethylamino)adenine at specific positions is described. A solution of N6-benzoyl-2'-deoxy-2-methyl sulfonyladenosine, ethyl diisopropylamine, methoxybisethoxy ethylamine, and DMF is heated to 100°C for 30 min and evaporated to dryness under reduced pressure. Recrystallization of the residue affords N6-benzoyl-2'-doxyadenosine with the ethylene-glycol derivative in the 2-position. This material is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'- $\beta$ -cyanoethyl phosphoramidite derivative which is inserted into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis. Modified sequences synthesized will be:

GAA GTC ACT GGA aCG  
GaA GTC ACT GGA aCG  
GaA GTC aCT GGA aCG  
GAA GTC aCT GGA ACG  
GAA GTC ACT GGA aCG  
GaA GTC ACT GGA ACG  
GAA GTC ACT GGa ACG  
GAa GTC ACT GGA ACG and  
Gaa GTC aCT GGa aCG

where a is 2-(methoxybisethoxyethylamino)adenine.

**EXAMPLE 166**

Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 2-octanoxymadenine at specific positions is described. A solution of N6-benzoyl-2'-deoxy-2-methyl-sulfonyladenosine, ethyl diisopropylamine, octanoylamine, and DMF is heated to 100°C for 30 min and evaporated to dryness under reduced pressure. Recrystallization of the residue affords N6-benzoyl-2'- deoxyadenosine with the carbon chain in the 2-position. This material is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'- $\beta$ -cyanoethyldiisopropylphosphoramidite derivative which is inserted into the 5LO-sequence 5'-

- 160 -

GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

Modified sequences synthesized will be:

- GAA GTC ACT GGA aCG  
5 GAA GTC ACT GGA aCG  
GAA GTC aCT GGA aCG  
GAA GTC aCT GGA ACG  
GAA GTC ACT GGA aCG  
GAA GTC ACT GGA ACG  
10 GAA GTC ACT GGA ACG  
GAA GTC ACT GGA ACG and  
Gaa GTC aCT GGA aCG  
where a is 2-octanoylaminoadenine.

#### EXAMPLE 167

- 15 Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 2'-deoxy-6'-(imidazol-4-yl-ethoxy)carbocyclic guanosine at specific positions is described. A solution of 2'-deoxy-3',5'- di-O-benzyl-6'-a-hydroxy-N6-methoxy-ethoxy carbocyclic guanosine, Journal of Chemical  
20 Society, Chemical Communications, London, pp. 1083-1084 (1987), is isobutrylated by the transient method (pyridine, chlorotrimethylsilane, then isobutryl chloride, then methanol/ammonia. This material is reacted with 2-(1-benzoyl imidazol-4-yl)ethanol,  
25 triphenylphosphine/DEAD, and acetonitrile at room temperature for three hours. The solution is evaporated to dryness under reduced pressure. The residue is recrystallized from ethanol providing product which is debenzylated and deblocked with palladium/charcoal,  
30 hydrogen, and hydrochloric acid to provide 2'-deoxy-6'- $\alpha$ -(imidazol-4-yl-ethoxy)-N2-isobutrylcarbocyclic guanosine. This material is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'- $\beta$ -cyano-ethylphosphoramidite derivative which is inserted into  
35 the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

Modified sequences synthesized will be:

- 161 -

- gAA GTC ACT GGA ACg  
gAA GTC ACT gGA ACg  
gAA gTC ACT GGA ACg  
GAA GTC ACT GGA ACg  
5 GAA GTC ACT ggA ACG  
GAA gTC ACT GGA ACG  
gAA GTC ACT GGA ACG  
GAA GTC ACT GgA ACG  
GAA gTC ACT GGA ACG and  
10 gAA gTC ACT ggA ACg  
where g is 2'-deoxy-6'-(imidazol-4-yl-ethoxy)carbocyclic  
guanosine.

**EXAMPLE 168**

- Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3'  
15 containing 2'-deoxy-6'- $\alpha$ -(N1,N1,N2-trimethoxycarbonyl-  
N2-methylcarbamoylethoxy)ethylenediamine) carbocyclic  
guanosine at specific positions is described. A solution  
of 2'-deoxy-3',5'-di-O-benzyl- 6'- $\alpha$ -hydroxy-N6-methoxy-  
ethoxy carbocyclic guanosine, Journal of Chemical  
20 Society, Chemical Communications, London, pp. 1083-1084  
(1987), is isobutrylated by the transient method  
pyridine/ chlorotrimethylsilane, then isobutryl chloride,  
then methanol/ammonia. The purified material is reacted  
with N1,N1,N2-trimethoxycarbonyl-N2-methylcarbamoyl-  
25 ethoxy) ethylenediamine diphenyldiazine, and acetonitrile  
at room temperature for three hours. The solution is  
evaporated to dryness under reduced pressure. The  
residue is recrystallized from ethanol providing product  
which is debenzylated and deblocked with palladium/  
30 charcoal, hydrogen, and hydrochloric acid to provide  
2'-deoxy-6'- $\alpha$ -(N1,N1,N2-trimethoxycarbonyl-N2-methylcar-  
bamoylethoxy)ethylenediamine)-N2-isobutrylcarbocyclic  
guanosine. This material is converted by procedures of  
examples 107 and 108 above into its 5'-DMT-3'- $\beta$ -cyano-  
35 ethylphosphoramidite derivative which is inserted into  
the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated,  
solid phase DNA synthesis.

- 162 -

Modified sequences synthesized will be:

- gAA GTC ACT GGA ACg  
gAA GTC ACT gGA ACg  
gAA gTC ACT GGA ACg  
5 GAA GTC ACT GGA ACg  
GAA GTC ACT ggA ACG  
GAA gTC ACT GGA ACG  
gAA GTC ACT GGA ACG  
GAA GTC ACT GgA ACG  
10 GAA gTC ACT GGA ACG and  
gAA gTC ACT ggA ACg

where g is 2'-deoxy-6'- $\alpha$ -(N1,N1,N2-trimethoxycarbonyl-N2-methylcarbamoylethoxy)ethylenediamine)carbocyclic guanosine.

15 **EXAMPLE 169**

- Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 2'-deoxy-2'-(imidazo-4-yl)ethoxy- $\alpha$ -D-adenosine at specific positions was performed. The 3',5'-positions of N6-Benzoyl- $\alpha$ -adenosine are simultaneously protected  
20 with 1,3-dichloro-1,1,3,3-tetraisopropyldisiloxane according to the known procedure. Nucleic Acid Chemistry, Improved and New Synthetic Procedures, Methods, and Techniques, Part 3, page 229 (1986). The 2'-hydroxy group is coupled with (1-benzoylimidazol-4-yl)ethanol  
25 with triphenylphosphine/DEAD according to the Mitsunobu procedure. Synthesis 1 (1981). The purified product is silyl deprotected with tetrabutylammonium fluoride and the resulting nucleoside is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'- $\beta$ - cyano-ethylphosphoramidite derivative which is inserted into  
30 the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

Modified sequences synthesized will be:

- GAA GTC ACT GGA aCG  
35 GaA GTC ACT GGA aCG  
GaA GTC aCT GGA aCG  
GAA GTC aCT GGA ACG

- 163 -

GAA GTC ACT GGA aCG  
GaA GTC ACT GGA ACG  
GAA GTC ACT GGa ACG  
GAa GTC ACT GGA ACG and

5 Gaa GTC aCT GGA aCG

where a is 2'-deoxy-2'-(imidazo-4-yl)ethoxy- $\alpha$ -D-adenosine.

#### EXAMPLE 170

Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 2'-deoxy-2'-(N1,N1,N2-trimethoxycarbonyl-N2-methylcarbamoylethoxyethylenediamine)- $\alpha$ -D-adenosine at specific positions was performed. A solution of N6-benzoyl-3',5'-(1,1,3,3-tetraisopropylidisiloxanyl)-2'-trifluoromethylsulfonyl-9-(7- $\alpha$ -D-arabinofuranosyl) adenine is coupled with (N1,N1,N2-trimethoxy carbonyl-N2-methylcarbamoylethanol) ethylenediamine with according to the Mitsunobu procedure. This material is deprotected with fluoride ion as of examples 107 and 108 above. The resultant material is converted by procedures described above into its 5'-DMT-3'- $\beta$ - cyanoethyldiisopropyl-phosphoramidite derivative which is inserted into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

Modified sequences synthesized will be:

25 GAa GTC ACT GGA aCG  
GaA GTC ACT GGA aCG  
GaA GTC aCT GGA aCG  
GAA GTC aCT GGA ACG  
GAA GTC ACT GGA aCG  
30 GaA GTC ACT GGA ACG  
GAA GTC ACT GGa ACG  
GAa GTC ACT GGA ACG and  
Gaa GTC aCT GGa aCG

where a is 2'-deoxy-2'-(N1,N1,N2-trimethoxycarbonyl-N2-methylcarbamoylethoxyethylenediamine)- $\alpha$ -D-adenosine.

- 164 -

**EXAMPLE 171**

Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 2'-deoxy-2'-(9-[3-chloro-6-methoxyacridinyl] amino-n-pentyloxy- $\alpha$ -D-erythropentofuranosyl)adenine at specific positions is performed. A solution of N6-benzoyl-3',5'-(1,1,3,3-tetraisopropylidisiloxanyl)-2'-trifluoromethylsulfonyl-9-( $\beta$ -D-arabinofuranosyl) adenine is coupled with chloro-6-methoxyacridinyl- 9-amino-n-pentanol according to the Mitsunobu procedure. The disilyl protection group is removed with fluoride ion and the resulting material is converted by procedures of examples 107 and 108 above into its 5'-DMT-3'- $\beta$ -cyano-ethyl-diisopropylphosphoramidite derivative which is inserted into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

Modified sequences synthesized will be:

GAA GTC ACT GGA aCG  
GaA GTC ACT GGA aCG  
GaA GTC aCT GGA aCG  
20 GAA GTC aCT GGA ACG  
GAA GTC ACT GGA aCG  
GaA GTC ACT GGA ACG  
GAA GTC ACT GGa ACG  
GAa GTC ACT GGA ACG and  
25 Gaa GTC aCT GGa aCG

where a is 2'-deoxy-2'-(9-[3-chloro-6-methoxyacridinyl] amino- n-pentyloxy)adenosine.

**EXAMPLE 172**

Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 2'-deoxy-2'-[dimethylamino-bis(ethylamino) ethoxy] - $\alpha$ -D-adenosine at specific positions was performed. A solution of N6-benzoyl-3',5'-(1,1,3,3-tetraisopropylidisiloxanyl)-2'-trifluoromethylsulfonyl-9-( $\beta$ -D-arabinofuranosyl)adenine is coupled with dimethyl-amino-bis(ethylamino)ethanol] according to the Mitsunobu procedure. The disilyl protection group is removed with fluoride ion and the resulting material is converted by

- 165 -

procedures of examples 107 and 108 above into its 5'-DMT-3'- $\beta$ -cyanoethylphosphoramidite derivative which was inserted into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

5 Modified sequences synthesized will be:

GAA GTC ACT GGA aCG

GaA GTC ACT GGA aCG

GaA GTC aCT GGA aCG

GAA GTC aCT GGA ACG

10 GAA GTC ACT GGA aCG

GaA GTC ACT GGA ACG

GAA GTC ACT GGa ACG

GAA GTC ACT GGA ACG and

Gaa GTC aCT GGa aCG

15 where a is 2'-deoxy-2'-[dimethylamino-bis(ethylamino)-ethoxy] - $\alpha$ -D-adenosine.

#### EXAMPLE 173

Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 1'-(imidazo-4-yl)ethoxymethyladenosine at  
20 specific positions was performed. Psicofuranine, Carbohydr. Res., Vol. 44, p. 112 (1975) is N6-benzoylated via the transient procedure as set forth by R. A. Jones in Oligonucleotide Synthesis -- A Practical Approach, M. J. Gait, Ed., (IRL Press, Washington, DC, 1985) and  
25 subsequently the 3',5'-hydroxyls groups are simultaneously protected with 1,3-dichloro-1,1,3,3-tetra-isopropylidisiloxane according to a known procedure. Nucleic Acid Chemistry, Improved and New Synthetic Procedures, Methods, and Techniques, Part 3, page 229,  
30 (1986). The 1'-hydroxymethyl group is treated with triphenylmethyl chloride in pyridine to afford protection of the primary alcohol. 2'-Deoxygenation is performed as described by M. J. Robins, et al, in J. Am. Chem. Soc., Vol. 105, p. 4059 (1983). The resulting 2'-deoxy nucleo-  
35 side is selectively detritylated with dilute trifluoroacetic acid. The 1'-hydroxymethyl group is coupled with 1-(benzoyl-imidazo-4-yl) ethanol according to the



- 166 -

Mitsunobu conditions. Removal of the silyl protecting group and subsequent conversion of the resulting nucleoside to the 5'-DMT and the 3'-cyanoethyldiisopropylphosphite according to the procedures of examples 107 and 5 108 will afford monomers suitable for insertion into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid phase DNA synthesis.

Modified sequences synthesized will be:

GAA GTC ACT GGA aCG  
10 GaA GTC ACT GGA aCG  
GaA GTC aCT GGA aCG  
GAA GTC aCT GGA ACG  
GAA GTC ACT GGA aCG  
GaA GTC ACT GGA ACG  
15 GAA GTC ACT GGa ACG  
GAa GTC ACT GGA ACG and  
Gaa GTC aCT GGa aCG

where a is 1'-(imidazo-4-yl)ethoxymethyladenosine.

#### EXAMPLE 174

20 Synthesis of 5LO sequence 5'- GAA GTC ACT GGA ACG -3' containing 4'-(imidazo-4-yl)ethoxymethylthymine at specific positions was preformed. 4'-Hydroxymethylthymidine, J. Org. Chem., Vol. 44, p. 1309 (1979) is selectively 4',5' protected with the TIPS group and then 25 converted to its 3'-THP derivative. This material is silyl deprotected with fluoride ion and subsequently selectively 5'-tritylated. The 3'-hydroxy group is coupled to 2-(1-benzoyl imidazo-4-yl)ethanol via the Mitsunobu reaction conditions. This material is treated 30 with acid to remove the trityl and THP groups and subsequently converted, as of examples 107 and 108 above, into the 5'-DMT and 3'-b-cyanoethoxydiisopropylphosphoramidite groups. This material is inserted into the 5LO-sequence 5'- GAA GTC ACT GGA ACG via automated, solid 35 phase DNA synthesis.

Modified sequences synthesized will be:

GAA GTC ACT GGA ACG

- 167 -







GAA GTC ACt GGA ACG and

GAA GtC ACt GGA ACG

where t = 4'-(imidazo-4-yl)ethoxymethylthymine

What is claimed is:

1. A composition represented by one of the Formulas 2, 3, 4, 5, 6 and 7:



where G and K are, independently, C or N;

J is N or CR<sub>3</sub>;

R<sub>1</sub> is OH or NH<sub>2</sub>;

- 169 -

$R_2$  and  $R_3$  are H,  $NH_2$ , lower alkyl, substituted lower alkyl, lower alkenyl, aralkyl, alkylamino, aralkylamino, substituted alkylamino, heterocycloalkyl, heterocyclo-alkylamino, aminoalkylamino, heterocycloalkylamino, poly-alkylamino, a RNA cleaving moiety, a group for improving the pharmacokinetic properties of an oligonucleotide or a group for improving the pharmacodynamic properties of an oligonucleotide;

10  $R_4$  and  $R_5$  are H, OH,  $NH_2$ , lower alkyl, substituted lower alkyl, substituted amino, a RNA cleaving moiety, a group for improving the pharmacokinetic properties of an oligonucleotide or a group for improving the pharmacodynamic properties of an oligonucleotide;

15  $R_6$  and  $R_7$  are H, OH,  $NH_2$ , SH, halogen,  $CONH_2$ ,  $C(NH)NH_2$ ,  $C(O)O$ -alkyl,  $CSNH_2$ , CN,  $C(NH)NHOH$ , lower alkyl, substituted lower alkyl, substituted amino, a RNA cleaving moiety, a group for improving the pharmacokinetic properties of an oligonucleotide or a group for improving the pharmacodynamic properties of an oligonucleotide;

20 X is a sugar or a sugar analog moiety where said sugar analog moiety is a sugar substituted with at least one substituent comprising a RNA cleaving moiety, a group for improving the pharmacodynamic properties of an oligonucleotide, or a group for improving the pharmacokinetic properties of an oligonucleotide;

provided that when said composition is represented by Formula 2 and when G is N and J is N then X is said sugar analog moiety substituted with at least one group other than F and  $OCH_3$ ;

further provided when said composition is represented by Formula 2 and G is N and J is  $CR_3$  and  $R_3$  is H then X is said sugar analog moiety;

- 170 -



further provided that when said composition is represented by Formula 4 and when G is N and X is said sugar,  $R_2$  is not H;

further provided that when said composition is represented by Formula 6 and  $R_6$  is H and  $R_2$  is  $\text{NH}_2$  and  $R_7$  is  $\text{CONH}_2$ ,  $\text{CSNH}_2$ ,  $\text{C(O)O-alkyl}$ ,  $\text{C(NH)NH}_2$  or  $\text{C(NH)NHOH}$  then X is said sugar analog moiety;

further provided that when said composition is represented by Formula 6 and  $R_6$  is H, OH or SH and  $R_7$  is  $\text{C(O)O-alkyl}$  or  $\text{C(NH)NH}_2$  and  $R_2$  is  $-\text{CH}_2\text{CN}$  then X is said sugar analog moiety; and

further provided that when said composition is represented by Formula 7 and  $R_3$  is H and G is C, X is said sugar analog moiety.

2. The composition of claim 1 wherein X is represented by one of the Formulas:



where Q is O or  $\text{CHR}_{11}$ ;

$R_8$  and  $R_9$  are H, lower alkyl, substituted lower alkyl, a RNA cleaving moiety, a group for improving the pharmacodynamic properties of an oligonucleotide or a group for improving the pharmacokinetic properties of an oligonucleotide;

$R_{10}$  is H, OH, lower alkyl, substituted lower alkyl, F, Cl, Br, CN,  $\text{CF}_3$ ,  $\text{OCF}_3$ ,  $\text{OCN}$ , O-alkyl, S-alkyl,  $\text{SOMe}$ ,  $\text{SO}_2\text{Me}$ ,  $\text{ONO}_2$ ,  $\text{NO}_2$ ,  $\text{N}_3$ ,  $\text{NH}_2$ , NH-alkyl,  $\text{OCH}_2\text{CH}=\text{CH}_2$ ,  $\text{OCH}=\text{CH}_2$ ,  $\text{OCH}_2\text{CCH}$ ,  $\text{OCCH}$ , aralkyl, heteroaralkyl, heterocycloalkyl, amino-alkylamino, heterocycloalkyl, polyalkylamino, substituted silyl, a RNA cleaving moiety, a group for improving the pharmacodynamic properties of an

- 171 -

oligonucleotide or a group for improving the pharmacokinetic properties of an oligonucleotide; and

- 5  $R_{11}$  is H, OH lower alkyl, substituted lower alkyl, a RNA cleaving moiety, a group for improving the pharmaco-kinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligo-nucleotide.

3. The composition of claim 1 wherein said RNA cleaving moiety comprises a portion reactive with said  
10 RNA.

4. The composition of claim 3 wherein the reactive portion comprises a basic functionality capable of catalyzing the hydrolysis of phosphodiester bonds in RNA.

- 15 5. The composition of claim 3 wherein said reactive portion comprises a coordination complex of a metal.

6. The composition of claim 3 wherein said reactive portion comprises a moiety capable of  
20 stabilizing the transition state formed in hydrolysis of a phosphodiester bond of RNA.

7. The composition of claim 6 wherein the moiety is capable of providing an octahedral coordination complex having two vacant sites.

- 25 8. The composition of claim 3 wherein said reactive portion comprises an acidic functionality capable of catalyzing the hydrolysis of phosphodiester bonds in RNA.

9. The composition of claim 3 wherein said  
30 reactive portion comprises a heterocycle.



- 172 -

10. The composition of claim 9 wherein the heterocycle is either a tetrazole or triazole.
11. The composition of claim 3 wherein said RNA cleaving moiety comprises an imidazole.
- 5 12. The composition of claim 3 wherein said reactive portion comprises an alkylating functionality.
13. The composition of claim 12 wherein the alkylating functionality is selected from the group consisting of sulfonyl halide, aziridines, nitrogen  
10 mustard compositions and Michael reaction accepting compositions.
14. The composition of claim 3 wherein said reactive portion comprises a functionality capable of forming free radicals.
- 15 15. The composition of claim 14 wherein the free radical functionality is selected from the group consisting of quinones, streptonigrins, mitomycins, coumarins, and active oxygen producing agents.
16. The composition of claim 1 wherein said RNA  
20 cleaving moiety further comprises a tether portion for attaching the reactive portion to the balance of the composition.
17. The composition of claim 1 wherein said RNA cleaving moiety comprises a nitrogen heterocycle.
- 25 18. The composition of claim 1 wherein said RNA cleaving moiety comprises an imidazole.
19. An oligonucleotide analog comprising at least one composition in accordance with claim 1.

20. A composition for modulating the activity of RNA comprising an oligonucleotide comprising at least one composition in accordance with claim 1.

21. A composition for modulating the activity of  
5 RNA comprising an oligonucleotide comprising at least one composition in accordance with claim 2.

22. A composition of represented by one of the Formulas:



wherein,

10 Q is O or CR<sub>11</sub>;

R<sub>8</sub> and R<sub>9</sub> are H, lower alkyl, substituted lower alkyl, a RNA cleaving moiety or a group for improving the pharmacodynamic properties of an oligonucleotide;

R<sub>10</sub> is H, OH, lower alkyl, substituted lower alkyl,  
15 F, Cl, Br, CN, CF<sub>3</sub>, OCF<sub>3</sub>, OCN, O-alkyl, S-alkyl, SOME, SO<sub>2</sub>Me, ONO<sub>2</sub>, NO<sub>2</sub>, N<sub>3</sub>, NH<sub>2</sub>, NH-alkyl, OCH<sub>2</sub>CH=CH<sub>2</sub>, OCH=CH<sub>2</sub>, OCH<sub>2</sub>CCH, OCCH, aralkyl, heteroaralkyl, heterocycloalkyl, poly-alkylamino, substituted silyl, a RNA cleaving moiety or a group for improving the pharmacodynamic  
20 properties of an oligonucleotide or a group for improving the pharmacokinetic properties of an oligonucleotide;

R<sub>11</sub> is H, OH, lower alkyl, substituted lower alkyl, a RNA cleaving moiety or a group for improving the  
25 pharmacodynamic properties of an oligonucleotide;

Z is a naturally-occurring or synthetic base moiety;  
and



- 174 -

provided that when Z = adenine, guanine, uracil, thymine or cytosine and R<sub>8</sub> and R<sub>9</sub> are H then R<sub>10</sub> is not F, Cl, Br, NH<sub>2</sub>, N<sub>3</sub>, O-CH<sub>3</sub>, O-CH<sub>2</sub>CH<sub>3</sub>, O-allyl or O-benzyl.

23. A composition of claim 22 wherein:

5 Z is a pyrimidinyl-1 or purinyl-9 moiety.



24. A composition of claim 22 wherein:

Z is a nucleoside or deoxynucleoside base.

25. A composition of claim 22 wherein:

10 B is adenine, guanine, uracil, thymine, cytosine or 5-methylcytosine.

26. A mixed sequenced oligonucleotide having at least one modified nucleotide incorporated therein wherein said modified nucleotide being represented by one of the Formulas:



15 wherein,

Q is O or CR<sub>11</sub>;

R<sub>8</sub> and R<sub>9</sub> are H, lower alkyl, substituted lower alkyl, a RNA cleaving moiety or a group for improving the pharmacodynamic properties of an oligonucleotide;

20

R<sub>10</sub>, is H, OH, lower alkyl, substituted lower alkyl, F, Cl, Br, CN, CF<sub>3</sub>, OCF<sub>3</sub>, OCN, O-alkyl, S-alkyl, SOME, SO<sub>2</sub>Me, ONO<sub>2</sub>, NO<sub>2</sub>, N<sub>3</sub>, NH<sub>2</sub>, NH-alkyl, OCH<sub>2</sub>CH=CH<sub>2</sub>, OCH=CH<sub>2</sub>, OCH<sub>2</sub>CCH, OCCH, aralkyl, heteroaralkyl,

25 heterocyclo-alkyl, poly-alkylamino, substituted silyl, a RNA cleaving moiety or a group for improving the

- 175 -

pharmacodynamic properties of an oligonucleotide or a group for improving the pharmaco-kinetic properties of an oligonucleotide;


5  $R_{11}$  is H, lower alkyl, substituted lower alkyl, a RNA cleaving moiety or a group for improving the pharmacodynamic properties of an oligonucleotide;

Z is a naturally-occurring or synthetic base moiety; and provided that if B is adenine, guanine, uracil, thymine or cytosine and both  $R_8$  and  $R_9$  are H, Q is O and  
10 all of the sugar linking groups are unsubstituted phosphodiester linkages, and if said oligonucleotide includes only one of said modified nucleotides then  $R_{10}$  is not O-CH<sub>3</sub> or F.

27. The oligonucleotide of claim 26 wherein at  
15 least some of the phosphodiester bonds of said oligonucleotide are substituted by phosphorothionate, methyl phosphonate, or alkylphosphate.


28. The oligonucleotide of claim 26 wherein  
at least some of the phosphodiester bonds of said  
20 oligonucleotide are substituted by substantially non-chiral, substantially non-ionic moieties.

29. A mixed sequenced oligonucleotide having at least one modified nucleotide incorporated therein wherein said modified nucleotide is represented by one of the Formulas 2, 3, 4, 5, 6 and 7:




5


2



3




4




5

- 177 -



6



7

where G and K are, independently, C or N;

J is N or CR<sub>2</sub>CR<sub>3</sub>

R<sub>1</sub> is OH or NH<sub>2</sub>;

- 5 R<sub>2</sub> and R<sub>3</sub> are H, NH<sub>2</sub>, lower alkyl, substituted lower alkyl, lower alkenyl, aralkyl, alkylamino, aralkylamino, substituted alkylamino, heterocycloalkyl, heterocyclo-alkylamino, aminoalkylamino, hetrocycloalkylamino, poly-alkylamino, a RNA cleaving  
10 moiety, a group for improving the pharmacokinetic properties of an oligonucleotide or a group for improving the pharmacodynamic properties of an oligonucleotide;

- R<sub>4</sub> and R<sub>5</sub> are H, OH, NH<sub>2</sub>, lower alkyl, substituted  
15 lower alkyl, substituted amino, a RNA cleaving moiety, a group for improving the pharmacokinetic properties of an oligonucleotide or a group for improving the pharmaco-dynamic properties of an oligonucleotide;

- R<sub>6</sub> and R<sub>7</sub> are H, OH, NH<sub>2</sub>, SH, halogen, CONH<sub>2</sub>,  
20 C(NH)NH<sub>2</sub>, C(O)O-alkyl, CSNH<sub>2</sub>, CN, C(NH)NHOH, lower alkyl, substi-tuted lower alkyl, substituted amino, a RNA cleaving moiety, a group for improving the pharmacokinetic proper-ties of an oligonucleotide or a group for improving the pharmacodynamic properties of  
25 an oligonucleotide;

X is a sugar or a sugar analog moiety where said sugar analog moiety is a sugar substituted with at

- 178 -

least one substituent comprising a RNA cleaving moiety, a group for improving the pharmacodynamic properties of an oligonucleotide, or a group for improving the pharmacokinetic properties of an

5 oligonucleotide;

provided that when said composition is represented by Formula 2 and when G is N and J is N then X is said sugar analog moiety substituted with at least one group other than F and  $\text{OCH}_3$ ;

10 further provided when said composition is represented by Formula 2 and G is N and J is  $\text{CR}_3$  and  $\text{R}_3$  is H then X is said sugar analog moiety;

further provided that when said composition is represented by Formula 4 and when G is N and X is said

15 sugar,  $\text{R}_2$  is not H;

further provided that when said composition is represented by Formula 6 and  $\text{R}_6$  is H and  $\text{R}_2$  is  $\text{NH}_2$  and  $\text{R}_7$  is  $\text{CONH}_2$ ,  $\text{CSNH}_2$ ,  $\text{C(O)O-alkyl}$ ,  $\text{C(NH)NH}_2$  or  $\text{C(NH)NHOH}$  then X is said sugar analog moiety;

20 further provided that when said composition is represented by Formula 6 and  $\text{R}_6$  is H, OH or SH and  $\text{R}_7$  is  $\text{C(O)O-alkyl}$  or  $\text{C(NH)NH}_2$  and  $\text{R}_2$  is  $-\text{CH}_2\text{CN}$  then X is said sugar analog moiety; and

further provided that when said composition is represented by Formula 7 and  $\text{R}_3$  is H and G is C, X is said sugar analog moiety;

25 further provided that if said oligonucleotide includes only one of said modified nucleotide and  $\text{G} = \text{N}$  and  $\text{J} = \text{N}$  of said modified nucleoside and all of the sugar linking groups are unsubstituted phosphodiester linkages then X is not a sugar or a sugar analog moiety containing a  $2' - \text{OCH}_3$  or  $2' - \text{F}$ .

30. The oligonucleotide of claim 29 wherein at least some of the phosphodiester bonds of said  
35 oligonucleotide are substituted by phosphorothionate, methyl phosphonate, or alkyl phosphate.

- 179 -

31. The oligonucleotide of claim 29 wherein at least some of the phosphodiester bonds of said oligonucleotide are substituted by substantially non-chiral, substantially non-ionic moieties.
- 5 32. A process for the preparation of 2'-substituted nucleosides comprising:  
treating a nucleoside having unblocked 2', 3' and 5' hydroxyl groups with metal hydride in a solvent; and  
10 reacting said treated nucleoside with a non-activated halide compound selected from the group consisting of alkyl halide, substituted alkyl halide, aralkyl halide and heterocycloalkyl halide.
33. A composition for modulating the activity of an RNA comprising:  
15 a targeting portion specifically hybridizable with a preselected nucleotide sequence of the RNA; and  
a moiety capable of cleaving a phosphodiester bond in said RNA.
- 20 34. The composition of claim 33 adapted for placement of said RNA cleaving moiety into the minor groove site of the hybrid structure formed from said RNA and said composition.
35. The composition of claim 33 wherein said  
25 targeting portion is an oligonucleotide or oligonucleotide analog comprising from about 3 to about 50 base units.
36. The composition of claim 33 wherein said  
30 targeting portion is an oligonucleotide or oligonucleotide analog comprising from about 8 to 40 base units.

- 180 -

37. The composition of claim 33 wherein said targeting portion is an oligonucleotide or oligonucleotide analog comprising from about 12 to 25 base units.
- 5 38. The composition of claim 33 wherein said targeting portion is an oligonucleotide analog wherein at least some phosphodiester bonds of an oligonucleotide have been substituted by phosphothionate bonds.
- 10 39. The composition of claim 33 wherein said targeting portion is an oligonucleotide analog wherein at least some phosphodiester bonds of an oligonucleotide have been substituted by substantially non-chiral, substantially non-ionic moieties.
- 15 40. The composition of claim 33 wherein said RNA cleaving moiety comprises a portion reactive with said RNA.
41. The composition of claim 33 wherein said RNA cleaving moiety further comprises a tether portion for  
20 attaching the reactive portion to the balance of the composition.
42. The composition of claim 33 wherein said RNA cleaving moiety comprises a basic functionality for catalyzing the hydrolysis of phosphodiester bonds in  
25 RNA.
43. The composition of claim 33 wherein said RNA cleaving moiety comprises a coordination complex of a metal
44. The composition of claim 33 wherein said RNA  
30 cleaving moiety is capable of stabilizing the

- 181 -

transition state formed in the hydrolysis of a phosphodiester bond of RNA.

45. The composition of claim 33 wherein said RNA cleaving moiety is capable of providing an octahedral  
5 coordination complex having two vacant sites.

46. The composition of claim 33 wherein said RNA cleaving moiety comprises an acidic functionality capable of catalyzing the hydrolysis of phosphodiester bonds in RNA.

10 47. The composition of claim 46 wherein said acidic functionality comprises a heterocycle.

48. The composition of claim 46 wherein said acidic functionality is a nitrogen heterocycle.

49. The composition of claim 47 wherein the  
15 heterocycle is a tetrazole or triazoles.

50. The composition of claim 33 wherein said RNA cleaving moiety comprises an alkylating functionality.

51. The composition of claim 33 wherein the alkylating functionality is selected from the group  
20 consisting of sulfonyl halides, aziridines, nitrogen mustard compositions and Michael reaction accepting compositions.

52. The composition of claim 33 wherein said RNA cleaving moiety comprises a functionality capable of  
25 forming free radicals.

53. The composition of claim 52 wherein the functionality is selected from the group consisting of



- 182 -

quinones, streptonigrins, mitomycins, coumarins, and active oxygen producing agents.

54. A composition for modulating the activity of RNA comprising:

5 a targeting portion specifically hybridizable with a preselected nucleotide sequence of the RNA; and

a reactive portion capable of covalently bonding with or cleaving said RNA;

10 the composition being adapted for placement of said reactive portion into the minor groove site of the hybrid structure formed from said RNA and said composition.

55. The composition of claim 54 wherein said  
15 reactive portion comprises an alkylating functionality.

56. The composition of claim 55 wherein the alkylating functionality is selected from the group consisting of sulfonyl halides, aziridines, nitrogen mustard compositions and Michael reaction accepting  
20 compositions.

57. The composition of claim 54 wherein said reactive portion comprises a functionality capable of forming free radicals.

58. The composition of claim 57 wherein the  
25 functionality is selected from the group consisting of quinones, streptonigrins, mitomycins, coumarins, and active oxygen producing agents.

59. A composition for modulating the activity of an RNA comprising:

- 183 -

a targeting portion specifically hybridizable  
with a preselected nucleotide sequence of the RNA;  
and

a portion capable of improving the  
5 pharmacodynamic or pharmacokinetic properties of  
said oligonucleotide;

the composition being adapted for placement of said  
improving portion into the minor groove site of the  
hybrid structure formed from said RNA and said  
10 composition.

60. A method for modulating the production of a  
protein by an organism comprising:

contacting said organism with a composition  
comprising:

15 a targeting portion specifically hybridizable  
with a preselected nucleotide sequence of the RNA;  
and

a reactive portion capable of catalyzing the  
cleavage of a phosphodiester bond in said RNA;

20 61. The method of claim 60 wherein said composition  
is adapted for placement of said reactive portion into  
the minor groove site of the hybrid structure formed  
from said RNA and said composition.

62. The method of claim 60 wherein said targeting  
25 portion is an oligonucleotide or oligonucleotide analog  
comprising from about 3 to about 50 base units.

63. The method of claim 60 wherein said targeting  
portion is an oligonucleotide or oligonucleotide analog  
comprising from about 8 to 40 base units.

30 64. The method of claim 60 wherein said targeting  
portion is an oligonucleotide or oligonucleotide analog  
comprising from about 12 to 25 base units.

65. The method of claim 60 wherein said targeting portion is an oligonucleotide analog wherein at least some phosphodiester bonds of said oligonucleotide have been substituted by phosphorothionate.
- 5 66. The method of claim 60 wherein said targeting portion is an oligonucleotide analog wherein at least some phosphodiester bonds of said oligonucleotide have been substituted by substantially non-chiral, substantially non-ionic moieties.
- 10 67. The method of claim 60 wherein said reactive portion comprises a basic functionality capable of catalyzing the hydrolysis of phosphodiester bonds in RNA.
68. The method of claim 60 wherein said reactive  
15 portion comprises a coordination complex of a metal.
69. The method of claim 60 wherein said reactive portion comprises a moiety capable of stabilizing the transition state formed in hydrolysis of a phosphodiester bond in RNA.
- 20 70. The method of claim 69 where in the moiety is capable of providing an octahedral coordination complex having two vacant sites.
71. The method of claim 60 wherein said reactive portion comprises an acidic functionality capable of  
25 catalyzing the hydrolysis of phosphodiester bonds in RNA.
72. The method of claim 71 wherein said acidic functionality comprises a heterocycle.

- 185 -

73. The method of claim 71 wherein said acidic functionality is a nitrogen heterocycle.

74. The method of claim 73 wherein the heterocycle is either a tetrazole or a triazoles.

5 75. The method of claim 60 wherein said reactive portion comprises an alkylating functionality.

76. The method of claim 75 wherein the alkylating functionality is selected from the group consisting of sulfonyl halide, aziridines, nitrogen mustards and  
10 Michael reaction accepting moieties.

77. The method of claim 60 wherein said reactive portion comprises a functionality capable of forming free radicals.

78. The method of claim 77 wherein the  
15 functionality is selected from the group consisting of quinones, streptonigrins, mitomycins, coumarins, and active oxygen producing agents.

79. A method for modulating the production of a protein by an organism comprising:

20 contacting said organism with a composition comprising:

a targeting portion specifically hybridizable with a preselected nucleotide sequence of the RNA; and

25 a reactive portion capable of covalently bonding with or cleaving said RNA;

the composition being adapted for placement of said reactive portion into the minor groove site of the  
30 hybrid structure formed from said RNA and said composition.

- 186 -

80. The method of claim 79 further comprising  
preselecting said nucleotide sequence of the RNA.

81. A method of treating an animal having a disease  
characterized by undesired production of a protein  
5 comprising:

contacting the animal with a composition  
comprising:

a targeting portion specifically  
hybridizable with a preselected nucleotide  
10 sequence of the RNA; and  
a reactive portion capable of covalently  
bonding with or cleaving said RNA.

82. A method of treating an animal having a disease  
characterized by undesired production of a protein  
15 comprising:

contacting the animal with a composition  
comprising:

a targeting portion specifically  
hybridizable with a preselected nucleotide  
20 sequence of the RNA; and  
a reactive portion capable of covalently  
bonding with or cleaving said RNA;

the composition being adapted for placement of said  
reactive portion into the minor groove site of the  
25 hybrid structure formed from said RNA and said  
composition.

83. A method for detecting the presence or absence  
of an RNA in a sample suspected of containing it  
comprising

30 contacting the sample with a composition comprising:  
a targeting portion specifically hybridizable  
with a preselected nucleotide sequence of the RNA;  
and

- 187 -

a reactive portion capable of covalently bonding with or cleaving said RNA; the composition being adapted for placement of said reactive portion into the minor groove site of the hybrid structure formed from said RNA and said composition; and detecting cleavage of the RNA by the composition.

84. A nuclease resistant oligonucleotide or oligonucleotide analog for modulating the activity of a selected sequence of RNA or DNA having a sequence of nucleotide bases specifically hybridizable with said selected sequence and having at least one modified 2'-deoxyfuranosyl moiety.

85. The oligonucleotide or oligonucleotide analog of claim 84 wherein said modification comprises substitution by H, OH, halo, azido, amino, substituted amino, cyano, halomethyl, isocyanato, alkoxyl, thioalkoxyl, halo-alkoxyl, alkyl sulfide, alkyl sulfonate, nitrate, ammonium, allyloxy or alkeneoxy.

86. The oligonucleotide or oligonucleotide analog of claim 84 wherein said modification comprises substitution by H, OH, halo, azido, amino, allyloxy, methoxy, or alkyl.

87. The oligonucleotide or oligonucleotide analog of claim 84 having from about 5 to about 50 nucleotide bases.

88. The oligonucleotide or oligonucleotide analog of claim 84 wherein said modification is at the 3' end of said oligonucleotide.

- 188 -

89. The oligonucleotide or oligonucleotide analog of claim 84 wherein at least one of the sugar linking groups is replaced with carbon or ether linkages.
90. The oligonucleotide or oligonucleotide analog  
5 of claim 84 wherein at least one of the nucleotides of said oligonucleotide has 5'-methylene group or a sugar 4'-oxygen removed.
91. The oligonucleotide or oligonucleotide analog of claim 84 further modified such that at least some of  
10 the sugar linking groups comprise a phosphorothioate, methyl phosphonate, or phosphate alkylate.
92. The oligonucleotide or oligonucleotide analog of claim 84 in a pharmaceutically acceptable carrier.
93. The oligonucleotide or oligonucleotide analog  
15 of claim 84 wherein said selected sequence of RNA or DNA comprises a portion of a HIV genome.
94. The oligonucleotide or oligonucleotide analog of claim 84 wherein said selected sequence of RNA or DNA comprises a portion of a herpes virus genome.
- 20 95. The oligonucleotide or oligonucleotide analog of claim 84 wherein said selected sequence of RNA or DNA comprises a portion of a papilloma virus genome.
96. A method for modulating the production of a protein by an organism comprising contacting the  
25 organism with an oligonucleotide or oligonucleotide analog having a sequence of nucleotide bases specifically hybridizable with a selected sequence of RNA or DNA coding for said protein, and having at least one modified nucleoside moiety.

- 189 -

97. The method of claim 96 wherein said modified nucleoside moiety is a 2'-modified-2'-deoxyribofuranosyl moiety.

98. The method of claim 96 wherein said  
5 modification comprises substitution by H, OH, halo, azido, amino, sub-stituted amino, cyano, halomethyl, isocyanato, alkoxyl, thioalkoxyl, haloalkoxyl, alkyl sulfide, alkyl sulfonate, nitrite, ammonium, allyloxy, or alkeneoxy.

10 99. The method of claim 96 wherein said modification comprises substitution by hydrogen of a hydroxide, halo, azido, amino, methoxy, allyloxy or alkyl group.

100. The method of claim 96 wherein said  
15 oligonucleotide has from about 5 to about 50 nucleotide bases.

101. The method of claim 96 wherein said modification is at the 3' end of said oligonucleotide.

102. The method of claim 96 wherein at least one of  
20 the sugar linking groups of said modified oligonucleotide is replaced with a carbon or ether linkage.

103. The method of claim 96 wherein at least one of the nucleotides of said oligonucleotide has 5'-  
25 methylene group or a sugar 4'-oxygen removed.

104. The method of claim 96 wherein the oligonucleotide is further modified such that at least some of the sugar linking groups of said oligonucleotide are modified to comprise a



- 190 -

phosphorothioate, methyl phosphonate, or phosphate alkylate.

105. The method of claim 96 wherein the oligonucleotide is in a pharmaceutically acceptable carrier.

5 106. The method of claim 96 wherein said selected sequence of RNA or DNA comprises a portion of a HIV genome.

107. The method of claim 96 wherein said selected sequence of RNA or DNA comprises a portion of a herpes  
10 virus genome.

108. The method of claim 96 wherein said selected sequence of RNA or DNA comprises a portion of a papilloma virus genome.

109. A method for treating an organism having a  
15 disease characterized by the undesired production of a protein comprising contacting the organism with a nuclease resistant oligonucleotide or oligonucleotide analog having a sequence of bases specifically hybridizable with a RNA or DNA sequence coding for said  
20 protein, wherein at least one of the nucleoside moieties is modified, either alone or in combination with a pharmaceutically acceptable carrier.

110. The method of claim 109 wherein said modification comprises substitution by H, OH, halo,  
25 azido, amino, substituted amino, cyano, halomethyl, isocyanato, alkoxyl, thioalkoxyl, haloalkoxyl, alkyl sulfide, alkyl sulfonate, nitrate, nitrite, ammonium, allyloxy or alkeneoxy.

- 191 -

111. The method of claim 109 wherein said modification comprises substitution by H, OH, halo, azido, amino, methoxy, allyloxy or alkyl.
112. The method of claim 109 wherein said modified  
5 nucleoside moiety is a 2'-modified-2'-deoxyribofuranosyl moiety.
113. The method of claim 109 wherein said oligonucleotide has from about 5 to about 50 nucleotide bases.
- 10 114. The method of claim 109 wherein said modification is at the 3' end of said oligonucleotide.
115. The method of claim 109 wherein at least one of the sugar linking groups of said modified oligonucleotide is replaced with a carbon or ether  
15 linkage.
116. The method of claim 115 wherein at least one of the nucleotides of said oligonucleotide has 5'-methylene group or a sugar 4'-oxygen removed.
117. The method of claim 109 wherein at least some  
20 of the sugar linking groups of said modified oligonucleotide are modified to comprise a phosphorothioate, methyl phosphonate, or phosphate alkylate.
118. The method of claim 109 wherein said  
25 oligonucleotide is in a pharmaceutically acceptable carrier.
119. The method of claim 109 wherein said sequence of RNA or DNA comprises a portion of a HIV genome.

- 192 -

120. The method of claim 109 wherein said sequence of RNA or DNA comprises a portion of a herpes virus genome.

121. The method of claim 109 wherein said sequence  
5 of RNA or DNA comprises a portion of a papilloma virus genome.

# INTERNATIONAL SEARCH REPORT

International Application No. PCT/US91/00243

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (If several classification symbols apply, indicate all) <sup>6</sup>		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC(5): C07H 1/00; A61K 31/70; C12Q 1/68		
U.S. CL.: 536/23, 29; 514/44; 435/6		
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched <sup>7</sup>		
Classification System	Classification Symbols	
US.	514/44; 536/23, 29; 435/6	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched <sup>8</sup>		
Databases Searched: APS, CAS ONLINE		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT <sup>9</sup></b>		
Category <sup>*</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
Y	US, A, 4,876,335 (Yamane et al.) 24 October 1989, see entire document.	1-59, 83-95
Y,P	US, A, 4,965,350 (Inoue et al.) 23 October 1990, see entire document.	1-59, 83-95
Y	US, A, 4,760,017 (McCormick) 26 July 1988, see entire document.	83-95
Y	US, A, 4,689,320 (Kaji) 25 August 1987, see entire document	60-82, 96-121
Y	US, A, 4,511,713 (Miller et al.) 16 April 1985, see entire document	60-82, 96-121
<p><sup>*</sup> Special categories of cited documents: <sup>10</sup></p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"A" document member of the same patent family</p>		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
11 March 1991		07 MAY 1991
International Searching Authority		Signature of Authorizing Officer
ISA/US		James O. Wilson